

REGULARITY OF ABSOLUTELY CONTINUOUS INVARIANT MEASURES FOR PIECEWISE EXPANDING UNIMODAL MAPS.

FABIÁN CONTRERAS, DMITRY DOLGOPYAT

CONTENTS

1. Introduction	1
2. Preliminary	3
2.1. Piecewise Expanding Unimodal Maps	3
2.2. Auxiliary facts and Transfer Operators	3
3. Repeated Derivatives of the density function	5
3.1. Explicit formulas for the first and the second derivatives.	5
3.2. Higher order derivatives.	8
4. Differentiability set for the density.	13
4.1. Saltus part.	13
4.2. Absolute continuity.	15
4.3. Differentiability points.	16
4.4. Whitney smoothness	18
4.5. Nondifferentiability set.	19
References	20

1. INTRODUCTION

An important discovery of the 20th century mathematics is that many deterministic systems exhibit stochastic behavior. The stochasticity is caused by exponential divergence of nearby trajectories. This instability causes many important objects associated to dynamical systems, such as attractors and invariant measures, to be fractal.

Piecewise expanding maps of the interval are among the simplest and most studied examples of chaotic systems. They admit absolutely continuous invariant measures (a.c.i.m) [10] which are ergodic, mixing, enjoy exponential decay of correlations and the Central Limit Theorem for Hölder observables (see e.g. [1, 19]). In this paper, we consider a class of simplest piecewise expanding maps, so called piecewise expanding unimodal maps (PEUMs)¹ of the unit interval. PEUMs are piecewise expanding maps with only two branches. We study regularity of the density of a.c.i.m for PEUMs. A classical result of A. Lasota and J. Yorke [10] says that the density, which we denote by ρ , is of bounded variation. Recall that a bounded variation function is differentiable almost everywhere (See e.g., [6], Corollary 6.6). Therefore the set of non-differentiability of ρ is a natural

¹The precise definition of PEUMs is given at the beginning of Section 2.

fractal set associated to our PEUM. Let us describe the previous results about the differentiability. In the smooth case, R. Sacksteder [15] and K. Krzyzewski [9] proved that when map f is expanding of class C^r then ρ is of class C^{r-1} . Later, B. Szewc [18] showed that if f is a piecewise expanding continuous map with finitely many critical points (those points where the derivative of f is not defined) then a density function will belong to the space

$$\{\phi \in BV[0, 1] : \phi \in C^{r-1} \text{ in } [0, 1] \setminus B\},$$

where B is the union of the closures of the critical orbits. In this paper, we improve on $r = 1$ case of the Szewc's theorem for PEUMs by showing that the set where ρ is differentiable is larger. Namely, we need to discard not all points in the closure of the critical orbit, but only points which are approached by the critical orbits exponentially fast. We also obtain a partial converse, by showing that if x is approached exponentially fast by the critical orbit *and the exponent is sufficiently large* then ρ is not differentiable at x .

We also show that a similar improvement is possible for $r > 1$ if we consider smoothness in the sense of Whitney, that is, we study the points where of the density admits a Taylor expansion of order r . (Of course Szewc's result is optimal for classical smoothness since the the set where the density is not differentiable is dense in B). This leads to the question of describing the Taylor coefficients of the density. Here we make use of the recent result of V. Baladi [2]² saying that the density ρ belongs to the set

$$BV_1 = \{\phi \in BV[0, 1] : \phi' \in BV[0, 1]\}.$$

In other words, the derivative of ρ coincides with a function of bounded variation almost everywhere. Accordingly, we can differentiate that function almost everywhere and call the result the second derivative of ρ . We then show that this procedure can be continued recursively and that the resulting functions indeed provide the Taylor coefficients of ρ .

More precisely, the main results of our paper can be summarized as follows. Let f be a PEUM such that both branches of f are C^{k+2} .

Theorem 1.1. *There is a sequence of functions $\rho_0, \rho_1, \dots, \rho_k \in BV$ such that $\rho_0 = \rho$ and for $j < k, \rho'_j = \rho_{j+1}$ almost everywhere.*

Theorem 1.2. (A) *The set of points where ρ is non differentiable has Hausdorff dimension zero.*

(B) *If the critical orbit is dense then the set of points where ρ is non differentiable is uncountable.*

(C) *There is a set \mathcal{N}_k such that $\mathcal{HD}(\mathcal{N}_k) = 0$ and ρ is k differentiable in the sense of Whitney on $[0, 1] - \mathcal{N}_k$. That is, if $\bar{x} \notin \mathcal{N}_k$ then*

$$\rho(x) - \rho(\bar{x}) = \sum_{m=1}^k \frac{\rho_m(\bar{x})}{m!} (x - \bar{x})^m + o\left((x - \bar{x})^k\right).$$

²Baladi was motivated by the question of regularity of invariant measure with respect to parameters raised in the work of D. Ruelle [12, 13, 14]. Applications of Baladi's result to Ruelle's question are described in [2, 3, 4]. Our results also have applications to the regularity question as will be detailed elsewhere.

Note that since $[0, 1] - \mathcal{N}_k$ is not closed, ρ in general *can not* be extended to a smooth function on $[0, 1]$.

Remark 1.3. *The set \mathcal{N}_k is typically much smaller than the set B used in [18]. Indeed, if f_t is a family of PEUMs satisfying a certain transversality condition then $B(f_t)$ contains an interval for almost all t (see e.g. [16, 17]).*

The paper is organized as follows:

In Section 2, we give the necessary definitions. In particular, we introduce a special family of transfer operators used in the proof of Theorem 1.1. We then prove several auxiliary facts of independent interest.

Section 3 starts with some explicit formulas for the first and second derivatives³ of ρ which are proven to belong to $BV[0, 1]$. Then we extend our analysis to repeated differentiation of arbitrary order proving Theorem 1.1.

Section 4 begins with some results on the regularity of the saltus part of ρ . Then we show that the regular part of ρ is not only continuous but also absolutely continuous. In the remaining subsections we prove Theorem 1.2. That is we show that ρ admits a Taylor expansion after we remove an exceptional set of zero Hausdorff dimension.

2. PRELIMINARY

2.1. Piecewise Expanding Unimodal Maps. We work with piecewise expanding unimodal maps. $f : [0, 1] \rightarrow [0, 1]$ is a piecewise expanding unimodal map (PEUM) if there is a point c called the critical point, a number $\varepsilon > 0$ and a constant $\lambda > 1$ such that

$$(1) \quad f(x) = \begin{cases} f_1(x) & \text{if } x \leq c \\ f_2(x) & \text{if } x \geq c \end{cases}$$

where f_1 is a C^2 map defined on $[0, c + \varepsilon]$ and f_2 is a C^2 map defined on $[c - \varepsilon, 1]$ such that $f_1(c) = f_2(c)$ and $|Df_j(x)| \geq \lambda$ for all x from the domain of f_j .

PEUMs have unique a.c.i.m. [10] which is ergodic (in fact it is mixing and even exponentially mixing [1, 19]). Let us denote by ρ the density of the a.c.i.m. ρ is a function of bounded variation.

From now on, λ will mean $\lambda := \inf_{x \neq c} |Df(x)|$.

2.2. Auxiliary facts and Transfer Operators. Denote by $\xi(z) = \frac{D^2f(z)}{Df(z)}$. In the arguments of this section we will need to represent $D(|Df^m y|)$ as a sum. Namely we have

$$D(|Df^m y|) = \frac{|Df^m y|}{Df^m y} \sum_{j=0}^{m-1} \xi(f^j y) Df^j y \quad \text{and} \quad D(Df^m y) = \sum_{j=0}^{m-1} \xi(f^j y) Df^j y.$$

Both formulas are easy consequences of the chain rule.

We need to introduce a family of transfer operators acting on the space $BV[0, 1]$ of functions of bounded variation. $BV[0, 1]$ it is a Banach space with the norm $\|\cdot\|_{BV} = \|\cdot\|_\infty + \text{var}(\cdot)$, where $\|\cdot\|_\infty$ is the usual supremum norm and $\text{var}(\cdot)$ is the total variation (cf. [6], page 116).

³the derivatives are understood in the sense of Theorem 1.1

The first operator in our family is $\tilde{\mathcal{L}}(\phi)(x) = \sum_{f(y)=x} \frac{\phi(y)}{Df(y)|Df(y)|}$. Note that $\tilde{\mathcal{L}}$ preserves $BV[0, 1]$. More generally, we shall use the following transfer operators acting on $BV[0, 1]$.

Definition 2.1. For $\phi \in BV[0, 1]$, define the operator $\tilde{\mathcal{L}}^m(\phi)$ by

$$\tilde{\mathcal{L}}^m(\phi)(x) = \sum_{f(y)=x} \frac{\phi(y)}{(Df(y))^m |Df(y)|},$$

where m is a nonnegative integer.

Note that $\tilde{\mathcal{L}}^1 = \tilde{\mathcal{L}}$.

Definition 2.2. Let k, i_1, \dots, i_k and $m_1 > \dots > m_k$ be positive integers. For functions h_1, \dots, h_k , define $\mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}$ at (h_1, \dots, h_k) by

$$\mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k) = \left(\tilde{\mathcal{L}}^{m_1} \right)^{i_1} \left(h_1 \left(\tilde{\mathcal{L}}^{m_2} \right)^{i_2} \left(h_2 \cdots \left(\tilde{\mathcal{L}}^{m_k} \right)^{i_k} (h_k) \right) \right).$$

We shall often use

Proposition 2.3.

$$\|\mathfrak{D}_{m_1, m_2, \dots, m_k}^{i_1, i_2, \dots, i_k}(h_1, \dots, h_k)\| \leq M(\lambda^{-i_1})^{m_1} (\lambda^{-i_2})^{m_2} \dots (\lambda^{-i_k})^{m_k}$$

Proof. We use induction on k . For $k = 1$ we have

$$\begin{aligned} |\tilde{\mathcal{L}}^m(h)(x)| &= \left| \sum_{f^i y=x} \frac{h}{(Df^i(y))^m |Df^i(y)|} \right| \leq \sum_{f^i y=x} \left| \frac{|h|}{|Df^i(y)|^m |DF^i(y)|} \right| \\ &\leq \frac{|h|}{\lambda^{im}} |\mathcal{L}^i(1)(x)| \leq \frac{C\|h\|}{\lambda^{im}}. \end{aligned}$$

Now, let us suppose the result is true for $k - 1$, we have

$$\begin{aligned} |\mathfrak{D}_{m, m_2, \dots, m_k}^{i, i_2, \dots, i_k}(h_1, \dots, h_k)(x)| &= \left| \left(\tilde{\mathcal{L}}^m \right)^i (h_1 \mathfrak{D}_{m_2, \dots, m_k}^{i_2, \dots, i_k}(h_2, \dots, h_k)) \right| \\ &\leq \left\| \sum_{f^i y=x} \frac{h_1(y) \mathfrak{D}_{m_2, \dots, m_k}^{i_2, \dots, i_k}(h_2, \dots, h_k)(y)}{(Df^i(y))^m |Df^i(y)|} \right\| \leq \lambda^{-im} \sum_{f^i y=x} \frac{\|h_1 \mathfrak{D}_{m_2, \dots, m_k}^{i_2, \dots, i_k}(h_2, \dots, h_k)\|}{|Df^i(y)|} \\ &\leq \lambda^{-im} \|h_1 \mathfrak{D}_{m_2, \dots, m_k}^{i_2, \dots, i_k}(h_2, \dots, h_k)\| \|\mathcal{L}^i(1)\| \leq \tilde{M}(\lambda^{-i})^m \|\mathfrak{D}_{m_2, \dots, m_k}^{i_2, \dots, i_k}(h_2, \dots, h_k)\| \\ &\leq \tilde{K}(\lambda^{-i})^m \tilde{M}(\lambda^{-i_2})^{m_2} \dots (\lambda^{-i_s})^{m_s} \leq M(\lambda^{-i})^m (\lambda^{-i_2})^{m_2} \dots (\lambda^{-i_s})^{m_s} \end{aligned}$$

Thus the claim holds for k and the Proposition is proven by induction. \square

If a series consisting of functions in $BV[0, 1]$ converges to a function g , then the series of the derivatives of each term does not always converge to the derivative of g . However, assuming that the series of derivatives converges in L^1 we have the following result.

Lemma 2.4. *If $\sum_{k=1}^n g_k \rightarrow g$ in BV and $\sum_{k=1}^n g'_k \rightarrow h$ in L_1 — then $g' = h$ a. e.*

Proof. Let $\epsilon > 0$. Then, there exists $N > 0$ such that, for all $n \geq N$,

$$\|g - \sum_{k=1}^n g_k\|_{BV} \leq \epsilon$$

Since $\|f'\|_{L_1} \leq \|f\|_{BV}$ for any function $f \in BV$, then

$$\|g' - \sum_{k=1}^n g'_k\|_{L_1} \leq \|g - \sum_{k=1}^n g_k\|_{BV} \leq \epsilon$$

Therefore, $\sum_{k=1}^n g'_k$ converges to g' in L_1 , hence $g' = h$ as claimed. \square

Another simple but useful fact is the following.

Lemma 2.5. *Let $g(s, r)$ be a function from $(\mathbb{Z} \cup \{0\}) \times (\mathbb{Z} \cup \{0\})$ to \mathbb{R} . Suppose the*

series $\sum_{i=1}^{\infty} \sum_{j=0}^{i-1} g(i-j, j)$ converges absolutely. Then,

$$\sum_{i=1}^{\infty} \sum_{j=0}^{i-1} g(i-j, j) = \sum_{c=1}^{\infty} \sum_{d=0}^{\infty} g(c, d)$$

We leave the proof to the reader.

3. REPEATED DERIVATIVES OF THE DENSITY FUNCTION

3.1. Explicit formulas for the first and the second derivatives. Before analyzing repeated derivatives of ρ of arbitrary order, we will start by giving explicit formulas for ρ' and ρ'' .

Let us define

$$\rho_1 = - \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i(\xi \cdot \rho)$$

Note that the series converges (absolutely) by Proposition 2.3 since ρ and ξ belong to $BV[0, 1]$.

Lemma 3.1. (a) *Let ρ be the density of the invariant measure of f . Then, $\rho' = \rho_1$ almost everywhere.*

(b) *$(\mathcal{L}^n \rho)'(x)$ converges to $\rho_1(x)$ uniformly for x which are not on the orbit of c .*

Proof. Since ρ is a fixed point of \mathcal{L} , then $\rho = \mathcal{L}^n(\rho)$ for all n . Because ρ is of bounded variation so is $\mathcal{L}^n(\rho)$, hence both are differentiable almost everywhere. In fact, differentiating both sides, we get $\rho' = (\mathcal{L}^n \rho)'$ almost everywhere. Next if $h \in BV$ then

$$(\mathcal{L}^n h)'(x) = \sum_{f^n y=x} \frac{h'(y)}{Df^n(y)|Df^n(y)|} - \sum_{f^n(y)=x} \frac{h(y) \cdot D(|Df^n(y)|)}{|Df^n(y)|^2} \quad \text{a. e.}$$

Note that

$$\left| \sum_{f^n y=x} \frac{h'(y)}{Df^n(y)|Df^n(y)|} \right| = \left| \tilde{\mathcal{L}}^n(h') \right| \leq \lambda^{-n} (\mathcal{L}^n(|h'|))(x),$$

converges to 0 in L^1 and almost everywhere. Thus we focus on $\sum_{f^n(y)=x} \frac{h(y) \cdot D(|Df^n(y)|)}{|Df^n(y)|^2}$.

Assuming that $y \notin \{c, f(c), \dots, f^{n-1}(c)\}$ for each y with $f^n y = x$ we have

$$\begin{aligned} & \sum_{f^n(y)=x} \frac{h(y) \cdot D(|Df(y)|)}{|Df^n(y)|^2} = \sum_{f^n(y)=x} \frac{h(y)}{|Df^n(x)|^2} D \left(\prod_{a=0}^{n-1} |Df(f^a y)| \right) \\ &= \sum_{f^n(y)=x} \frac{h(y)}{|Df^n(y)|^2} \frac{|Df^n(y)|}{Df^n(y)} \sum_{a=0}^{n-1} \xi(f^a(y)) Df^a(y) = \sum_{f^n(y)=x} \frac{h(y)}{|Df^n(y)|} \sum_{a=0}^{n-1} \frac{\xi(f^a(y))}{Df^{n-a}(f^a(y))} \\ &= \sum_{a=0}^{n-1} \sum_{f^{n-a}(z)=x} \frac{\xi(z)}{Df^{n-a}(z)} \sum_{f^a(y)=z} \frac{h(y)}{|Df^n(y)|} = \sum_{a=0}^{n-1} \sum_{f^{n-a}(z)=x} \frac{\xi(z)}{Df^{n-a}(z)|Df^{n-a}(z)|} \sum_{f^a(y)=z} \frac{h(y)}{|Df^a(y)|} \\ &= \sum_{a=0}^{n-1} \sum_{f^{n-a}(z)=x} \frac{\xi(z)}{Df^{n-a}(z)|Df^{n-a}(z)|} \mathcal{L}^a(h)(z) = \sum_{a=0}^{n-1} \tilde{\mathcal{L}}^{n-a}(\xi(\mathcal{L}^a h))(x) \\ &= \sum_{i=1}^n \tilde{\mathcal{L}}^i(\xi(\mathcal{L}^{n-i} h))(x) \end{aligned}$$

Proposition 2.3 shows that we can take the limit $n \rightarrow \infty$ term-by-term. Since

$$\lim_{n \rightarrow \infty} (\mathcal{L}^{n-i} h)(x) = \left(\int_0^1 h(z) dz \right) \rho(x)$$

both parts (a) and (b) follow. \square

At this point, we could get ρ_2 by differentiating each term in (3.1). This is possible due to Lemma 2.4.

Proposition 3.2. *The function ρ_1 is almost everywhere differentiable and*

$$(2) \quad \rho'_1 = 3 \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \tilde{\mathcal{L}}^j(\xi \tilde{\mathcal{L}}^j(\xi \rho)) + 2 \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^j(\xi^2 \rho) - \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^j(\xi' \rho)$$

In particular, there exists $\rho_2 \in BV$ such that $\rho'_1 = \rho_2$ almost everywhere.

Proof. By Lemma 3.1 $\rho_1 = -\sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i(\xi \rho)$ almost everywhere. Therefore by Lemma 2.4

$$\rho'_1(x) = -\sum_{i=1}^{\infty} \left(\sum_{f^i y=x} \frac{\xi(y) \rho(y)}{Df^i(y)|Df^i(y)|} \right)' = -\sum_{i=1}^{\infty} \sum_{f^i y=x} \left(\frac{\xi(y) \rho(y)}{Df^i(y)|Df^i(y)|} \right)'$$

almost everywhere. Decompose

$$\left(\frac{\xi(y)\rho(y)}{Df^i(y)|Df^i(y)|} \right)' = \underbrace{\frac{(\xi(y)\rho(y))'}{Df^i(y)|Df^i(y)|}}_{(I)} - \underbrace{\frac{\xi(y)\rho(y)(Df^i(y)|Df^i(y)|)'}{(Df^i(y))^2|Df^i(y)|^2}}_{(II)}.$$

Let us first work on (I). We have

$$\begin{aligned} \sum_{i=1}^{\infty} \sum_{f^i y=x} (I) &= \sum_{i=1}^{\infty} \sum_{f^i y=x} \frac{\xi'(y)Dy\rho(y) + \xi(y)\rho'(y)Dy}{Df^i(y)|Df^i(y)|} \\ &= \sum_{i=1}^{\infty} \sum_{f^i y=x} \left(\frac{\xi'(y)\rho(y)}{Df^i(y)|Df^i(y)|} + \frac{\xi(y)\rho'(y)}{Df^i(y)|Df^i(y)|} \right) \\ &= \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i (\xi'\rho + \xi\rho')(x) \end{aligned}$$

By Lemma 3.1

$$\sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i (\xi\rho') = - \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \tilde{\mathcal{L}}^i (\xi\tilde{\mathcal{L}}^j(\xi\rho)).$$

Therefore

$$(3) \quad \sum_{i=1}^{\infty} \sum_{f^i y=x} (I) = \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i (\xi'\rho) - \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i (\xi\tilde{\mathcal{L}}^j(\xi\rho)).$$

Now, let us analyze (II).

$$\begin{aligned} \sum_{i=1}^{\infty} \sum_{f^i y=x} (II) &= \sum_{i=1}^{\infty} \sum_{f^i y=x} \frac{\xi(y)\rho(y)[D(Df^i y)|Df^i y| + D(|Df^i y|)(Df^i)(y)]}{(Df^i y)^2|Df^i y|^2} (Df^i y)^2|Df^i y|^2 \\ &= \sum_{i=1}^{\infty} \sum_{f^i y=x} \frac{\xi(y)\rho(y) \left[2|Df^i y| \sum_{j=0}^{i-1} \xi(f^j y) Df^j y \right]}{(Df^i y)^2|Df^i y|^2} = 2 \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \sum_{f^i y=x} \frac{\xi(y)\rho(y) Df^j y \xi(f^j y)}{(Df^i y)^2|Df^i y|} \end{aligned}$$

By making the change of variable $z = f^j y$, we obtain

$$\begin{aligned} \sum_{i=1}^{\infty} \sum_{f^i y=x} (II) &= 2 \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \sum_{f^i y=x} \frac{\xi(y)\rho(y)(Df^j)(y)\xi(z)}{(Df^i y)^2|Df^i y|} \\ &= 2 \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \sum_{f^i y=x} \frac{\xi(y)\rho(y)(Df^j)(y)\xi(z)}{(Df^{i-j} z)^2(Df^j y)^2|Df^{i-j} z||Df^j y|} = 2 \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \sum_{f^i y=x} \frac{\xi(y)\rho(y)\xi(z)}{(Df^{i-j} z)^2 Df^j y |Df^{i-j} z| |Df^j y|} \\ &= 2 \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \sum_{f^{i-j} z=x} \frac{\xi(z)}{(Df^{i-j} z)^2 |Df^{i-j} z|} \sum_{f^i y=z} \frac{\xi(y)\rho(y)}{Df^j y |Df^j y|} = 2 \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \left(\tilde{\mathcal{L}} \right)^{i-j} \left(\xi\tilde{\mathcal{L}}^j(\xi\rho) \right). \end{aligned}$$

By Lemma 2.5

$$\sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \left(\tilde{\mathcal{L}} \right)^{i-j} \left(\xi \tilde{\mathcal{L}}^j(\xi\rho) \right) = \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} \left(\tilde{\mathcal{L}} \right)^{i-j} \left(\xi \tilde{\mathcal{L}}^j(\xi\rho) \right).$$

Therefore

$$(4) \quad \sum_{i=1}^{\infty} \sum_{f^i y=x} (II) = \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} \left(\tilde{\mathcal{L}} \right)^{i-j} \left(\xi \tilde{\mathcal{L}}^j(\xi\rho) \right).$$

Combining (3) and (4), we finally obtain

$$\begin{aligned} \rho'_1 &= - \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i(\xi'\rho) + \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \tilde{\mathcal{L}}^j(\xi \tilde{\mathcal{L}}^j(\xi\rho)) + \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} \left(\tilde{\mathcal{L}} \right)^j \left(\xi \tilde{\mathcal{L}}^j(\xi\rho) \right) \\ &= 3 \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \tilde{\mathcal{L}}^j(\xi \tilde{\mathcal{L}}^j(\xi\rho)) + 2 \sum_{i=1}^{\infty} \left(\tilde{\mathcal{L}} \right)^i \left(\xi^2 \rho \right) - \sum_{i=1}^{\infty} \tilde{\mathcal{L}}^i(\xi'\rho) \end{aligned}$$

almost everywhere as claimed. \square

3.2. Higher order derivatives. Lemma 3.1 shows that ρ' is in BV . Then we saw in Proposition 3.2 that $\rho'_1 = \rho_2 \in BV$. Here we show that these results can be extended to repeated differentiation of arbitrary order. We start with the following general result.

Proposition 3.3. *Let k, i_1, \dots, i_k and $m_1 > \dots > m_k$ be positive integers with $i_1, \dots, i_k \geq 1$. Let h_1, \dots, h_k be BV functions whose derivatives are in L^∞ .*

(a) *If $n \geq 1$, the derivative of $\sum_{k \leq i_1 + \dots + i_k \leq n} \mathfrak{D}_{m, m_2, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)$ is a finite sum of functions of the type⁴ $\sum_{\tilde{k} \leq \tilde{i}_1 + \dots + \tilde{i}_{\tilde{k}} \leq n} \mathfrak{D}_{m+1, \tilde{m}_2, \dots, \tilde{m}_{\tilde{k}}}^{\tilde{i}_1, \dots, \tilde{i}_{\tilde{k}}}(\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}})$, where $k \leq \tilde{k} \leq k+1, \tilde{i}_1, \dots, \tilde{i}_{\tilde{k}} \geq 1$,*

$\tilde{m}_1 > \dots > \tilde{m}_{\tilde{k}}$ are positive integers and $\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}} \in \{h_1, \dots, h_k, h'_1, \dots, h'_k, \xi, \xi'\}$.

(b) *The derivative of*

$$\sum_{i_1=1}^{\infty} \dots \sum_{i_k=1}^{\infty} \mathfrak{D}_{m, m_2, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)$$

equals almost everywhere to a finite sum of functions of the type $\sum_{\tilde{i}_1} \dots \sum_{\tilde{i}_{\tilde{k}}} \mathfrak{D}_{m+1, \tilde{m}_2, \dots, \tilde{m}_{\tilde{k}}}^{\tilde{i}_1, \dots, \tilde{i}_{\tilde{k}}}(\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}})$,

where $k \leq \tilde{k} \leq k+1, \tilde{i}_1, \dots, \tilde{i}_{\tilde{k}} \geq 1, \tilde{m}_1 > \dots > \tilde{m}_{\tilde{k}}$ are positive integers and

$$\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}} \in \{h_1, \dots, h_k, h'_1, \dots, h'_k, \xi, \xi'\}.$$

⁴That is, the sums coincide at the points where both of them are defined.

Proof. We will prove (a) by induction on m . For $m = 1$, we need to compute $\sum_{i=1}^n D\left(\tilde{\mathcal{L}}^i(h)\right)$,

so let us work on $D\left(\tilde{\mathcal{L}}^i(h)\right)$. Then

$$\begin{aligned} D\left(\tilde{\mathcal{L}}^i(h)\right)(x) &= \sum_{f^i(y)=x} D\left(\frac{h(y)}{Df^i y|Df^i y|}\right) = \sum_{f^i(y)=x} \left[\frac{h'(y)}{Df^i y|Df^i y|} - \frac{h(y)D(Df^i y|Df^i y|)}{(Df^i y)^2|Df^i y|^2} \right] \\ &= \sum_{f^i(y)=x} \left[\frac{h'(y)Dy}{Df^i y|Df^i y|} - \frac{h(y)(D(Df^i y|Df^i y|) + Df^i yD(|Df^i y|))}{(Df^i y)^2|Df^i y|^2} \right] \\ &= \sum_{f^i(y)=x} \left[\frac{h'(y)}{(Df^i y)^2|Df^i y|} - \frac{h(y)(2|Df^i y|\sum_{j=0}^{i-1}\xi(f^j(x))Df^j(x))}{(Df^i y)^2|Df^i y|^2} \right] \\ &= \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{j=0}^{i-1}\sum_{f^i(y)=x} \frac{h(y)\xi(f^j(y))Df^j(y)}{(Df^i y)^2|Df^i y|}. \end{aligned}$$

Let $z = f^j(y)$. Then

$$\begin{aligned} D\left(\tilde{\mathcal{L}}^i(h)\right)(x) &= \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{j=0}^{i-1}\sum_{f^i(y)=x} \frac{h(y)\xi(f^j(y))Df^j(y)}{(Df^i y)^2|Df^i y|} \\ &= \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{j=0}^{i-1}\sum_{f^i(y)=x} \frac{h(y)\xi(z)}{(Df^{i-j}z)^2|Df^{i-j}z|(Df^j y)|Df^j y|} \\ &= \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{j=0}^{i-1}\sum_{f^{i-j}z=x} \frac{\xi(z)}{(Df^{i-j}z)^2|Df^{i-j}z|} \sum_{f^j y=z} \frac{h(y)}{(Df^j y)|Df^j y|} \\ &= \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{j=0}^{i-1}\tilde{\mathcal{L}}^{i-j}(\xi\tilde{\mathcal{L}}^j(h)) \end{aligned}$$

Hence

$$\begin{aligned} \sum_{i=1}^n D\left(\tilde{\mathcal{L}}^i(h)\right) &= \sum_{i=1}^n \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{i=1}^n \sum_{j=0}^{i-1} \tilde{\mathcal{L}}^{i-j}(\xi\tilde{\mathcal{L}}^j(h)) \\ &= \sum_{i=1}^n \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{i=1}^n \tilde{\mathcal{L}}^i(\xi h) - 2\sum_{i=1}^n \sum_{j=1}^{i-1} \tilde{\mathcal{L}}^{i-j}(\xi\tilde{\mathcal{L}}^j(h)) \\ &= \sum_1^n \tilde{\mathcal{L}}^i(h')(x) - 2\sum_{i=1}^n \tilde{\mathcal{L}}^i(\xi h) - 2\sum_{\substack{2 \leq i+j \leq n \\ 1 \leq i, 1 \leq j}} \tilde{\mathcal{L}}^i(\xi\tilde{\mathcal{L}}^j(h)) \end{aligned}$$

Therefore, the derivative is a finite sum of terms as described in the statement.

Assume the statement is true for $l < m$. Let us prove that it also holds for m . We are interested in the derivative of

$$(5) \quad \sum_{k+1 \leq i+i_1+\dots+i_k \leq n} \mathfrak{D}_{m,m_1,\dots,m_k}^{i,i_2,\dots,i_k}(h, h_1, \dots, h_k)$$

with $i \geq 1, i_1 \geq 1, \dots, i_k \geq 1$. For this, note that

$$\sum_{k+1 \leq i+i_1+\dots+i_k \leq n} \mathfrak{D}_{m,m_1,\dots,m_k}^{i,i_1,\dots,i_k}(h, h_1, \dots, h_k) = \sum_{i=1}^{n-k} \left(\widetilde{L} \right)^i \left(h \sum_{k \leq i_1+\dots+i_k \leq n-i} \mathfrak{D}_{m_1,\dots,m_k}^{i_1,\dots,i_k}(h_1, \dots, h_k) \right)$$

Thus, if we are interested in the derivative of (5), we need to analyze

$$\begin{aligned} & \sum_{i=1}^{n-k} D \left[\left(\widetilde{L} \right)^i \left(h \sum_{k \leq i_1+\dots+i_k \leq n-i} \mathfrak{D}_{m_1,\dots,m_k}^{i_1,\dots,i_k}(h_1, \dots, h_k) \right) \right] \\ &= \sum_{i=1}^{n-k} \sum_{f^i y=x} D \left[\frac{h(y) \sum_{k \leq i_1+\dots+i_k \leq n-i} \mathfrak{D}_{m_1,\dots,m_k}^{i_1,\dots,i_k}(h_1, \dots, h_k)(y)}{(Df^i y)^m |Df^i y|} \right] = \sum_{i=1}^{n-k} \sum_{f^i y=x} (I) - (II) \end{aligned}$$

where

$$\begin{aligned} (I) &= \frac{D \left[h(y) \widehat{\sum} \mathfrak{D}_{m_1,\dots,m_k}^{i_1,\dots,i_k}(h_1, \dots, h_k)(y) \right]}{(Df^i y)^m |Df^i y|}, \\ (II) &= \frac{h(y) \widehat{\sum} \mathfrak{D}_{m_1,\dots,m_k}^{i_1,\dots,i_k}(h_1, \dots, h_k)(y) D \left[(Df^i y)^{2m} |Df^i y|^2 \right]}{(Df^i y)^{2m} |Df^i y|^2} \end{aligned}$$

and $\widehat{\sum}$ means $\sum_{k \leq i_1+\dots+i_k \leq n-i}$.

Let us first work on (II). Note that

$$\begin{aligned} & D \left[(Df^i y)^m |Df^i y| \right] = m(Df^i y)^{m-1} D(Df^i y) |Df^i y| + (Df^i y)^m D(|Df^i y|) \\ &= m(Df^i y)^{m-1} |Df^i y| \sum_{j=0}^{i-1} \xi(f^j y) Df^j y + (Df^i y)^m \frac{|Df^i y|}{(Df^i y)} \sum_{j=0}^{i-1} \xi(f^j y) Df^j y \\ &= m(Df^i y)^{m-1} |Df^i y| \sum_{j=0}^{i-1} \xi(f^j y) Df^j y + (Df^i y)^{m-1} |Df^i y| \sum_{j=0}^{i-1} \xi(f^j y) Df^j y \\ &= (m+1)(Df^i y)^{m-1} |Df^i y| \sum_{j=0}^{i-1} \xi(f^j y) Df^j y \end{aligned}$$

Then $\sum_{i=1}^{n-k} \widehat{\sum}_{f^i y=x} \sum (II)$ equals

$$\begin{aligned} & \sum_{i=1}^{n-k} \widehat{\sum}_{f^i y=x} \sum \frac{(m+1)h(y) \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) (Df^i y)^{m-1} |Df^i y| \sum_{j=0}^{i-1} \xi(f^j y) Df^j y}{(Df^i y)^{2m} |Df^i y|^2} \\ &= \sum_{i=1}^{n-k} \sum_{j=0}^{i-1} \widehat{\sum}_{f^i y=x} \sum \frac{(m+1)h(y) \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) \xi(f^j y) Df^j y}{(Df^i y)^{m+1} |Df^i y|}. \end{aligned}$$

Let $z = f^j y$. Then $\sum_{i=1}^{n-k} \widehat{\sum}_{f^i y=x} \sum (II)$ equals

$$\begin{aligned} &= \sum_{i=1}^{n-k} \sum_{j=0}^{i-1} \widehat{\sum}_{f^i y=x} \sum \frac{(m+1)h(y) \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) \xi(f^j y) Df^j y}{(Df^i y)^{m+1} |Df^i y|} \\ &= \sum_{i=1}^{n-k} \sum_{j=0}^{i-1} \widehat{\sum}_{f^i y=x} \sum \frac{(m+1)h(y) \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) \xi(z) Df^j y}{(Df^{i-j} z)^{m+1} |Df^{i-j} z| (Df^j y)^{m+1} |Df^j y|} \\ &= \sum_{i=1}^{n-k} \sum_{j=0}^{i-1} \widehat{\sum}_{f^i y=x} \sum \frac{(m+1)h(y) \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) \xi(z)}{(Df^{i-j} z)^{m+1} |Df^{i-j} z| (Df^j y)^m |Df^j y|} \\ &= \sum_{i=1}^{n-k} \sum_{j=0}^{i-1} \widehat{\sum}_{f^{i-j} z=x} \sum \frac{\xi(z)}{(Df^{i-j} z)^{m+1} |Df^{i-j} z|} \sum_{f^j y=z} \frac{(m+1)h(y) \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y)}{(Df^j y)^m |Df^j y|} \\ &= (m+1) \sum_{i=1}^{n-k} \sum_{j=0}^{i-1} \widehat{\sum} \left(\frac{m+1}{\tilde{L}} \right)^{i-j} \left(\xi \left(\frac{m}{\tilde{L}} \right)^j \left(h \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k) \right) \right) (x) \\ &= (m+1) \sum_{i=1}^{n-k} \widehat{\sum} \left(\frac{m+1}{\tilde{L}} \right)^i \left(\xi \left(h \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k) \right) \right) (x) + \\ & \quad (m+1) \sum_{\substack{1 \leq i+j \leq n-k \\ 1 \leq i, 1 \leq j}} \widehat{\sum} \left(\frac{m+1}{\tilde{L}} \right)^{i-j} \left(\xi \left(\frac{m}{\tilde{L}} \right)^j \left(h \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k) \right) \right) (x) = A + B. \end{aligned}$$

The last two terms can be rewritten as

$$\begin{aligned} A &= (m+1) \sum_{1 \leq i+i_1+\dots+i_k \leq n} \left(\frac{m+1}{\tilde{L}} \right)^i \left(\xi \left(h \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k) \right) \right) (x), \\ B &= (m+1) \sum_{k+2 \leq i+j+i_1+\dots+i_k \leq n} \left(\frac{m+1}{\tilde{L}} \right)^i \left(\xi \left(\frac{m}{\tilde{L}} \right)^j \left(h \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k) \right) \right) (x). \end{aligned}$$

Therefore $\sum_{i=1}^{n-k} \sum_{f^i y=x} (II)$ is a sum of terms described in the statement.

Now, let us analyze (I). Note that $\sum_{i=1}^{n-k} \sum_{f^i y=x} (I)$ equals to

$$\begin{aligned} & \sum_{i=1}^{n-k} \sum_{f^i y=x} \frac{h'(y) Dy \widehat{\sum} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y)}{(Df^i y)^m |Df^i y|} - \frac{h(y) D \left[\widehat{\sum} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) \right]}{(Df^i y)^m |Df^i y|} \\ &= \sum_{i=1}^{n-k} \sum_{f^i y=x} \frac{h'(y) \widehat{\sum} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y)}{(Df^i y)^{m+1} |Df^i y|} - \frac{h(y) D \left[\widehat{\sum} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) \right]}{(Df^i y)^m |Df^i y|} \\ &= \sum_{i=1}^{n-k} \binom{m+1}{\tilde{\mathcal{L}}}^i \left(h' \widehat{\sum} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k) \right)(y) - \sum_{i=1}^{n-k} \sum_{f^i y=x} \frac{h(y) \widehat{\sum} D \left[\mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)(y) \right]}{(Df^i y)^m |Df^i y|}. \end{aligned}$$

Using our inductive hypothesis, the derivative of $\widehat{\sum} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k)$ is the finite sum of terms of the type $\sum_{\tilde{k} \leq \tilde{i}_1 + \dots + \tilde{i}_{\tilde{k}} \leq n-i} \mathfrak{D}_{m_1+1, \tilde{m}_2, \dots, \tilde{m}_{\tilde{k}}}^{\tilde{i}_1, \tilde{i}_2, \dots, \tilde{i}_{\tilde{k}}}(\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}})$. Hence, let us take one of these terms and analyze the expression

$$\begin{aligned} & \sum_{i=1}^{n-k} \sum_{\tilde{k} \leq \tilde{i}_1 + \dots + \tilde{i}_{\tilde{k}} \leq n-i} \sum_{f^i y=x} \frac{h(y) \mathfrak{D}_{m_1+1, \tilde{m}_2, \dots, \tilde{m}_{\tilde{k}}}^{i_1+1, \tilde{i}_2, \dots, \tilde{i}_{\tilde{k}}}(\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}})(y) \cdot Dy}{(Df^i y)^m |Df^i y|} \\ &= \sum_{i=1}^{n-k} \sum_{\tilde{k} \leq \tilde{i}_1 + \dots + \tilde{i}_{\tilde{k}} \leq n-i} \sum_{f^i y=x} \frac{h(y) \mathfrak{D}_{m_1+1, \tilde{m}_2, \dots, \tilde{m}_{\tilde{k}}}^{i_1+1, \tilde{i}_2, \dots, \tilde{i}_{\tilde{k}}}(\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}})(y)}{(Df^i y)^{m+1} |Df^i y|} \\ &= \sum_{i=1}^{n-k} \sum_{\tilde{k} \leq \tilde{i}_1 + \dots + \tilde{i}_{\tilde{k}} \leq n-i} \binom{m+1}{\tilde{\mathcal{L}}}^i \left(h \mathfrak{D}_{m_1+1, \tilde{m}_2, \dots, \tilde{m}_{\tilde{k}}}^{i_1+1, \tilde{i}_2, \dots, \tilde{i}_{\tilde{k}}}(\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}}) \right) \\ &= \sum_{\tilde{k}+1 \leq 1 + \tilde{i}_1 + \dots + \tilde{i}_{\tilde{k}} \leq n} \binom{m+1}{\tilde{\mathcal{L}}}^i \left(h \mathfrak{D}_{m_1+1, \tilde{m}_2, \dots, \tilde{m}_{\tilde{k}}}^{i_1+1, \tilde{i}_2, \dots, \tilde{i}_{\tilde{k}}}(\tilde{h}_1, \dots, \tilde{h}_{\tilde{k}}) \right). \end{aligned}$$

Since we have a finite sums of terms as above, we obtained that our proposition also holds for m . Therefore, part (a) is established by induction.

(b) Lemma 2.4 and Proposition 2.3 allow us to take the limit $n \rightarrow \infty$. Then the condition $k \leq i_1 + \dots + i_k \leq n$ becomes $k \leq i_1 + \dots + i_k \leq \infty$ and using the condition $i_1 \geq 1, \dots, i_k \geq 1$ the sum $k \leq i_1 + \dots + i_k \leq n$ converges to

$$\sum_{i_1=1}^{\infty} \cdots \sum_{i_k=1}^{\infty}.$$

Therefore part (b) follows from part (a). \square

Proposition 3.3(b) allows us to derive Theorem 1.1.

Proof of Theorem 1.1. We have already defined ρ_1 in Lemma 3.1 and ρ_2 in Proposition 2. Continuing this procedure, Suppose that we have already defined $\rho_{k-1} \in BV$. Let $\rho_k = \rho'_{k-1}$. Then Proposition 3.3(b) can be used to show inductively that

$$\rho_k = \sum_{\text{finite } i_1, \dots, i_s \geq 1} \sum_{k, m_2, \dots, m_s} \mathfrak{D}_{k, m_2, \dots, m_s}^{i_1, \dots, i_s}(h_1, \dots, h_{s-1}, \rho)$$

where $s \geq 1$, $k-1 > m_2 > \dots > m_s$, and h_1, \dots, h_{s-1} are C^1 away from c (in fact h_j are rational functions of derivatives of f). Now Proposition 2.3 implies that $\rho_k \in BV$ as claimed. \square

4. DIFFERENTIABILITY SET FOR THE DENSITY.

4.1. Saltus part. Any function of bounded variation ϕ can be decomposed as

$$\phi = \phi_r + \phi_s$$

where ϕ_r is a continuous function, called the regular part, and ϕ_s is constant except at discontinuities of ϕ . ϕ_s is called the saltus part, it is discontinuous on a countable set (see [11], page 14)

In fact, in the case of ρ , ρ_s can be explicitly written as ([2])

$$\rho_s = \sum_{j \geq 1} \alpha_j H_{c_j}$$

where $c_j = f^j(c)$, $\alpha_j = \lim_{x \uparrow c_j} \rho(x) - \lim_{x \downarrow c_j} \rho(x)$ and H_{c_j} is defined as

$$(6) \quad H_{c_j}(x) = \begin{cases} 1 & \text{if } x < c_j \\ \frac{1}{2} & \text{if } x = c_j \\ 0 & \text{if } x > c_j \end{cases}$$

Lemma 4.1. *If c is not periodic then*

$$\alpha_j = \pm \rho(c) \left[\frac{1}{|Df_+^j(c)|} + \frac{1}{|Df_-^j(c)|} \right],$$

where the expression takes the sign $+$ (resp. the sign $-$) if f has a maximum (resp. minimum) at c .

Proof. We have

$$\alpha_j = \lim_{x \uparrow c_j} \rho(x) - \lim_{x \downarrow c_j} \rho(x).$$

Using the fact that ρ is a fixed point of \mathcal{L} and $\mathcal{L}^j \rho(x) = \sum_{f^j y = x} \frac{\rho(y)}{Df^j(y)}$, we can see

that ρ has a discontinuity at $x = c_j$. In fact, among all the y 's in the set $\{f^{-j}c_j\}$, the discontinuity comes from $y = c$, therefore

$$\alpha_j = \lim_{y \uparrow c} \frac{\rho(y)}{Df^j(y)} - \lim_{y \downarrow c} \frac{\rho(y)}{Df^j(y)}. \quad \square$$

Proposition 4.2. *For $k \geq 0$, the element ρ_k of the sequence from Theorem 1.1 can be decomposed as $(\rho_k)_r + (\rho_k)_s$, where $(\rho_k)_r$ is a continuous function and $(\rho_k)_s = \sum_{m \geq 1} \alpha_{k,j} H_{c_j}$, with H_{c_j} defined in (6) and $\alpha_{k,j} = \lim_{x \uparrow c_j} \rho_k(x) - \lim_{x \downarrow c_j} \rho_k(x)$. Moreover there exists $\theta < 1$ such that $|\alpha_{k,j}| \leq K\theta^j$*

Proof. The existence of decomposition follows from the fact that, due to Theorem 1.1, $\rho_k \in BV$ -function. We need to show that all discontinuities of ρ_k lie on the critical orbit and bound the size of discontinuity.

Let z be a discontinuity point of ρ_k which is different from c_i for $i = 1 \dots j$. Let $\bar{\rho} = \mathcal{L}^j(1)$. In the proof of Proposition 1.1 we saw that

$$\begin{aligned} \rho_k &= \sum_{\text{finite } i, i_2, \dots, i_s \geq 1} \sum \mathfrak{D}_{k, m_2, \dots, m_k}^{i, i_2, \dots, i_s}(h_1, \dots, h_{s-1}, \rho) \\ &= \sum_{\text{finite } i, i_2, \dots, i_s \geq 1} \sum \mathfrak{D}_{k, m_2, \dots, m_k}^{i, i_2, \dots, i_s}(h_1, \dots, h_{s-1}, \bar{\rho}) + \sum_{\text{finite } i, i_2, \dots, i_s \geq 1} \sum \mathfrak{D}_{k, m_2, \dots, m_k}^{i, i_2, \dots, i_s}(h_1, \dots, h_{s-1}, \rho - \bar{\rho}). \end{aligned}$$

Denote $\Delta(h) = \lim_{x \uparrow z} h(x) - \lim_{x \downarrow z} h(x)$. Then

$$\Delta \left(\sum_{\text{finite } i, i_2, \dots, i_s \geq 1} \sum \mathfrak{D}_{k, m_2, \dots, m_k}^{i, i_2, \dots, i_s}(h_1, \dots, h_{s-1}, \rho - \bar{\rho}) \right) = O(\theta^j)$$

in view of Proposition 2.3 and the fact that $\rho - \bar{\rho} = O(\theta^j)$.

Note that if $i, i_2, \dots, i_s < j$ then $\left(\tilde{\mathcal{L}} \right)^i$ and $\left(\tilde{\mathcal{L}} \right)^{i_r}$ are continuous at z for $r = 2, \dots, s$, so

$$\sum_{\text{finite } i, i_2, \dots, i_k < j} \Delta \left(\sum \mathfrak{D}_{k, m_2, \dots, m_s}^{i, i_2, \dots, i_s}(h_1, \dots, h_{s-1}, \bar{\rho}) \right) = 0.$$

Applying Proposition 2.3 again we see that

$$\sum_{\text{finite } \max(i, i_2, \dots, i_s) > j} \sum \mathfrak{D}_{k, m_2, \dots, m_s}^{i, i_2, \dots, i_s}(h_1, \dots, h_{s-1}, \bar{\rho}) = O \left(\sum_{\max(i, i_2, \dots, i_s) > j} \lambda^{-(i+i_2+\dots+i_s)} \right)$$

and since the expression in the right side is $O(j^s \lambda^{-j})$, we have

$$\Delta \left(\sum_{\text{finite } \max(i, i_2, \dots, i_s) > j} \sum \mathfrak{D}_{k, m_2, \dots, m_s}^{i, i_2, \dots, i_s}(h_1, \dots, h_{s-1}, \bar{\rho}) \right) \leq O(j^s \lambda^{-j}).$$

In particular if z is not on the critical orbit then $\Delta \rho_k = 0$ and if $z = c_j$ then $\Delta \rho_k$ is exponentially small in j as claimed. \square

4.2. Absolute continuity. As we mentioned before, the regular part of ρ is continuous. In fact, it is absolutely continuous.

Theorem 4.3. *The regular part of ρ is absolutely continuous. That is*

$$\rho_r(x_2) - \rho_r(x_1) = \int_{x_1}^{x_2} \rho'(x) dx.$$

Proof. Let $n \geq 1$ and let $x_2, x_1 \in [0, 1]$. Then

$$(7) \quad \begin{aligned} & (\mathcal{L}^n(1))(x_2) - (\mathcal{L}^n(1))(x_1) \\ &= \int_{x_1}^{x_2} (\mathcal{L}^n(1))'(x) dx + \sum_{\substack{j \leq n \\ c_j \in [x_1, x_2]}} \Delta_j(\mathcal{L}^n(1)), \end{aligned}$$

where $\Delta_j(\mathcal{L}^n(1)) = \lim_{x \uparrow c_j} \mathcal{L}^n(1)(x) - \lim_{x \downarrow c_j} \mathcal{L}^n(1)(x)$.

As $n \rightarrow \infty$, $(\mathcal{L}^n(1))(x) \rightarrow \rho(x)$. Hence, $\Delta_j(\mathcal{L}^n(1)) \rightarrow \Delta_j \rho$. By Lemma 3.1 $(\mathcal{L}^n(1))' \rightarrow \rho_1$ as $n \rightarrow \infty$. Thus letting $n \rightarrow \infty$ in (7) we get

$$\begin{aligned} \rho(x_2) - \rho(x_1) &= \int_{x_1}^{x_2} \rho_1(x) dx + \sum_{c_j \in [x_1, x_2]} \Delta_j \rho \\ &= \int_{x_1}^{x_2} \rho_1(x) dx + \rho_s(x_2) - \rho_s(x_1) \end{aligned}$$

Therefore $\rho_r(x_2) - \rho_r(x_1) = \int_{x_1}^{x_2} \rho_1(x) dx$. □

Proposition 4.4. *There exist constants $K \geq 1$, $D \geq 1$ and $\varsigma < 1$ such that if \bar{x} satisfies*

$$(8) \quad d(c_j, \bar{x}) > \epsilon,$$

for $j \leq n$ and $d(x, \bar{x}) < \epsilon$, then

$$|\rho(x) - \rho(\bar{x})| \leq K\epsilon + D\varsigma^n.$$

Proof. Decompose

$$(9) \quad \rho(x) - \rho(\bar{x}) = (\rho_r(x) - \rho_r(\bar{x})) + (\rho_s(x) - \rho_s(\bar{x})).$$

Combining Theorem 4.3 with the fact that $\rho' = \rho_1 \in BV[0, 1]$, we get

$$(10) \quad |\rho_r(x) - \rho_r(\bar{x})| \leq K\epsilon.$$

Also, (8) implies

$$(11) \quad \rho_s(x) - \rho_s(\bar{x}) = \sum_{j \geq n} \alpha_j [H_{c_j}(x) - H_{c_j}(\bar{x})].$$

By Lemma 4.1 $|\alpha_j| \leq \frac{2\|\rho\|_\infty}{\lambda^j}$. Hence, we can bound (11) as

$$|\rho_s(x) - \rho_s(\bar{x})| \leq \sum_{j \geq n} |\alpha_j| \left| H_{c_j}(x) - H_{c_j}(\bar{x}) \right| \leq 2\|\rho\|_\infty \sum_{j \geq n} \frac{1}{\lambda^j}$$

$$= 2\|\rho\|_\infty \frac{1}{\lambda^n} \sum_{j \geq 1} \frac{1}{\lambda^j} = 2\|\rho\|_\infty \left(\frac{\lambda}{\lambda-1} \right) \frac{1}{\lambda^n}$$

Taking $D = 2\|\rho\|_\infty \left(\frac{\lambda}{\lambda-1} \right)$, $\varsigma = \frac{1}{\lambda}$, we have

$$(12) \quad |\rho_s(x) - \rho_s(\bar{x})| \leq D\varsigma^n.$$

Combining (9), (10) and (12) we obtain the result. \square

4.3. Differentiability points. Recall that there is a constant $\theta < 1$ such that

$$\mathcal{L}^n h = \left[\int h(z) dz \right] \rho(x) + O(\theta^n \|h\|_{BV}).$$

(see e.g. [1], Proposition 3.5, item 4)

Theorem 4.5. *If $1 > \beta > \max(\theta, 1/\lambda)$ and if \bar{x} is a point such that $d(\bar{x}, c_j) \geq \beta^j$ for all $j \geq j_0$ then ρ_k is differentiable at \bar{x} .*

Proof. Let $\epsilon > 0$ and let x such that $d(x, \bar{x}) = \epsilon$.

Let n be the maximal number such that

$$(13) \quad c_j \notin [x; \bar{x}] \text{ for all } j \leq n.$$

Then $\epsilon \geq \beta^n$, hence $\epsilon\lambda^n \geq \beta^n \lambda^n$ and $\frac{\epsilon}{\theta^n} \geq \frac{\beta^n}{\theta^n}$.

By definition of β , $\beta\lambda > 1$ and $\frac{\beta}{\theta} > 1$. Hence, $\beta^n \lambda^n \rightarrow \infty$ and $\frac{\beta^n}{\theta^n} \rightarrow \infty$ as $n \rightarrow \infty$. Therefore,

$$\epsilon\lambda^n \rightarrow \infty$$

and

$$\frac{\epsilon}{\theta^n} \rightarrow \infty.$$

as $n \rightarrow \infty$.

By Theorem 1.1

$$\rho_k(x) = \sum_{\text{finite } i_1, \dots, i_k=1}^{\infty} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \rho).$$

Let $\bar{\rho} = \mathcal{L}^n(1)$. Since $\rho = \bar{\rho} + O(\theta^n)$, Proposition 2.3 implies that we can write the above expression as

$$\rho_k(x) = \sum_{\substack{\text{finite } k \leq i_1, \dots, i_k < n \\ 1 \leq i_1, \dots, 1 \leq i_k}} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho}) + O(\lambda^{-n} + \theta^n).$$

Therefore

$$(14) \quad \rho_k(x) - \rho_k(\bar{x}) =$$

$$\sum_{\substack{k \leq i_1, \dots, i_k < n \\ 1 \leq i_1, \dots, 1 \leq i_k}} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho})(x) - \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho})(\bar{x}) + O(\lambda^{-n} + \theta^n).$$

Note that $\mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho})$ is differentiable in $[x; \bar{x}]$ since $h_1 \dots h_k$ are C^1 away from c and (13) ensures that $f^{-n}[x, \bar{x}]$ does not contain c .

Thus

$$(15) \quad \begin{aligned} & \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho})(x) - \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho})(\bar{x}) \\ &= \int_{\bar{x}}^x \left(\mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho}) \right)'(s) ds \end{aligned}$$

By Proposition 3.3

$$\left(\sum_{\substack{k \leq i_1, \dots, i_k < n \\ 1 \leq i_1, \dots, 1 \leq i_k}} \mathfrak{D}_{m_1, \dots, m_k}^{i_1, \dots, i_k}(h_1, \dots, h_k, \bar{\rho}) \right)' = \sum_{finite} \sum_{\tilde{i}_1, \dots, \tilde{i}_k} \mathfrak{D}_{m_1+1, \dots, m_k}^{\tilde{i}_1, \dots, \tilde{i}_k}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n),$$

where $\tilde{h}_1, \dots, \tilde{h}_k \in \{h_1, h_2, \dots, h_k, h'_1, \dots, h'_k, \xi, \xi'\}$ and $\Upsilon_n \in \{\bar{\rho}, \bar{\rho}'\}$. Hence

$$(15) = \int_{\bar{x}}^x \sum_{finite} \sum_{\tilde{i}_1, \dots, \tilde{i}_k} \mathfrak{D}_{m_1+1, \dots, m_k}^{\tilde{i}_1, \dots, \tilde{i}_k}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)(s) ds$$

Decompose the last integral as

$$\begin{aligned} & \int_{\bar{x}}^x \sum_{finite} \sum_{\tilde{i}_1, \dots, \tilde{i}_k} \mathfrak{D}_{m_1+1, \dots, m_k}^{\tilde{i}_1, \dots, \tilde{i}_k}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)(s) ds = \sum_{finite} \sum_{\tilde{i}_1, \dots, \tilde{i}_k} \mathfrak{D}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)(\bar{x})(x - \bar{x}) + \\ & + \int_{\bar{x}}^x \left[\mathfrak{D}_{m_1+1, \dots, m_k}^{\tilde{i}_1, \dots, \tilde{i}_k}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)(s) - \mathfrak{D}_{m_1+1, \dots, m_k}^{\tilde{i}_1, \dots, \tilde{i}_k}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)(\bar{x}) \right] ds \end{aligned}$$

We now invoke Proposition 3.3 again which together with (13) implies that $\mathfrak{D}_{m_1+1, \dots, m_k}^{\tilde{i}_1, \dots, \tilde{i}_k}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)$ is differentiable on $[x; \bar{x}]$. Moreover, by Proposition 2.3 its derivative is bounded by a constant M . Hence the last integrand in the above formula is $O(\epsilon)$ and so the integral is $O(\epsilon^2)$. Accordingly

$$(15) = (x - \bar{x}) \sum_{finite} \sum_{\tilde{i}_1, \dots, \tilde{i}_k} \mathfrak{D}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)(\bar{x}) + O(\epsilon^2).$$

Hence

$$\lim_{x \rightarrow \bar{x}} \frac{\rho_k(x) - \rho_k(\bar{x})}{x - \bar{x}} = \lim_{x \rightarrow \bar{x}} \sum_{finite} \sum_{1 \leq i_1, \dots, i_k < n} \sum_{finite} \sum_{\tilde{i}_1, \dots, \tilde{i}_k} \mathfrak{D}(\tilde{h}_1, \dots, \tilde{h}_k, \Upsilon_n)(\bar{x}) + O\left(\epsilon + \frac{\lambda^{-n} + \theta^n}{\epsilon}\right).$$

As x approaches \bar{x} , n goes to ∞ , hence Υ_n converges to ρ or ρ_1 . Thus,

$$\lim_{x \rightarrow \bar{x}} \frac{\rho_k(x) - \rho_k(\bar{x})}{x - \bar{x}} = \sum_{\text{finite } i_1, \dots, i_k=1} \sum_{i=1}^{\infty} \sum_{\text{finite } \tilde{i}_1, \dots, \tilde{i}_k} \mathfrak{D}(\tilde{h}_1, \dots, \tilde{h}_k, \tilde{\rho})(\bar{x}) = \rho_{k+1}(\bar{x}). \quad \square$$

In particular, we have the following result which also follows from [18].

Corollary 4.6. *If c is periodic of period p , then ρ is differentiable except for a finite set of points.*

Proof. If \bar{x} does not belong to the orbit of c (which is a finite set) then we can pick any $\beta > \max(\theta, 1/\lambda)$ and pick $j_0 \geq 1$ large enough so that $d(\bar{x}, \bar{c}) \geq \beta^j$ for all $j \geq j_0$, where $\bar{c} = \max\{c_1, c_2, \dots, c_p\}$. \square

4.4. Whitney smoothness.

Proof of Theorem 1.2, part (C). The case $k = 1$ follows from Theorem 3.3.

Let $k \geq 2$ and pick $1 > \beta > \max\{\lambda^{-\frac{n}{k}}, \theta^{\frac{n}{k}}\}$. Let $\bar{x} \in \mathcal{N}_\beta$, let $\epsilon > 0$ be very small.

Once again, let n be the maximal number such that $c_j \notin [x; \bar{x}]$ for all $j \leq n$.

Then, similar to the proof of Theorem 4.5,

$$(16) \quad \epsilon^k > \lambda^{-n} \quad \text{and} \quad \epsilon^k > \theta^n$$

Since $\rho = \mathcal{L}^n(1) + O(\theta^n)$, for $0 \leq s \leq k-1$, Proposition 3.3 implies

$$\rho_s = \sum_{\substack{\text{finite } k \leq i_1, \dots, i_k < n \\ i_1 \geq 1, \dots, i_k \geq 1}} \mathcal{D}_{m_1, \dots, m_j}^{i_1, \dots, i_j}(h_{1,s}, \dots, h_{j-1,s}, \mathcal{L}^n(1)) + O(\lambda^{-n} + \theta^n).$$

To simplify the notation, let

$$\rho_{s,n} = \sum_{\substack{\text{finite } k \leq i_1, \dots, i_k < n \\ i_1 \geq 1, \dots, i_k \geq 1}} \mathcal{D}_{m_1, \dots, m_j}^{i_1, \dots, i_j}(h_{1,s}, \dots, h_{j-1,s}, \mathcal{L}^n(1)).$$

By definition of n and since $f \in C^{k+2}$, $\rho_{k-1,n}$ is C^2 in $B(\bar{x}, \epsilon) = \{y : |y - \bar{x}| < \epsilon\}$. Hence, if $x \in B(\bar{x}, \epsilon)$,

$$\begin{aligned} \rho_{k-1}(x) - \rho_{k-1}(\bar{x}) &= \rho_{k-1,n}(x) - \rho_{k-1,n}(\bar{x}) + O(\lambda^{-n} + \theta^n) \\ &= \int_{\bar{x}}^x \rho'_{k-1,n}(y) dy + O(\lambda^{-n} + \theta^n) \\ &= \int_{\bar{x}}^x \rho'_{k-1,n}(y) - \rho'_{k-1,n}(\bar{x}) dy + \rho_{k-1,n}(\bar{x})(x - \bar{x}) + O(\lambda^{-n} + \theta^n) \\ &= O(\epsilon^2) + \rho'_{k-1,n}(\bar{x})(x - \bar{x}) + O(\lambda^{-n} + \theta^n). \end{aligned}$$

By Proposition 3.3, $\rho'_{k-1,n}(\bar{x}) = \rho'_k(\bar{x}) + O(\lambda^{-n} + \theta^n)$. Thus

$$\rho_{k-1}(x) - \rho_{k-1}(\bar{x}) = \rho_k(\bar{x})(x - \bar{x}) + O(\epsilon(\lambda^{-n} + \theta^n)) + O(\lambda^{-n} + \theta^n) + O(\epsilon^2).$$

Inequalities (16) imply that

$$(17) \quad \rho_{k-1}(x) - \rho_{k-1}(\bar{x}) = \rho_k(\bar{x})(x - \bar{x}) + O(\epsilon^{k+1}) + O(\epsilon^k) + O(\epsilon^2) = \rho_k(\bar{x})(x - \bar{x}) + O(\epsilon^2).$$

Now, note that if $x \in B(\bar{x}, \epsilon)$, then

$$(18) \quad \begin{aligned} \rho_{k-2}(x) - \rho_{k-2}(\bar{x}) &= \rho_{k-2,n}(x) - \rho_{k-2,n}(\bar{x}) + O(\lambda^{-n} + \theta^n) \\ &= \int_{\bar{x}}^x \rho'_{k-2,n}(y) dy + O(\lambda^{-n} + \theta^n). \end{aligned}$$

By Proposition 3.3, $\rho'_{k-2,n}(y) = \rho_{k-1}(y) + O(\lambda^{-n} + \theta^n)$. Combining (18) with (17) and using that $\epsilon^{k+1} < \epsilon^k < \epsilon^3$ we get

$$\begin{aligned} \rho_{k-2}(x) - \rho_{k-2}(\bar{x}) &= \int_{\bar{x}}^x \rho_{k-1}(y) dy + O(\epsilon(\lambda^{-n} + \theta^n)) + O(\lambda^{-n} + \theta^n) \\ &= \int_{\bar{x}}^x \rho_{k-1}(\bar{x}) + \rho_k(\bar{x})(y - \bar{x}) dy + O(\epsilon^3) + O(\epsilon(\lambda^{-n} + \theta^n)) + O(\lambda^{-n} + \theta^n) \\ &= \rho_{k-1}(\bar{x})(x - \bar{x}) + \rho_k(\bar{x}) \frac{(x - \bar{x})^2}{2} + O(\epsilon^3). \end{aligned}$$

Continuing this recursive argument we get

$$\rho_s(x) - \rho_s(\bar{x}) = \left(\sum_{j=0}^{k-s-1} \rho_{k-j}(\bar{x}) \frac{(x - \bar{x})^{k-s-j}}{(k-s-j)!} \right) + O(\epsilon^{k-s+1})$$

for all $s = 0, \dots, k-1$. In particular, when $s = 0$, we have the desired result. \square

4.5. Nondifferentiability set. As we saw in Proposition 4.5, if the critical orbit does not approach a point x exponentially fast, then the density function ρ is differentiable at x . In this subsection, we obtain a partial converse to this statement that is, if the critical point does approach exponentially fast with sufficiently high exponent then we cannot have differentiability.

Definition 4.7. For $\beta < 1$, define

$$\mathcal{N}_\beta = \{\bar{x} : d(c_n, \bar{x}) \leq \beta^n \text{ for infinitely many } n\}.$$

Proposition 4.8. $\mathcal{HD}(\mathcal{N}_\beta) = 0$ where \mathcal{HD} denotes the Hausdorff dimension.

Proof. Define U_n as the ball centered at c_n of radius β^n . Given $\epsilon > 0$ let $n_0 \geq 1$ such that $\beta^{n_0} \leq \epsilon$. Then, $\{U_n\}_{n \geq n_0}$ is an ϵ -cover of \mathcal{N}_β .

Note that $|U_n| = 2\beta^n$. Hence, for any $s \geq 0$ we have that

$$\mathcal{H}_\epsilon^s(\mathcal{N}_\beta) \leq \sum_{n \geq n_0} |U_n|^s \leq \sum_{n \geq n_0} |U_n|^s = \frac{2\beta^{n_0 s}}{1 - \beta^s} < \infty.$$

Therefore $\mathcal{HD}(\mathcal{N}_\beta) = 0$. \square

Proposition 4.9. If $\{c_n\}$ is dense in some interval $I \subset [0, 1]$ then \mathcal{N}_β is uncountable for all $\beta < 1$.

We have already mentioned in Remark 1.3 the closure of $\{c_n\}$ contains an interval for a typical PEUM.

Proof. Define $L_n = [c_n - \beta^n, c_n + \beta^n]$.

Since $\{c_n\}$ is dense, there exists c_{n_1} such that L_{n_1} is strictly contained in I . Set $M_1 = L_{n_1}$.

Now, again using the density of $\{c_n\}$, there exist $c_{n_{(1,1)}} \in (c_{n_1} - \beta^{n_1}, c_{n_1})$ and $c_{n_{(1,2)}} \in (c_{n_1}, c_{n_1} + \beta^{n_1})$ such that $L_{n_{(1,1)}}$ and $L_{n_{(1,2)}}$ are strictly contained in $(c_{n_1} - \beta^{n_1}, c_{n_1})$ and $(c_{n_1}, c_{n_1} + \beta^{n_1})$ respectively. Set $M_2 = L_{n_{(1,1)}} \cup L_{n_{(1,2)}}$.

Continuing this procedure we inductively define M_n and set $M = \bigcap_{n \geq 1} M_n$. M is a Cantor set which is contained in \mathcal{N}_β . Since M is uncountable, so is \mathcal{N}_β . \square

Lemma 4.10. *If*

$$(19) \quad \beta(\max_x |f'(x)|) < 1$$

and $\bar{x} \in \mathcal{N}_\beta$ then ρ is non-differentiable at \bar{x}

Proof. Suppose ρ is differentiable at \bar{x} . Since $\bar{x} \in \mathcal{N}_\beta$, there exists a sequence n_j $d(\bar{x}, c_{n_j}) \leq \beta^{n_j}$. Without loss of generality, assume $\bar{x} < c_{n_j}$.

Let y_1 and y_2 be two arbitrary points such that

$$\bar{x} < y_1 < c_{n_j} < y_2 < c_{n_j} + \beta^{n_j}.$$

Since ρ is assumed to be differentiable at \bar{x} , we have that $|\rho(y_i) - \rho(\bar{x})| \leq M\beta^{n_j}$ for $i = 1, 2$ and hence

$$|\rho(y_1) - \rho(y_2)| \leq 2M\beta^{n_j}.$$

Accordingly

$$\frac{c}{(\max |f'|)^{n_j}} \leq |\alpha_{n_j}| = \lim_{y_1 \uparrow c_{n_j}, y_2 \downarrow c_{n_j}} |\rho(y_2) - \rho(y_1)| \leq 2M\beta^{n_j}$$

where the first inequality follows from Lemma 4.1. For large j this inequality is incompatible with (19). Hence ρ can not be differentiable at \bar{x} . \square

Parts (A) and (B) of Theorem 1.2 follows from Theorem 4.5, Proposition 4.9 and Lemma 4.10. Since part (C) was proven in subsection 4.4, the proof of Theorem 1.2 is complete.

REFERENCES

- [1] Baladi V. *Positive transfer operators and decay of correlations*, Advanced Ser. Nonlin. Dyn. **16** (2000) World Scientific, River Edge, NJ, x+314 pp.
- [2] Baladi V. *On the susceptibility function of piecewise expanding interval maps*, Comm. Math. Phys. **275** (2007) 839–859.
- [3] Baladi V., Smiana D. *Linear response formula for piecewise expanding unimodal maps*, Nonlinearity **21** (2008) 677–711.
- [4] Baladi V., Smiana D. *Smooth deformations of piecewise expanding unimodal maps*, Discrete Contin. Dyn. Syst. **23** (2009) 685–703.
- [5] Baladi V., Smiana D. *Alternative proofs of linear response for piecewise expanding unimodal maps*, Erg. Th. Dynam. Sys. **30** (2010) 1–20.
- [6] Fitzpatrick F., Royden H. L. *Real Analysis* (4) 2010, Prentice Hall.
- [7] Keller G., Liverani C. *Stability of the spectrum for transfer operators*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **28** (1999) 141–152.

- [8] Keller G., Liverani C. *Rare events, escape rates and quasistationarity: some exact formulae*, J. Stat. Phys. **135** (2009) 519–534.
- [9] Krzyzewski K. *Some results on expanding mappings*, Asterisque 50 (1977), 205-218
- [10] Lasota A., Yorke J. A. *On the existence of invariant measures for piecewise monotonic transformations*, Trans. AMS **186** (1973), 481–488.
- [11] Riesz F., Sz.-Nagy B. *Functional analysis*, 1990 (reprint of the 1955 original), Dover Books on Advanced Mathematics, Dover, New York.
- [12] Ruelle D. *Smooth dynamics and new theoretical ideas in nonequilibrium statistical mechanics*, J. Statist. Phys. **95** (1999) 393–468.
- [13] Ruelle D. *Application of hyperbolic dynamics to physics: some problems and conjectures*, Bull. Amer. Math. Soc. **41** (2004) 275–278.
- [14] Ruelle D. *Differentiating the a.c.i.m. of an interval map with respect to f* , Comm. Math. Phys. **258** (2005) 445–453.
- [15] Sacksteder R. *The measures invariant under an expanding map*, In Springer Lecture Notes in Maths 392 (1974).
- [16] Schnellmann, D. *Typical points for one-parameter families of piecewise expanding maps of the interval*, Discrete Contin. Dyn. Syst. 31 (2011), no.3, 877-911.
- [17] Schnellmann, D. *Law of iterated logarithm and invariance principle for one-parameter families of interval maps*, Arxiv, preprint, 2013.
- [18] Szewc B. *The Perron-Frobenius operator in spaces of smooth functions on an interval*, Ergodic Theory Dyn. Syst. 4, 613-643 (1984).
- [19] Viana M. *Lecture notes on attractors and physical measures*, IMCA Monogr. **8** (1999) Lima, iv+101 pp.