# MATH241 Fall 2023 Exam 3 (Justin W-G)

Name (Neatly):	
UID (Neatly):	

#### Instructions:

- 1. Please do all problems on the pages and in the spaces provided. This exam will be scanned into Gradescope and if your answers are not in the correct locations they will not be found or graded!
- 2. Only simplify Calculus 3 related calculations unless otherwise specified.

1. Write TRUE or FALSE in the box to the right. No justification is required. Unreadable or [10 pts] ambiguous answers will be marked as incorrect.

# Solution:

Statement	TRUE/FALSE
Every vertically simple region is also horizontally simple.	
In spherical, $x = \rho \sin \phi \cos \theta$ .	
We can never use a double integral to find volume.	
$\iint_R 1  dA$ yields volume.	
The parameterization of a surface requires two variables.	

2. The following integral can be evaluated without integrating but by understanding something about the shape and integrand:

$$\int_{0}^{4} \int_{0}^{4-x} 3 \, dy \, dx$$

(a) Describe the shape as best you can. Either words or a picture are acceptable. [5 pts] Solution:

(b) Give the simplified numerical value for the integral. **Solution:** 


[5 pts]

3. Given the following iterated integral:

$$\int_0^{\pi/2} \int_0^{2\cos\theta} 2\,dr\,d\theta$$

Note: Don't worry about the fact that there's no "polar r" there. Not a typo.

(a) Draw the region *R* corresponding to the integral. All you need to do is draw the region. [5 pts] **Solution:** 



(b) Evaluate the integral. Solution:

[10 pts]

4. Suppose R is the region above  $y = x^2$ , below y = 9, and in the first quadrant. Set up the iterated [10 pts] double integral in rectangular coordinates treating R as horizontally simple for:

$$\iint_R xy \, dA$$

### DO NOT EVALUATE!

**Note:** A picture is not necessary but can earn partial credit. **Solution:** 

5. Suppose R is the region inside  $r = 3\cos\theta$  and outside  $r = 1 + \cos\theta$ . Set up the iterated double [10 pts] integral in polar coordinates for:

$$\iint_R x \, dA$$

### DO NOT EVALUATE!

**Note:** A picture is not necessary but can earn partial credit. **Solution:** 

6. Suppose D is the solid between  $r = \sin \theta$  and  $r = 2 \sin \theta$  and between z = 1 and  $z = 3 + x^2 + y^2$ . [15 pts] Write down the iterated triple integral in cylindrical coordinates for the volume of D.

## DO NOT EVALUATE!

Note: Pictures are not necessary but can earn partial credit.

7. Suppose D is the solid inside  $z = \sqrt{3x^2 + 3y^2}$ , below z = 10 and outside  $x^2 + y^2 + z^2 = 2$ . If the [15 pts] density is given by  $f(x, y, z) = z^4$ , write down the iterated triple integral in spherical coordinates for the mass of D.

## DO NOT EVALUATE!

Note: Pictures are not necessary but can earn partial credit.

8. Write down a parameterization of the part of the plane y = 4 - x in the first octant and between [5 pts] z = 0 and z = 7. No sketch is required.

Solution:

9. Sketch the surface parameterized by:

[5 pts]

$$\bar{\boldsymbol{r}}(x,y) = x\,\hat{\boldsymbol{\imath}} + y\,\hat{\boldsymbol{\jmath}} + (9 - y^2)\,\hat{\boldsymbol{k}}$$
$$0 \le x \le 4$$
$$0 \le y \le 3$$

10. Let R be the region bounded by the lines y = 1,  $y = \frac{1}{4}x$ , and x - 3y = 2. Consider the integral: [15 pts]

$$\iint_R \frac{y}{x - 3y} \, dA$$

Use the substitution x = 3u + v and y = u to convert this integral to an iterated integral in the *uv*-plane.

### DO NOT EVALUATE!

Note: Pictures are not necessary but can earn partial credit.