

MATH241 Exam 4 Spring 2022 (Justin Wyss-Gallifent)
May the 4th be with you!

In the box below NEATLY enter your name as it appears in Gradescope:

NAME:

In the box below NEATLY enter your UID:

UID:

Directions: Please do work in the spaces provided and do not spill over to other pages - the exams will be scanned into Gradescope and auto-tagged this way. Show work and use only methods taught in this course and show work as indicated. No calculators or other devices permitted. Numerical answers do not need to be simplified. Good luck!

Extra Credit: Write the date, time, building and room of your final exam.

[\leq 2 pts]

Solution:

1. Write the word TRUE or FALSE in the box to the right. No justification is required. [10 pts]

(a) If C is closed and \mathbf{F} is conservative then $\int_C \mathbf{F} \cdot d\mathbf{r} \neq 0$.

(b) If \mathbf{F} is conservative then $\nabla \times \mathbf{F} = \mathbf{0}$.

(c) Green's Theorem applies if C is not closed.

(d) We only have one way to evaluate $\int_C f ds$.

(e) The expression $\nabla \times f(x, y, z)$ makes sense.

2. Let C be the curve parametrized by $\mathbf{r}(t) = 3 \cos t \hat{\mathbf{i}} + 4 \sin t \hat{\mathbf{j}}$ for $0 \leq t \leq \frac{\pi}{2}$. Without [10 pts]
integrating, explain how you know that $\int_C 1 ds > 5$.

Hint: Sketch it!

Solution:

3. Let Σ be the part of the plane $2x + y + z = 4$ in the first octant. If the mass density at a point [20 pts] is given by $f(x, y, z) = x$, write down and evaluate the integral for the mass of Σ .

You Should Evaluate Your Result!

Solution:

4. Suppose $f(x, y, z) = \frac{x}{y} - xz^2$ and C is the straight line segment from $(1, 2, 3)$ to $(8, 10, 100)$. [5 pts]
Evaluate the line integral:

$$\int_C (\nabla f(x, y, z)) \cdot d\mathbf{r}$$

You Should Evaluate Your Result!

Solution:

5. Use Green's Theorem to evaluate the line integral $\int_C 4y dx + 6x dy$ where C is the curve [15 pts]
parametrized by $\mathbf{r}(t) = (1 + \cos t)\hat{i} + (2 + \sin t)\hat{j}$ for $0 \leq t \leq 2\pi$.

You Should Evaluate Your Result!

Solution:

6. Let Σ be the part of $z = x^2 + y^2$ inside $z = 8 - x^2 - y^2$ as well as the part of $z = 8 - x^2 - y^2$ inside $z = x^2 + y^2$, oriented outwards. Apply the Divergence Theorem to construct an iterated triple integral in cylindrical coordinates for the following: [15 pts]

$$\iint_{\Sigma} (xy \hat{i} + y \hat{j} + xz \hat{k}) \cdot \mathbf{n} \, dS$$

You Should Not Evaluate Your Result!

Solution:

7. Let Σ be the part of the vertical plane $x + 2y = 4$ in the first octant and below $z = 5$. Let C [25 pts]
be the edge of Σ with clockwise orientation when viewed from the first octant looking towards
the origin. Consider the integral:

$$\int_C z^2 dx + x dy + z dz$$

Use Stokes' Theorem to rewrite the line integral as a surface integral and then as an iterated
double integral.

You Should Not Evaluate Your Result!

Solution: