

AMSC 614
NUMERICAL METHODS FOR STATIONARY PDEs
 HOMEWORK # 4 (Pbs 1-3 due Nov 13, Pbs 4-5 due Nov 18)

1. *Bogner-Fox-Schmit rectangle*: Let R be a rectangle with vertices $\{\mathbf{x}_i\}_{i=1}^4$ in \mathbb{R}^2 .

(a) Show that the following nodal variables determine $Q_3(R)$, i.e. that the corresponding set \mathcal{N} is unisolvent:

$$p(\mathbf{x}_i), \quad \partial_1 p(\mathbf{x}_i), \quad \partial_2 p(\mathbf{x}_i), \quad \partial_{12}^2 p(\mathbf{x}_i) \quad \forall 1 \leq i \leq 4.$$

(b) Show that the corresponding finite element space \mathbb{V}_h satisfies $\mathbb{V}_h \subset C^1(\bar{\Omega}) \cap H^2(\Omega)$.

2. *Raviart-Thomas element (of lowest order)*: This problem illustrates how to design finite elements for the space $H(\text{div}; \Omega)$ where Ω is a polygonal domain in \mathbb{R}^2 .

(a) $H(\text{div}; \Omega)$ is the space of vector fields \mathbf{p} in Ω such that $\mathbf{p} \in [L^2(\Omega)]^2$ and weak divergence $\text{div } \mathbf{p} \in L^2(\Omega)$. Show that $H(\text{div}; \Omega)$ is a Hilbert space with the inner product $\langle \mathbf{p}, \mathbf{q} \rangle := \int_{\Omega} \mathbf{p} \cdot \mathbf{q} + \text{div } \mathbf{p} \text{ div } \mathbf{q}$.

(b) Consider the following space \mathcal{P} of vector-valued polynomials over a triangle T in Ω :

$$\mathcal{P} = \mathbb{P}_0(T)^2 + \mathbf{x} \mathbb{P}_0(T).$$

Hence a function $\mathbf{p} \in \mathcal{P}$ is of the form $\mathbf{p}(\mathbf{x}) = \mathbf{a} + b\mathbf{x}$ with $\mathbf{a} \in \mathbb{R}^2$ and $b \in \mathbb{R}$ constants. Consider the following nodal variables for each side S of T :

$$N_S(\mathbf{p}) = \int_S \mathbf{p} \cdot \nu_S$$

where ν_S is the unit normal to S . Prove that the set \mathcal{N} of nodal variables is unisolvent. To this end show that the product $\mathbf{p} \cdot \nu_S$ is constant for all sides S of T .

(c) Prove that all functions \mathbf{p} in the finite element space resulting from pasting together affine equivalent triangles are in $H(\text{div}; \Omega)$. Note however that \mathbf{p} is discontinuous across interelement boundaries. Hint: show that the normal components of discrete vector fields are continuous across interelement boundaries and that this implies the assertion.

3. *Dual basis*: Consider a simplex T in \mathbb{R}^d and let $\mathcal{N}_1(T) = \{N_i\}_{i=0}^d \subset \mathbb{P}_1^*(T)$ be the Lagrange nodal variables (or nodal evaluation). By the Riesz representation theorem, there exist functions $\lambda_i^* \in \mathbb{P}_1(T)$ for each $0 \leq i \leq d$ such that

$$N_j(\phi_i) = \int_T \lambda_i \lambda_j^* = \delta_{ij}.$$

Show that

$$\lambda_i^* = \frac{(1+d)^2}{|T|} \lambda_i - \frac{1+d}{|T|} \sum_{j \neq i} \lambda_j \quad \forall 0 \leq i \leq d.$$

4. Use the MATLAB code `fem` to solve the following two problems on the L-shaped domain $\Omega = [-1, 1]^2 \setminus [0, 1] \times [0, -1]$ of \mathbb{R}^2 with exact solutions:

- *Smooth solution*: $u(x, y) = \cos(\pi x) \sin(\pi y)$, in cartesian coordinates;
- *Nonsmooth Solution*: $u(x, y) = r^{2/3} \sin(2\theta/3)$, in polar coordinates (r, θ) .

Assume Dirichlet condition $g_D = u$ on the entire boundary $\partial\Omega$ and $f = -\Delta u$.

(a) Read the tutorial by P. Morin about the implementation of the FEM for \mathbb{P}_1 Lagrange elements (see the website).

- (b) Generate the data files `vertex_coordinates.txt`, `elem_vertices.txt`, and `dirichlet.txt` using `gen_mesh_L_shape.m` for uniform refinement with meshsize $h = \frac{1}{N} = 2^{-k}$ and $k = 2, 3, 4, 5, 6, 7$. Find the corresponding solutions $U_{\mathcal{T}} = u_h$.
- (c) Show that the stiffness matrices for these meshes and those for finite differences with a 5-point stencil coincide. To this end consider a generic interior star.
- (d) Find the errors $|u - u_h|_{H_0^1(\Omega)}$ and $\|u - u_h\|_{L^2(\Omega)}$, and plot them vs the number of degrees of freedom N in a log-log plot. Explain the behavior $\|u - u_h\| \approx CN^{-\alpha}$ that you observe and find α . Relate this to the regularity of u and HW#3-Pb#2 about polynomial interpolation.

5. *The MINI element (of Arnold and Brezzi)*: This is an element for the Stokes problem. Let \mathbb{Q}_h be the space of continuous piecewise linear elements with zero mean; this is the space for pressure. Let \mathbb{V}_h be the space of vector-valued continuous piecewise polynomials \mathbf{v}_h of the form

$$\mathbf{w}_T + b_T \mathbf{c}_T \quad \forall T \in \mathcal{T},$$

where \mathbf{w}_T is linear in T , \mathbf{c}_T is constant, and b_T is the cubic bubble in T ; this is the space for velocity. Show that the pair $(\mathbb{V}_h, \mathbb{Q}_h)$ satisfies the discrete inf-sup property

$$\beta \|q_h\|_{L^2(\Omega)} \leq \inf_{\mathbf{v}_h \in \mathbb{V}_h} \frac{\int_{\Omega} q_h \operatorname{div} \mathbf{v}_h}{\|\mathbf{v}_h\|_{H_0^1(\Omega)}} \quad \forall q_h \in \mathbb{Q}_h$$

with $\beta > 0$ independent of h . Hint: Let $\mathbf{v} \in H_0^1(\Omega)$ be a function that satisfies the continuous inf-sup property for q_h . To discretize \mathbf{v} proceed as follows. First let $\mathbf{w}_h = I_h \mathbf{v}$ be an interpolant of \mathbf{v} with values in the space of continuous piecewise linears which is stable in $H_0^1(\Omega)$, namely

$$\|\mathbf{w}_h\|_{H_0^1(\Omega)} \leq \alpha \|\mathbf{v}\|_{H_0^1(\Omega)};$$

we will see two such interpolants. Choose the constant \mathbf{c}_T for each $T \in \mathcal{T}$ upon imposing the condition

$$\int_{\Omega} q_h \operatorname{div} (\mathbf{v} - \mathbf{v}_h) = 0.$$

To this end, integrate by parts $\sum_{T \in \mathcal{T}} \int_T q_h \operatorname{div} \mathbf{v}_h$ elementwise and note that the boundary terms vanish.