

NUMERICAL METHODS FOR STATIONARY PDEs

HOMEWORK # 3 (Pbs 1-3 due Th Oct 25, Pbs 4-5 due Tu Oct 30)

1 (20 pts). *Approximation with smooth functions*: Let $u \in W_p^1(\Omega)$ with $1 \leq p \leq \infty$ and dimension $d \geq 1$. Let $\rho \in C_0^\infty(\mathbb{R}^d)$ be a mollifier with properties:

$$\text{supp } \rho = B_1, \quad \rho \geq 0, \quad \int_{B_1} \rho = 1,$$

where B_1 is the unit ball centered at the origin. Let $\rho_\varepsilon(x) = \varepsilon^{-d} \rho(\varepsilon^{-1}x)$ and u_ε be defined as

$$u_\varepsilon(x) = \int_{\Omega} u(x-y) \rho_\varepsilon(y) dy \quad \forall x \in K,$$

where K is a compact set of Ω and ε is sufficiently small. Show the error estimate

$$\|u - u_\varepsilon\|_{L^p(K)} \leq C\varepsilon |u|_{W_p^1(\Omega)}.$$

2 (20 pts). *Equivalent norms (Deni-Lions)*: Consider the Sobolev space $W_p^{k+1}(\Omega)$ with $k \geq 0, 1 \leq p \leq \infty$ and a Lipschitz domain Ω in \mathbf{R}^d . Let $\{f_i\}_{i=1}^N$ be linear continuous functionals in $W_p^{k+1}(\Omega)$ such that for any polynomial $v \in \mathbb{P}_k$ of degree $\leq k$:

$$f_i(v) = 0 \quad \forall 1 \leq i \leq N \quad \iff \quad v = 0.$$

(a) Show that $\|v\|_{W_p^{k+1}(\Omega)}$ is equivalent to the seminorm

$$|v|_{W_p^{k+1}(\Omega)} + \sum_{i=1}^N |f_i(v)|.$$

Hint: Proceed by contradiction assuming that there is a sequence $\{v_n\} \subset W_p^{k+1}(\Omega)$ such that $\|v_n\|_{W_p^{k+1}(\Omega)} = 1$ but the latter seminorm tends to 0. Use that $W_p^{k+1}(\Omega)$ is compactly imbedded in $W_p^k(\Omega)$ (Rellich Theorem), namely that each bounded sequence in $W_p^{k+1}(\Omega)$ admits a convergence subsequence in $W_p^k(\Omega)$.

(b) Show the polynomial interpolation bound

$$\inf_{q \in \mathbb{P}_k} \|v - q\|_{W_p^{k+1}(\Omega)} \leq C(\Omega) |v|_{W_p^{k+1}(\Omega)} \quad \forall v \in W_p^{k+1}(\Omega).$$

3 (20 pts). *Nonhomogeneous Dirichlet Problem*: Given a bounded Lipschitz domain Ω in \mathbf{R}^n , and $g \in H^1(\Omega), f \in H^{-1}(\Omega)$, set

$$L(v) = \langle f, v \rangle - \int_{\Omega} \nabla g \nabla v \quad \forall v \in H^1(\Omega).$$

(a) Prove that there exists a unique solution to the variational problem

$$z \in H_0^1(\Omega) : \quad \int_{\Omega} \nabla z \nabla v = L(v) \quad \forall v \in H_0^1(\Omega).$$

(b) Show that such a problem is equivalent to the minimization of $J(v) = \int_{\Omega} \frac{1}{2} |\nabla v|^2 - \langle f, v \rangle$ over the subspace $V = \{v \in H^1(\Omega) : v - g \in H_0^1(\Omega)\}$.

(c) Prove that $u = z + g$ formally solves $-\Delta u = f$ in Ω with boundary condition $u = g$ on $\partial\Omega$.

4 (20 pts). *Third boundary value problem*: Given $f \in L^2(\Omega), g \in H^2(\Omega)$ and $0 < P_1 \leq p \leq P_2$ on $\partial\Omega$, consider the Robin problem

$$-\Delta u = f \quad \text{in } \Omega, \quad \partial_\nu u + p(u - g) = 0 \quad \text{on } \partial\Omega,$$

- (a) Find a variational formulation which amounts to solving this problem.
- (b) Show that Lax-Milgram theorem applies and conclude that there exists a unique solution $u \in H^1(\Omega)$. To this end, show that the bilinear form is coercive in $H^1(\Omega)$.
- (c) Suppose that $p = \epsilon^{-1} \rightarrow \infty$ and denote the corresponding solution by u_ϵ . Determine the boundary value problem satisfied by $u_0 = \lim_{\epsilon \rightarrow 0} u_\epsilon$.
- (d) Derive an error estimate for $\|u_\epsilon - u_0\|_{H^1(\Omega)}$.

5 (20 pts). *Darcy's flow*. Let u be the pressure and $\sigma = -K\nabla u$ be the flux of the model problem for flow in porous media, which can be written as

$$K^{-1}\sigma + \nabla u = 0, \quad \operatorname{div} \sigma = f.$$

- (a) Let $\mathbb{V} = H_0(\operatorname{div}; \Omega) := \{\tau \in [L^2(\Omega)]^d : \operatorname{div} \tau \in L^2(\Omega), \tau \cdot \nu = 0 \text{ on } \partial\Omega\}$ and $Q = L_0^2(\Omega)$. Write a variational formulation for this problem, and show that the inf-sup condition is satisfied.
- (b) Deduce existence, uniqueness, and stability of the solution pair (u, σ) .