

Lecture 2b: Phase Transition in Random Graphs

Radu Balan

Supplemental Material - 2023

Analytical Results

Distributions

Today we discuss about phase transition in random graphs. Recall on the *Erdős-Rényi class* $\mathcal{G}_{n,p}$ of random graphs, the probability mass function on \mathcal{G} , $P : \mathcal{G} \rightarrow [0, 1]$, is obtained by assuming that, as random variables, edges are independent from one another, and each edge occurs with probability $p \in [0, 1]$. Thus a graph $G \in \mathcal{G}$ with m vertices will have probability $P(G)$ given by

$$P(G) = p^m(1 - p)^{\binom{n}{2}}.$$

Analytical Results

Distributions

Today we discuss about phase transition in random graphs. Recall on the *Erdős-Rényi class* $\mathcal{G}_{n,p}$ of random graphs, the probability mass function on \mathcal{G} , $P : \mathcal{G} \rightarrow [0, 1]$, is obtained by assuming that, as random variables, edges are independent from one another, and each edge occurs with probability $p \in [0, 1]$. Thus a graph $G \in \mathcal{G}$ with m vertices will have probability $P(G)$ given by

$$P(G) = p^m (1 - p)^{\binom{n}{2} - m}.$$

Recall the expected number of q -cliques X_q is

$$\mathbb{E}[X_q] = \binom{n}{q} p^{q(q-1)/2}$$

Analytical Results

Distributions

We shall also use $\Gamma^{n,m}$ the set of all graphs on n vertices with m edges.
The set $\Gamma^{n,m}$ has cardinal

$$\binom{\binom{n}{2}}{m}.$$

In $\Gamma^{n,m}$ each graph is equally probable.

Analytical Results

Cliques

The case of 3-cliques: $\mathbb{E}[X_3] = \theta n^3 p^3$ ($\theta \sim \frac{1}{6}$).

The case of 4-cliques: $\mathbb{E}[X_4] = \theta n^4 p^6$ ($\theta \sim \frac{1}{24}$).

The first problem we consider is the size of the largest clique of a random graph.

Note, finding the size of the largest clique (called *the clique number*) is a NP-hard problem.

Analytical Results

Cliques

The case of 3-cliques: $\mathbb{E}[X_3] = \theta n^3 p^3$ ($\theta \sim \frac{1}{6}$).

The case of 4-cliques: $\mathbb{E}[X_4] = \theta n^4 p^6$ ($\theta \sim \frac{1}{24}$).

The first problem we consider is the size of the largest clique of a random graph.

Note, finding the size of the largest clique (called *the clique number*) is a NP-hard problem.

Idea: Analyze p so that $\mathbb{E}[X_q] \approx 1$.

- For $p > \frac{1}{n}$ and large n we expect that graphs will have a 3-clique;
- For $p > \frac{1}{n^{2/3}}$ and large n , we expect that graphs will have a 4-clique;

Analytical Results

Cliques

The case of 3-cliques: $\mathbb{E}[X_3] = \theta n^3 p^3$ ($\theta \sim \frac{1}{6}$).

The case of 4-cliques: $\mathbb{E}[X_4] = \theta n^4 p^6$ ($\theta \sim \frac{1}{24}$).

The first problem we consider is the size of the largest clique of a random graph.

Note, finding the size of the largest clique (called *the clique number*) is a NP-hard problem.

Idea: Analyze p so that $\mathbb{E}[X_q] \approx 1$.

- For $p > \frac{1}{n}$ and large n we expect that graphs will have a 3-clique;
- For $p > \frac{1}{n^{2/3}}$ and large n , we expect that graphs will have a 4-clique;

Question: How sharp are these thresholds?

Analytical Results

3-Cliques

Theorem

Let $p = p(n)$ be the edge probability in $\mathcal{G}_{n,p}$.

- 1 If $p \gg \frac{1}{n}$ (i.e. $\lim_{n \rightarrow \infty} np = \infty$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 3-clique}] \rightarrow 1.$
- 2 If $p \ll \frac{1}{n}$ (i.e. $\lim_{n \rightarrow \infty} np = 0$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 3-clique}] \rightarrow 0.$

Analytical Results

3-Cliques

Theorem

Let $p = p(n)$ be the edge probability in $\mathcal{G}_{n,p}$.

- ① If $p \gg \frac{1}{n}$ (i.e. $\lim_{n \rightarrow \infty} np = \infty$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 3-clique}] \rightarrow 1.$
- ② If $p \ll \frac{1}{n}$ (i.e. $\lim_{n \rightarrow \infty} np = 0$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 3-clique}] \rightarrow 0.$

Theorem

Let $m = m(n)$ be the number of edges in $\Gamma^{n,m}$.

- ① If $m \gg n$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n} = \infty$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 3-clique}] \rightarrow 1.$
- ② If $m \ll n$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n} = 0$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 3-clique}] \rightarrow 0.$

Analytical Results

4-Cliques

Theorem

Let $p = p(n)$ be the edge probability in $\mathcal{G}_{n,p}$.

- 1 If $p \gg \frac{1}{n^{2/3}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/3}p = \infty$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 4-clique}] \rightarrow 1$.
- 2 If $p \ll \frac{1}{n^{2/3}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/3}p = 0$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 4-clique}] \rightarrow 0$.

Analytical Results

4-Cliques

Theorem

Let $p = p(n)$ be the edge probability in $\mathcal{G}_{n,p}$.

- 1 If $p \gg \frac{1}{n^{2/3}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/3}p = \infty$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 4-clique}] \rightarrow 1$.
- 2 If $p \ll \frac{1}{n^{2/3}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/3}p = 0$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a 4-clique}] \rightarrow 0$.

Theorem

Let $m = m(n)$ be the number of edges in $\Gamma^{n,m}$.

- 1 If $m \gg n^{4/3}$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n^{4/3}} = \infty$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 4-clique}] \rightarrow 1$.
- 2 If $m \ll n^{4/3}$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n^{4/3}} = 0$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 4-clique}] \rightarrow 0$.

Analytical Results

q -Cliques

Theorem

Let $p = p(n)$ be the edge probability in $\mathcal{G}_{n,p}$. Let $q \geq 3$ be an integer.

- 1 If $p \gg \frac{1}{n^{2/(q-1)}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/(q-1)} p = \infty$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a } q\text{-clique}] \rightarrow 1$.
- 2 If $p \ll \frac{1}{n^{2/(q-1)}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/(q-1)} p = 0$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a } q\text{-clique}] \rightarrow 0$.

Analytical Results

q -Cliques

Theorem

Let $p = p(n)$ be the edge probability in $\mathcal{G}_{n,p}$. Let $q \geq 3$ be and integer.

- 1 If $p \gg \frac{1}{n^{2/(q-1)}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/(q-1)} p = \infty$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a } q - \text{clique}] \rightarrow 1$.
- 2 If $p \ll \frac{1}{n^{2/(q-1)}}$ (i.e. $\lim_{n \rightarrow \infty} n^{2/(q-1)} p = 0$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \mathcal{G}_{n,p} \text{ has a } q - \text{clique}] \rightarrow 0$.

Theorem

Let $m = m(n)$ be the number of edges in $\Gamma^{n,m}$. Let $q \geq 3$ be and integer.

- 1 If $m \gg n^{2(q-2)/(q-1)}$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n^{2(q-2)/(q-1)}} = \infty$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a } q - \text{clique}] \rightarrow 1$.
- 2 If $m \ll n^{2(q-2)/(q-1)}$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n^{2(q-2)/(q-1)}} = 0$) then $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a } q - \text{clique}] \rightarrow 0$.

Analytical Results

Markov and Chebyshev Inequalities

We want to control probabilities of the random event $X_3(G) > 0$. Two important tools:

- 1 (Markov's Inequality) Assume X is a non-negative random variable. Then $Prob[X \geq t] \leq \frac{\mathbb{E}[X]}{t}$.
- 2 (Chebyshev's Inequality) For any random variable X , $Prob[|X - E[X]| \geq t] \leq \frac{Var[X]}{t^2}$.

where $\mathbb{E}[X]$ is the mean of X , and $Var[X] = \mathbb{E}[X^2] - |\mathbb{E}[X]|^2$ is the variance of X .

Analytical Results

Markov and Chebyshev Inequalities

We want to control probabilities of the random event $X_3(G) > 0$. Two important tools:

- 1 (Markov's Inequality) Assume X is a non-negative random variable. Then $\text{Prob}[X \geq t] \leq \frac{\mathbb{E}[X]}{t}$.
- 2 (Chebyshev's Inequality) For any random variable X , $\text{Prob}[|X - \mathbb{E}[X]| \geq t] \leq \frac{\text{Var}[X]}{t^2}$.

where $\mathbb{E}[X]$ is the mean of X , and $\text{Var}[X] = \mathbb{E}[X^2] - |\mathbb{E}[X]|^2$ is the variance of X . *Quick Proof:*

$$\text{Prob}[X \geq t] = \int_t^\infty p_X(x) dx \leq \frac{1}{t} \int_t^\infty x p_X(x) dx \leq \frac{\mathbb{E}[X]}{t}.$$

$$\text{Prob}[|X - \mathbb{E}[X]| \geq t] = P[|X - \mathbb{E}[X]|^2 \geq t^2] \leq \frac{\mathbb{E}[|X - \mathbb{E}[X]|^2]}{t^2} = \frac{\text{Var}[X]}{t^2}.$$

Analytical Results

Proofs for the 3-clique case

For small probability: We shall use Markov's inequality to show

$Prob[X_3 > 0] \rightarrow 0$ when $p \ll \frac{1}{n}$:

$$Prob[X_3 > 0] = Prob[X_3 \geq 1] \leq \frac{E[X_3]}{1} = \frac{n(n-1)(n-2)}{6} p^3 = \theta n^3 p^3 \rightarrow 0.$$

Analytical Results

Proofs for the 3-clique case

For small probability: We shall use Markov's inequality to show

$Prob[X_3 > 0] \rightarrow 0$ when $p \ll \frac{1}{n}$:

$$Prob[X_3 > 0] = Prob[X_3 \geq 1] \leq \frac{E[X_3]}{1} = \frac{n(n-1)(n-2)}{6} p^3 = \theta n^3 p^3 \rightarrow 0.$$

For large probability: Since $\mathbb{E}[X_3] \rightarrow \infty$ it follows that $Prob[X_3 > 0] > 0$.

We need to show that $Prob[X_3 = 0] \rightarrow 0$. By Chebyshev's inequality:

$$Prob[X_3 = 0] \leq Prob[|X_3 - \mathbb{E}[X_3]| \geq \mathbb{E}[X_3]] \leq \frac{Var[X_3]}{|\mathbb{E}[X_3]|^2}$$

Analytical Results

Proofs for the 3-clique case

For small probability: We shall use Markov's inequality to show

$Prob[X_3 > 0] \rightarrow 0$ when $p \ll \frac{1}{n}$:

$$Prob[X_3 > 0] = Prob[X_3 \geq 1] \leq \frac{E[X_3]}{1} = \frac{n(n-1)(n-2)}{6} p^3 = \theta n^3 p^3 \rightarrow 0.$$

For large probability: Since $\mathbb{E}[X_3] \rightarrow \infty$ it follows that $Prob[X_3 > 0] > 0$.

We need to show that $Prob[X_3 = 0] \rightarrow 0$. By Chebyshev's inequality:

$$Prob[X_3 = 0] \leq Prob[|X_3 - \mathbb{E}[X_3]| \geq \mathbb{E}[X_3]] \leq \frac{Var[X_3]}{|\mathbb{E}[X_3]|^2}$$

Need the variance of $X_3 = \sum_{(i,j,k) \in S_3} 1_{i,j,k}$,

$$X_3^2 = \sum_{(i,j,k) \in S_3} \sum_{(i',j',k') \in S_3} 1_{i,j,k} 1_{i',j',k'}.$$

Analytical Results

Proofs for the 3-clique case

$$\begin{aligned}
 X_3^2 = & \sum_{(i,j,k) \in \mathcal{S}_3(n)} 1_{i,j,k} + \sum_{(i,j,k) \in \mathcal{S}_3(n)} \sum_{l \in \mathcal{S}_1(n-3)} (1_{i,j,k} 1_{i,j,l} + 1_{i,j,k} 1_{j,k,l} + 1_{i,j,k} 1_{k,i,l}) + \\
 & + \sum_{(i,j,k) \in \mathcal{S}_3(n)} \sum_{u,v \in \mathcal{S}_2(n-3)} (1_{i,j,k} 1_{i,u,v} + 1_{i,j,k} 1_{j,u,v} + 1_{i,j,k} 1_{k,u,v}) + \\
 & + \sum_{(i,j,k) \in \mathcal{S}_3(n)} \sum_{(i',j',k') \in \mathcal{S}_3(n-3)} 1_{i,j,k} 1_{i',j',k'}
 \end{aligned}$$

Analytical Results

Proofs for the 3-clique case

$$\begin{aligned}
 X_3^2 = & \sum_{(i,j,k) \in S_3(n)} 1_{i,j,k} + \sum_{(i,j,k) \in S_3(n)} \sum_{l \in S_1(n-3)} (1_{i,j,k} 1_{i,j,l} + 1_{i,j,k} 1_{j,k,l} + 1_{i,j,k} 1_{k,i,l}) + \\
 & + \sum_{(i,j,k) \in S_3(n)} \sum_{u,v \in S_2(n-3)} (1_{i,j,k} 1_{i,u,v} + 1_{i,j,k} 1_{j,u,v} + 1_{i,j,k} 1_{k,u,v}) + \\
 & + \sum_{(i,j,k) \in S_3(n)} \sum_{(i',j',k') \in S_3(n-3)} 1_{i,j,k} 1_{i',j',k'}
 \end{aligned}$$

$$\mathbb{E}[X_3^2] = |S_3| p^3 + 3|S_3|(n-3)p^5 + 3|S_3| \binom{n-3}{2} p^6 + |S_3| \binom{n-3}{3} p^6.$$

Thus

$$\text{Var}[X_3] = \mathbb{E}[X_3^2] - |\mathbb{E}[X_3]|^2 = \dots = \theta(n^3 p^3 + n^4 p^5 + n^5 p^6).$$

Analytical Results

Proofs for the 3-clique case

and:

$$\text{Prob}[X_3 = 0] \leq \frac{\theta(n^3 p^3 + n^4 p^5 + n^5 p^6)}{\theta(n^6 p^6)} = \frac{1}{(np)^3} + \frac{1}{n} \rightarrow 0.$$

Analytical Results

Proofs for the 3-clique case

and:

$$\text{Prob}[X_3 = 0] \leq \frac{\theta(n^3 p^3 + n^4 p^5 + n^5 p^6)}{\theta(n^6 p^6)} = \frac{1}{(np)^3} + \frac{1}{n} \rightarrow 0.$$

Similar proofs for the other cases (4-cliques and q -cliques).

Analytical Results

Behavior at the threshold

In general we obtain a "coarse threshold". Recall a Poisson random variable X with parameter λ has p.m.f. $\text{Prob}[X = k] = e^{-\lambda} \frac{\lambda^k}{k!}$.

Theorem

In $\mathcal{G}_{n,p}$,

- ① For $p = \frac{c}{n}$, X_3 is asymptotically Poisson with parameter $\lambda = c^3/6$.
- ② For $p = \frac{c}{n^{2/3}}$, X_4 is asymptotically Poisson with parameter $\lambda = c^6/24$.

Analytical Results

Behavior at the threshold

In general we obtain a "coarse threshold". Recall a Poisson random variable X with parameter λ has p.m.f. $Prob[X = k] = e^{-\lambda} \frac{\lambda^k}{k!}$.

Theorem

In $\mathcal{G}_{n,p}$,

- ① For $p = \frac{c}{n}$, X_3 is asymptotically Poisson with parameter $\lambda = c^3/6$.
- ② For $p = \frac{c}{n^{2/3}}$, X_4 is asymptotically Poisson with parameter $\lambda = c^6/24$.

Theorem

In $\Gamma^{n,m}$,

- ① For $m = cn$, X_3 is asymptotically Poisson with parameter $\lambda = 4c^3/3$.
- ② For $m = cn^{4/3}$, X_4 is asymptotically Poisson with parameter $\lambda = 8c^6/3$.

Analytical Results

Connected Components

$\mathcal{G}_{n,p}$ class of random graphs has a remarkable property in regards to the largest connected component. We shall express the result in the class $\Gamma^{n,m}$.

Analytical Results

Connected Components

Theorem

- ① Let $m = m(n)$ satisfies $m \ll \frac{1}{2}n \log(n)$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = 0$$

- ② Let $m = m(n)$ satisfies $m \gg \frac{1}{2}n \log(n)$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = 1$$

Analytical Results

Connected Components

Theorem

- ① Let $m = m(n)$ satisfies $m \ll \frac{1}{2}n \log(n)$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = 0$$

- ② Let $m = m(n)$ satisfies $m \gg \frac{1}{2}n \log(n)$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = 1$$

- ③ Assume $m = \frac{1}{2}n \log(n) + tn + o(n)$, where $o(n) \ll n$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = e^{-e^{-2t}}$$

Analytical Results

Connected Components

Theorem

- ① Let $m = m(n)$ satisfies $m \ll \frac{1}{2}n \log(n)$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = 0$$

- ② Let $m = m(n)$ satisfies $m \gg \frac{1}{2}n \log(n)$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = 1$$

- ③ Assume $m = \frac{1}{2}n \log(n) + tn + o(n)$, where $o(n) \ll n$. Then

$$\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ is connected}] = e^{-e^{-2t}}$$

In this case $\frac{1}{2}n \log(n)$ is known as a *strong threshold*. 

Numerical Results


3-cliques & Connectivity results

Results for $n = 1000$ vertices.

- 1 3-cliques. Recall $\mathbb{E}[X_3] \sim m^3$
- 2 Connectivity. Recall the connectivity threshold is $\frac{1}{2}n \log(n) = 3454$.


Numerical Results

3-cliques




Numerical Results

3-cliques




Numerical Results

Connectivity




Numerical Results

Connectivity




Numerical Results

Connectivity










Numerical Results

Connectivity



References

-  B. Bollobás, **Graph Theory. An Introductory Course**, Springer-Verlag 1979. **99(25)**, 15879–15882 (2002).
-  F. Chung, **Spectral Graph Theory**, AMS 1997.
-  F. Chung, L. Lu, The average distances in random graphs with given expected degrees, Proc. Nat.Acad.Sci. 2002.
-  R. Diestel, **Graph Theory**, 3rd Edition, Springer-Verlag 2005.
-  P. Erdős, A. Rényi, On The Evolution of Random Graphs
-  G. Grimmett, **Probability on Graphs. Random Processes on Graphs and Lattices**, Cambridge Press 2010.
-  J. Leskovec, J. Kleinberg, C. Faloutsos, Graph Evolution: Densification and Shrinking Diameters, ACM Trans. on Knowl.Disc.Data, **1(1)** 2007.