

Lecture 10: Review of graph modeling and inference

Radu Balan

Department of Mathematics, AMSC and NWC
University of Maryland, College Park, MD

April 27, 2023

Main Problems

Main Problem

Input data: a weighted graph $G = (\mathcal{V}, W)$ with n nodes.

Issues:

- 1 Decide how well the two random graph models explain the data.
- 2 Partition the graph into two communities.
- 3 Construct an embedding $\{y_1, \dots, y_n\} \subset \mathbb{R}^d$ such that $W_{i,j} \sim \varphi(\|y_i - y_j\|)$ for some monotonically decreasing function φ .

Typical weight functions:

- 1 Exponential model: $\varphi(t) = Ce^{-t^2}$, for some $C > 0$.
- 2 Power law: $\varphi(t) = \frac{C}{t^p}$, for some $C > 0$ and $p > 0$.

Analysis

Three studies need to be done:

- 1 *Random graph hypothesis*: Sort edges by weight: from the largest weight to the smallest weight. Then compare sample statistics of 3-cliques, 4-cliques with their expectations under the two stochastic models, Erdős-Rényi and SSBM.

Analysis

Three studies need to be done:

- 1 *Random graph hypothesis*: Sort edges by weight: from the largest weight to the smallest weight. Then compare sample statistics of 3-cliques, 4-cliques with their expectations under the two stochastic models, Erdős-Rényi and SSBM.
- 2 Community Detection/Partition/Image Segmentation: Two classes of algorithms: spectral methods and SDP relaxations.

Analysis

Three studies need to be done:

- 1 *Random graph hypothesis*: Sort edges by weight: from the largest weight to the smallest weight. Then compare sample statistics of 3-cliques, 4-cliques with their expectations under the two stochastic models, Erdős-Rényi and SSBM.
- 2 Community Detection/Partition/Image Segmentation: Two classes of algorithms: spectral methods and SDP relaxations.
- 3 Embeddings: *Laplacian eigenmaps*: The geometric graph is obtained by solving the bottom $d + 1$ eigenproblems for the normalized symmetric Laplacian $\tilde{\Delta} = I - D^{-1}WD^{-1/2}$. Additional algorithms: LLE and ISOMAP.

Distribution of Cliques

Expected Values

Let X_q denote the number of q -cliques in a random graph G . Then the expectation of X_q in $\mathcal{G}_{n,p}$ class is

$$\mathbb{E}[X_q] = \binom{n}{q} p^{q(q-1)/2}$$

The expectation of X_q in the class $\Gamma^{n,m}$ is approximated by the above formula for $p = \frac{2m}{n(n-1)}$:

$$\mathbb{E}[X_q] \approx \binom{n}{q} \left(\frac{2m}{n(n-1)} \right)^{q(q-1)/2} \sim \theta_q \frac{m^{q(q-1)/2}}{n^{q(q-2)}}$$

$$\mathbb{E}[X_3] \sim \theta \frac{m^3}{n^3}, \quad \mathbb{E}[X_4] \sim \theta \frac{m^6}{n^8}$$

3-Cliques and 4-cliques

Thresholds

Theorem

Let $m = m(n)$ be the number of edges in $\Gamma^{n,m}$.

- ① If $m \gg n$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n} = \infty$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 3-clique}] \rightarrow 1.$
- ② If $m \ll n$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n} = 0$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 3-clique}] \rightarrow 0.$

Theorem

Let $m = m(n)$ be the number of edges in $\Gamma^{n,m}$.

- ① If $m \gg n^{4/3}$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n^{4/3}} = \infty$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 4-clique}] \rightarrow 1.$
- ② If $m \ll n^{4/3}$ (i.e. $\lim_{n \rightarrow \infty} \frac{m}{n^{4/3}} = 0$) then
 $\lim_{n \rightarrow \infty} \text{Prob}[G \in \Gamma^{n,m} \text{ has a 4-clique}] \rightarrow 0.$

3-Cliques and 4-Cliques

Behavior at the threshold

In general we obtain a "coarse threshold". Recall a Poisson process X with parameter λ has p.m.f. $Prob[X = k] = e^{-\lambda} \frac{\lambda^k}{k!}$.

Theorem

In $\mathcal{G}_{n,p}$,

- ① For $p = \frac{c}{n}$, X_3 is asymptotically Poisson with parameter $\lambda = c^3/6$.
- ② For $p = \frac{c}{n^{2/3}}$, X_4 is asymptotically Poisson with parameter $\lambda = c^6/24$.

3-Cliques and 4-Cliques

Behavior at the threshold

In general we obtain a "coarse threshold". Recall a Poisson process X with parameter λ has p.m.f. $Prob[X = k] = e^{-\lambda} \frac{\lambda^k}{k!}$.

Theorem

In $\mathcal{G}_{n,p}$,

- 1 For $p = \frac{c}{n}$, X_3 is asymptotically Poisson with parameter $\lambda = c^3/6$.
- 2 For $p = \frac{c}{n^{2/3}}$, X_4 is asymptotically Poisson with parameter $\lambda = c^6/24$.

Theorem

In $\Gamma^{n,m}$,

- 1 For $m = cn$, X_3 is asymptotically Poisson with parameter $\lambda = 4c^3/3$.
- 2 For $m = cn^{4/3}$, X_4 is asymptotically Poisson with parameter $\lambda = 8c^6/3$.

Eigenvalues of Laplacians

$\Delta, L, \tilde{\Delta}$

What do we know about the set of eigenvalues of these matrices for a graph G with n vertices?

- ① $\Delta = \Delta^T \geq 0$ and hence its eigenvalues are non-negative real numbers.
- ② $eigs(\tilde{\Delta}) = eigs(L) \subset [0, 2]$.
- ③ 0 is always an eigenvalue and its multiplicity equals the number of connected components of G ,

$$\dim \ker(\Delta) = \dim \ker(L) = \dim \ker(\tilde{\Delta}) = \# \text{connected components.}$$

Eigenvalues of Laplacians

$\Delta, L, \tilde{\Delta}$

What do we know about the set of eigenvalues of these matrices for a graph G with n vertices?

- 1 $\Delta = \Delta^T \geq 0$ and hence its eigenvalues are non-negative real numbers.
- 2 $eigs(\tilde{\Delta}) = eigs(L) \subset [0, 2]$.
- 3 0 is always an eigenvalue and its multiplicity equals the number of connected components of G ,

$$\dim \ker(\Delta) = \dim \ker(L) = \dim \ker(\tilde{\Delta}) = \# \text{connected components.}$$

Let $0 = \lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1}$ be the eigenvalues of $\tilde{\Delta}$. Denote

$$\lambda(G) = \max_{1 \leq i \leq n-1} |1 - \lambda_i|.$$

Note $\sum_{i=1}^{n-1} \lambda_i = \text{trace}(\tilde{\Delta}) = n$. Hence the average eigenvalue is about 1. $\lambda(G)$ is called *the absolute gap* and measures the spread of eigenvalues

The spectral absolute gap

 $\lambda(G)$

The main result in [9]) says that for connected graphs w/h.p.:

$$\lambda_1 \geq 1 - \frac{C}{\sqrt{\text{Average Degree}}} = 1 - \frac{C}{\sqrt{p(n-1)}} = 1 - C\sqrt{\frac{n}{2m}}.$$

Theorem (For class $\mathcal{G}_{n,p}$)

Fix $\delta > 0$ and let $p > (\frac{1}{2} + \delta)\log(n)/n$. Let $d = p(n-1)$ denote the expected degree of a vertex. Let \tilde{G} be the giant component of the Erdős-Rényi graph. For every fixed $\varepsilon > 0$, there is a constant $C = C(\delta, \varepsilon)$, so that

$$\lambda(\tilde{G}) \leq \frac{C}{\sqrt{d}}$$

with probability at least $1 - Cn \exp(-(2 - \varepsilon)d) - C \exp(-d^{1/4} \log(n))$.

Connectivity threshold: $p \sim \frac{\log(n)}{n}$.

The spectral absolute gap

 $\lambda(G)$

The main result in [9] says that for connected graphs w/h.p.:

$$\lambda_1 \geq 1 - \frac{C}{\sqrt{\text{Average Degree}}} = 1 - \frac{C}{\sqrt{p(n-1)}} = 1 - C\sqrt{\frac{n}{2m}}.$$

Theorem (For class $\Gamma^{n,m}$)

Fix $\delta > 0$ and let $m > \frac{1}{2}(\frac{1}{2} + \delta)n \log(n)$. Let $d = \frac{2m}{n}$ denote the expected degree of a vertex. Let \tilde{G} be the giant component of the Erdős-Rényi graph. For every fixed $\varepsilon > 0$, there is a constant $C = C(\delta, \varepsilon)$, so that

$$\lambda(\tilde{G}) \leq \frac{C}{\sqrt{d}}$$

with probability at least $1 - Cn \exp(-(2 - \varepsilon)d) - C \exp(-d^{1/4} \log(n))$.

Connectivity threshold: $m \sim \frac{1}{2}n \log(n)$.

Isometric Embeddings with Partial Data

Linear constraints

Given any set of vectors $\{y_1, \dots, y_n\}$ and their associated matrix $Y = [y_1 | \dots | y_n]$ their invariant to the action of the rigid transformations (translations, rotations, and reflections) is the Gram matrix of the centered system:

$$G = \left(I - \frac{1}{n} \mathbf{1} \cdot \mathbf{1}^T\right) Y^T Y \left(I - \frac{1}{n} \mathbf{1} \cdot \mathbf{1}^T\right) =: LY^T YL, \quad L = I - \frac{1}{n} \mathbf{1} \cdot \mathbf{1}^T.$$

On the other hand, the distance between points i and j can be computed by:

$$d_{i,j}^2 = \|y_i - y_j\|^2 = G_{i,i} - G_{i,j} + G_{j,j} - G_{j,i} = e_{ij}^T G e_{ij}$$

where

$$e_{ij} = \delta_i - \delta_j = [0 \cdots 0 \ 1 \cdots -1 \ 0 \cdots 0]^T$$

where 1 is on position i , -1 is on position j , and 0 everywhere else.

Almost Isometric Embeddings with Partial Data

The SDP Problem

Reference [10] proposes to find the matrix G by solving the following Semi-Definite Program:

$$\begin{aligned} \min \quad & \text{trace}(G) \\ G = G^T \geq 0 \\ G \cdot \mathbf{1} = 0 \\ |\langle Ge_{ij}, e_{ij} \rangle - \tilde{d}_{i,j}^2| \leq \varepsilon, \quad (i, j) \in \Theta \end{aligned}$$

where $\tilde{d}_{i,j}^2$ are noisy estimates $d_{i,j}$ and ε is the maximum noise level. The trace promotes low rank in this optimization. However, this is basically a feasibility problem: Decrease ε to the minimum value where a feasible solution exists. With probability 1 that is unique.

How to do this: Use CVX with Matlab.

Geometric Graph Embedding

Gram matrix factorization: The Algorithm

Algorithm

Input: Symmetric $n \times n$ Gram matrix G .

- 1 *Compute the eigendecomposition of G , $G = Q\Lambda Q^T$ with diagonal of Λ sorted in a descending order;*
- 2 *Determine the number d of significant positive eigenvalues;*
- 3 *Partition*

$$Q = [Q_1 \quad Q_2] \quad , \text{ and } \Lambda = \begin{bmatrix} \Lambda_1 & 0 \\ 0 & \Lambda_2 \end{bmatrix}$$

where Q_1 contains the first d columns of Q , and Λ_1 is the $d \times d$ diagonal matrix of significant positive eigenvalues of G .

- 4 *Compute:*

$$Y = \Lambda_1^{1/2} Q_1^T$$

Output: Dimension d and $d \times n$ matrix Y of vectors $Y = [y_1 | \cdots | y_n]$

Nearly Isometric Embeddings with Partial Data

Stability to Noise

[10] proves the following stability result in the case of partial measurements. Here we denote $\Theta_r = \{(i, j), \|y_i - y_j\| \leq r\}$ the set of all pairs of points at distance at most r .

Theorem

Let $\{y_1, \dots, y_n\}$ be n nodes distributed uniformly at random in the hypercube $[-0.5, 0.5]^d$. Further, assume that we are given noisy measurement of all distances in Θ_r for some $r \geq 10\sqrt{d}(\log(n)/n)^{1/d}$ and the induced geometric graph of edges is connected. Let $\tilde{d}_{i,j}^2 = d_{i,j}^2 + \nu_{i,j}$ with $|\nu_{i,j}| \leq \varepsilon$. Then with high probability, the error distance between the estimated $\hat{Y} = [\hat{y}_1 | \dots | \hat{y}_n]$ returned by the SDP-based algorithm and the correct coordinate matrix $Y = [y_1 | \dots | y_n]$ is upper bounded as

$$\|L\hat{Y}^T\hat{Y}L - LY^TYL\|_1 \leq C_1(nr^d)^5 \frac{\varepsilon}{r^4}.$$

Optimization Criterion

Assume $\mathcal{G} = (\mathcal{V}, W)$ is a undirected weighted graph with n nodes and weight matrix W .

We interpret $W_{i,j}$ as the *similarity* between nodes i and j . The larger the weight the more similar the nodes, and the closer they are in a geometric graph embedding.

Thus we look for a dimension $d > 0$ and a set of points

$\{y_1, y_2, \dots, y_n\} \subset \mathbb{R}^d$ so that $d_{i,j} = \|y_i - y_j\|$'s is small for large weight $W_{i,j}$. This means we want to minimize

$$J(y_1, y_2, \dots, y_n) = \sum_{1 \leq i, j \leq n} W_{i,j} \|y_i - y_j\|^2,$$

To avoid trivial solution $Y = 0$ we impose a normalization condition:

$$YDY^T = I_d.$$

The Optimization Problem

Combining the criterion with the constraint:

$$(LE) : \begin{array}{ll} \text{minimize} & \text{trace} \left\{ Y \Delta Y^T \right\} \\ \text{subject to} & Y D Y^T = I_d \end{array}$$

we obtained the *Laplacian Eigenmap* problem.

Good news: The optimizer Y is obtained by solving an eigenproblem.

Laplacian Eigenmaps Embedding

Algorithm

Algorithm (Laplacian Eigenmaps)

Input: Weight matrix W , target dimension d

- 1 Construct the diagonal matrix $D = \text{diag}(D_{ii})_{1 \leq i \leq n}$, where $D_{ii} = \sum_{k=1}^n W_{i,k}$.
- 2 Construct the normalized Laplacian $\tilde{\Delta} = I - D^{-1/2} W D^{-1/2}$.
- 3 Compute the bottom $d + 1$ eigenvectors e_1, \dots, e_{d+1} , $\tilde{\Delta} e_k = \lambda_k e_k$, $0 = \lambda_1 \cdots \lambda_{d+1}$.

Laplacian Eigenmaps Embedding

Algorithm-cont's

Algorithm (Laplacian Eigenmaps - cont'd)

- ④ Construct the $d \times n$ matrix Y ,

$$Y = \begin{bmatrix} e_2 \\ \vdots \\ e_{d+1} \end{bmatrix} D^{-1/2}$$

- ⑤ The new geometric graph is obtained by converting the columns of Y into n d -dimensional vectors:

$$\begin{bmatrix} y_1 & | & \cdots & | & y_n \end{bmatrix} = Y$$

Output: Set of points $\{y_1, y_2, \dots, y_n\} \subset \mathbb{R}^d$.

Problem Formulation

Given: It is assumed that we are given a set of points $\{x_1, \dots, x_n\} \subset \mathbb{R}^N$, or a weight matrix W , where $W_{i,j}$ is inverse monotonically dependent to distances $\|x_i - x_j\|$; the smaller the distance $\|x_i - x_j\|$ the larger the weight $W_{i,j}$.

Target: We look for a dimension $d > 0$ and a set of points $\{y_1, y_2, \dots, y_n\} \subset \mathbb{R}^d$ so that all $d_{i,j} = \|y_i - y_j\|$'s are compatible with the raw data.

Approaches:

- 1 Principal Component Analysis
- 2 Independent Component Analysis
- 3 Laplacian Eigenmaps
- 4 Local Linear Embeddings (LLE)
- 5 Isomaps

Principal Component Analysis

Algorithm

Algorithm (Principal Component Analysis)

Input: Data vectors $\{x_1, \dots, x_n\} \in \mathbb{R}^N$; dimension d .

- ① *If affine subspace is the goal, append '1' at the end of each data vector.*
- ① *Compute the sample covariance matrix*

$$R = \sum_{k=1}^n x_k x_k^T$$

- ② *Solve the eigenproblems $Re_k = \sigma_k^2 e_k$, $1 \leq k \leq N$, order eigenvalues $\sigma_1^2 \geq \sigma_2^2 \geq \dots \geq \sigma_N^2$ and normalize the eigenvectors $\|e_k\|_2 = 1$.*

Principal Component Analysis

Algorithm - cont'ed

Algorithm (Principal Component Analysis)

- ③ *Construct the co-isometry*

$$U = \begin{bmatrix} e_1^T \\ \vdots \\ e_d^T \end{bmatrix}.$$

- ④ *Project the input data*

$$y_1 = Ux_1, \quad y_2 = Ux_2, \quad \dots, \quad y_n = Ux_n.$$

Output: Lower dimensional data vectors $\{y_1, \dots, y_n\} \in \mathbb{R}^d$.

The orthogonal projection is given by $P = \sum_{k=1}^d e_k e_k^T$ and the optimal subspace is $V = \text{Ran}(P)$

Dimension Reduction using Laplacian Eigenmaps

Algorithm

Algorithm (Dimension Reduction using Laplacian Eigenmaps)

Input: A geometric graph $\{x_1, x_2, \dots, x_n\} \subset \mathbb{R}^N$. Parameters: threshold τ , weight coefficient α , and dimension d .

- 1 *Compute the set of pairwise distances $\|x_i - x_j\|$ and convert them into a set of weights:*

$$W_{i,j} = \begin{cases} \exp(-\alpha\|x_i - x_j\|^2) & \text{if } \|x_i - x_j\| \leq \tau \\ 0 & \text{if otherwise} \end{cases}$$

- 2 *Compute the $d + 1$ bottom eigenvectors of the normalized Laplacian matrix $\tilde{\Delta} = I - D^{-1/2}WD^{-1/2}$, $\tilde{\Delta}e_k = \lambda_k e_k$, $1 \leq k \leq d + 1$, $0 = \lambda_0 \leq \dots \leq \lambda_{d+1}$, where $D = \text{diag}(\sum_{k=1}^n W_{i,k})_{1 \leq i \leq n}$.*

Dimension Reduction using Laplacian Eigenmaps

Algorithm - cont'd

Algorithm (Dimension Reduction using Laplacian Eigenmaps-cont'd)

- ③ Construct the $d \times n$ matrix Y ,

$$Y = \begin{bmatrix} e_2^T \\ \vdots \\ e_{d+1}^T \end{bmatrix} D^{-1/2}$$

- ④ The new geometric graph is obtained by converting the columns of Y into n d -dimensional vectors:

$$\begin{bmatrix} y_1 & | & \cdots & | & y_n \end{bmatrix} = Y$$

Output: $\{y_1, \dots, y_n\} \subset \mathbb{R}^d$.

Dimension Reduction using Isomaps

Algorithm

Algorithm (Dimension Reduction using Isomap)

Input: A geometric graph $\{x_1, x_2, \dots, x_n\} \subset \mathbb{R}^N$. Parameters: neighborhood size K and dimension d .

- 1 Construct the symmetric matrix S of squared pairwise distances:
 - 1 Construct the sparse matrix T , where for each node i find the nearest K neighbors \mathcal{V}_i and set $T_{i,j} = \|x_i - x_j\|_2$, $j \in \mathcal{V}_i$.
 - 2 For any pair of two nodes (i, j) compute $d_{i,j}$ as the length of the shortest path, $\sum_{p=1}^L T_{k_{p-1}, k_p}$ with $k_0 = i$ and $k_L = j$, using e.g. Dijkstra's algorithm.
 - 3 Set $S_{i,j} = d_{i,j}^2$.

Dimension Reduction using Isomaps

Algorithm - cont'd

Algorithm (Dimension Reduction using Isomap - cont'd)

- 2 Compute the Gram matrix G :

$$\rho = \frac{1}{2n} \mathbf{1}^T \cdot S \cdot \mathbf{1}, \quad \nu = \frac{1}{n} (S \cdot \mathbf{1} - \rho \mathbf{1})$$

$$G = \frac{1}{2} \nu \cdot \mathbf{1}^T + \frac{1}{2} \mathbf{1} \cdot \nu^T - \frac{1}{2} S$$

- 3 Find the top d eigenvectors of G , say e_1, \dots, e_d so that $GE = E\Lambda$, form the matrix Y and then collect the columns:

$$Y = \Lambda^{1/2} \begin{bmatrix} e_1^T \\ \vdots \\ e_d^T \end{bmatrix} = \begin{bmatrix} y_1 & | & \cdots & | & y_n \end{bmatrix}$$

Output: $\{v_1, \dots, v_n\} \subset \mathbb{R}^d$.

References



B. Bollobás, **Graph Theory. An Introductory Course**, Springer-Verlag 1979. **99**(25), 15879–15882 (2002).



S. Boyd, L. Vandenberghe, **Convex Optimization**, available online at: <http://stanford.edu/boyd/cvxbook/>



F. Chung, **Spectral Graph Theory**, AMS 1997.



F. Chung, L. Lu, The average distances in random graphs with given expected degrees, Proc. Nat.Acad.Sci. 2002.



F. Chung, L. Lu, V. Vu, The spectra of random graphs with Given Expected Degrees, Internet Math. **1**(3), 257–275 (2004).



R. Diestel, **Graph Theory**, 3rd Edition, Springer-Verlag 2005.



P. Erdős, A. Rényi, On The Evolution of Random Graphs



G. Grimmett, **Probability on Graphs. Random Processes on Graphs and Lattices**, Cambridge Press 2010.



C. Hoffman, M. Kahle, E. Paquette, Spectral Gap of Random Graphs and Applications to Random Topology, arXiv: 1201.0425 [math.CO] 17 Sept. 2014.



A. Javanmard, A. Montanari, Localization from Incomplete Noisy Distance Measurements, arXiv:1103.1417, Nov. 2012; also ISIT 2011.



J. Leskovec, J. Kleinberg, C. Faloutsos, Graph Evolution: Densification and Shrinking Diameters, ACM Trans. on Knowl.Disc.Data, **1(1)** 2007.



S.T. Roweis, L.K. Saul, Locally linear embedding, Science 290, 2323–2326 (2000).



J.B. Tenenbaum, V. de Silva, J.C. Langford, A global geometric framework for nonlinear dimensionality reduction, Science 290, 2319–2323 (2000).