

AFFINE DELIGNE-LUSZTIG VARIETIES IN AFFINE FLAG VARIETIES

ULRICH GÖRTZ, THOMAS J. HAINES, ROBERT E. KOTTWITZ, AND DANIEL C.
REUMAN

ABSTRACT. This paper studies affine Deligne-Lusztig varieties in the affine flag manifold of a split group. Among other things, it proves emptiness for certain of these varieties, relates some of them to those for Levi subgroups, extends previous conjectures concerning their dimensions, and generalizes the superset method.

1. INTRODUCTION

This paper, a continuation of [GHKR], investigates affine Deligne-Lusztig varieties $X_x(b)$ in the affine flag manifold $X = G(L)/I$ of a split group G . To provide some context we begin by discussing affine Deligne-Lusztig varieties $X_\mu(b)$ in the affine Grassmannian $G(L)/G(\mathfrak{o})$.

It is known that $X_\mu(b)$ is non-empty if and only if Mazur's inequality is satisfied, that is to say, if and only if the σ -conjugacy class $[b]$ of b is less than or equal to $[\epsilon^\mu]$ in the natural partial order on the set $B(G)$ of σ -conjugacy classes in $G(L)$. This was proved in two steps: the problem was reduced [KR] to one on root systems, which was then solved for classical split groups by C. Lucarelli [Lu] and now for all split groups by Q. Gashi [Ga].

A conjectural formula for $\dim X_\mu(b)$ was put forward by Rapoport [Ra], who pointed out its similarity to a conjecture of Chai's [Ch] on dimensions of Newton strata in Shimura varieties. In [GHKR] Rapoport's dimension conjecture was reduced to the superbasic case, which was then solved by Viehmann [V1].

Now we return to affine Deligne-Lusztig varieties $X_x(b)$ in the affine flag manifold. Here x denotes an element in the extended affine Weyl group \widetilde{W} of G . For some years now a challenging problem has been to "explain" the emptiness pattern one sees in the figures in [Re2] and [GHKR]. In other words, for a given b , one wants to understand the set of $x \in \widetilde{W}$ for which $X_x(b)$ is empty. Let us begin by discussing the simplest case, that in which $b = 1$ and x is *shrunk*, by which we mean that it lies in the union of the shrunken Weyl chambers (see [GHKR]). Then Reuman [Re2] observed that a simple rule explained the emptiness pattern in types A_1 , A_2 , and C_2 and conjectured that the same might be true in general. Computer calculations [GHKR] provided further evidence for the truth of Reuman's conjecture. However emptiness in the non-shrunken case remained quite mysterious.

1991 *Mathematics Subject Classification*. Primary 14L05; Secondary 11S25, 20G25, 14F30.
Haines was partially supported by NSF Grant FRG-0554254 and a Sloan Research Fellowship.
Kottwitz was partially supported by NSF Grant DMS-0245639.
Reuman was partially supported by United States NSF grant DMS-0443803.

In this paper, among other things, we give a precise conjecture (namely Conjecture 9.3.2) describing the whole emptiness pattern for any basic b . This is more general in two ways: we no longer require that $b = 1$ (though we do require that b be basic), and we no longer restrict attention to shrunken x . Computer calculations support this conjecture, and for shrunken x we show (see Proposition 9.4.5) that the new conjecture reduces to Reuman's. We prove (see Corollary 9.3.1) one direction of this new conjecture, namely that emptiness does occur when predicted; it remains a challenging problem to prove that non-emptiness occurs when predicted.

In fact Proposition 9.2.1 proves the emptiness of certain $X_x(b)$ even when b is not basic. However, in the non-basic case, there is a second cause for emptiness, stemming from Mazur's inequality. One might hope that these are the only two causes for emptiness. This is slightly too naive. Mazur's inequality works perfectly for $G(\sigma)$ -double cosets, but not for Iwahori double cosets, and would have to be improved slightly (in the Iwahori case) before it could be applied to give an optimal emptiness criterion. Although we do not yet know how to formulate Mazur's inequalities in the Iwahori case, in section 12 we are able to describe the information they should carry, whatever they end up being.

We now turn to the dimensions of non-empty affine Deligne-Lusztig varieties in the affine flag manifold. In [GHKR] we formulated two conjectures of this kind, and here we will extend both of them (in a way that is supported by computer evidence).

Conjecture 9.4.1(a) extends Conjecture 7.2.2 of [GHKR] from $b = 1$ to all basic b . This conjecture predicts the dimension of $X_x(b)$ for all shrunken x for which $X_x(b)$ is expected to be non-empty.

Conjecture 9.4.1(b) extends Conjecture 7.5.1 of [GHKR] from translation elements $b = \epsilon^\nu$ to all b . For this we need the following notation: b_b will denote a representative of the unique basic σ -conjugacy class whose image in Λ_G is the same as that of b . (Equivalently, $[b_b]$ is at the bottom of the connected component of $[b]$ in the poset $B(G)$.) In this second conjecture, it is the difference of the dimensions of $X_x(b)$ and $X_x(b_b)$ that is predicted. It is not required that x be shrunken, but $X_x(b)$ and $X_x(b_b)$ are required to be non-empty, and the length of x is required to be big enough. In the conjecture we phrase this last condition rather crudely as $\ell(x) \geq N_b$ for some (unspecified) constant N_b that depends on b . However the evidence of computer calculations suggests that for fixed b , having x such that $X_x(b)$ and $X_x(b_b)$ are both non-empty is almost (but not quite!) enough to make our prediction valid for x . It would be very interesting to understand this phenomenon better, though some insight into it is already provided by Beazley's work on Newton strata for $SL(3)$ [Be]. In addition, when $\ell(x) \geq N_b$, we conjecture that the non-emptiness of $X_x(b)$ is equivalent to that of $X_x(b_b)$.

The main result of this paper is Theorem 2.1.2, from which all our results on emptiness follow. This theorem states that for any semistandard parabolic subgroup $P = MN$ and any P -alcove (see Definition 2.1.1) $x\mathbf{a}$, every element of IxI is σ -conjugate under I to an element of $I_M x I_M$.

We also use Theorem 2.1.2 to prove an "Iwahori version" of the Hodge-Newton decomposition (see Theorem 2.1.4), in the form of a bijection

$$J_b^M \backslash X_x^M(b) \xrightarrow{\sim} J_b^G \backslash X_x^G(b),$$

valid when $x\mathbf{a}$ is a P -alcove (so that in particular $x \in \widetilde{W}_M$) and $b \in M(L)$. It is striking that the notion of P -alcove, discovered in the attempt to understand the entire emptiness pattern for the $X_x(b)$ when b is basic, is also precisely the notion needed for our Hodge-Newton decomposition.

For computer calculations, and perhaps also for theoretical reasons, it is useful to extend Reuman's superset method [Re2] from $b = 1$ to general b . To that end we introduce (see Definition 13.1.1) the notion of *fundamental alcove* $y\mathbf{a}$. We show that for each σ -conjugacy class $[b]$ there exists a fundamental alcove $y\mathbf{a}$ such that the whole double coset IyI is contained in $[b]$. We then explain why this allows one to use a superset method to analyze the emptiness of $X_x(b)$ for any x .

In addition we introduce a generalization of the superset method. The superset method is based on I -orbits in the affine flag manifold X . Now [GHKR] used orbits of $U(L)$, where U is the unipotent radical of a Borel subgroup containing our standard split maximal torus A . The generalized superset method interpolates between these two extremes, being based on orbits of $I_M N(L)$ on X , where $P = MN$ is a standard parabolic subgroup of G . Theorem 11.2.1 and the discussion preceding it explain how the generalized superset method can be used to study dimensions of affine Deligne-Lusztig varieties.

Finally, for any standard parabolic subgroup $P = MN$ and any basic $b \in M(L)$ Proposition 12.0.6 gives a formula for the dimension of $X_x(b)$ in terms of dimensions of affine Deligne-Lusztig varieties for M as well as intersections of I -orbits and $N'(L)$ -orbits for certain Weyl group conjugates N' of N . This generalizes Theorem 6.3.1 of [GHKR] and is also analogous to Proposition 5.6.1 of [GHKR], but with the affine Grassmannian replaced by the affine flag manifold.

The first and second named authors thank the American Institute of Mathematics for the invitation to the workshop on Buildings and Combinatorial Representation Theory which provided an opportunity to work on questions related to this paper. The second and third named authors thank the University of Bonn and the Max-Planck-Institut für Mathematik Bonn for providing other, equally valuable opportunities of this kind.

We are very grateful to Eva Viehmann for her helpful remarks on the manuscript.

Notation. We follow the notation of [GHKR], for the most part. Let k be a finite field with q elements, and let \bar{k} be an algebraic closure of k . We consider the field $L := \bar{k}((\epsilon))$ and its subfield $F := k((\epsilon))$. We write $\sigma : x \mapsto x^q$ for the Frobenius automorphism of \bar{k}/k , and we also regard σ as an automorphism of L/F in the usual way, so that $\sigma(\sum a_n \epsilon^n) = \sum \sigma(a_n) \epsilon^n$. We write \mathfrak{o} for the valuation ring $\bar{k}[[\epsilon]]$ of L .

Let G be a split connected reductive group over k , and let A be a split maximal torus of G . Write R for the set of roots of A in G . Put $\mathfrak{a} := X_*(A)_{\mathbb{R}}$. Write W for the Weyl group of A in G . Fix a Borel subgroup $B = AU$ containing A with unipotent radical U , and write R^+ for the corresponding set of positive roots, that is, those occurring in U . We denote by ρ the half-sum of the positive roots. For $\lambda \in X_*(A)$ we write ϵ^λ for the element of $A(F)$ obtained as the image of $\epsilon \in \mathbb{G}_m(F)$ under the homomorphism $\lambda : \mathbb{G}_m \rightarrow A$.

Let C_0 denote the dominant Weyl chamber, which by definition is the set of $x \in \mathfrak{a}$ such that $\langle \alpha, x \rangle > 0$ for all $\alpha \in R^+$. We denote by \mathbf{a} the unique alcove in the dominant Weyl chamber whose closure contains the origin, and call it the base alcove. As Iwahori subgroup I we choose the one fixing the base alcove \mathbf{a} ; I

is then the inverse image of the opposite Borel group of B under the projection $K := G(\mathfrak{o}) \rightarrow G(\bar{k})$. The opposite Borel arises here due to our convention that ϵ^λ acts on the standard apartment \mathfrak{a} by translation by λ (rather than by translation by the negative of λ), so that the stabilizer in $G(L)$ of $\lambda \in X_*(A) \subset \mathfrak{a}$ is $\epsilon^\lambda K \epsilon^{-\lambda}$. With this convention the Lie algebra of the Iwahori subgroup stabilizing an alcove \mathfrak{b} in the standard apartment is made up of affine root spaces $\epsilon^j \mathfrak{g}_\alpha$ for all pairs (α, j) such that $\alpha - j \leq 0$ on \mathfrak{b} (with \mathfrak{g}_α denoting the root subspace corresponding to α).

We will often think of alcoves in a slightly different way. Let Λ_G denote the quotient of $X_*(A)$ by the coroot lattice. The apartment \mathcal{A} corresponding to our fixed maximal torus A can be decomposed as a product $\mathcal{A} = \mathcal{A}_{\text{der}} \times V_G$, where $V_G := \Lambda_G \otimes \mathbb{R}$ and where \mathcal{A}_{der} is the apartment corresponding to $A_{\text{der}} := G_{\text{der}} \cap A$ in the building for G_{der} . By an *extended alcove* we mean a subset of the apartment \mathcal{A} of the form $\mathfrak{b} \times c$, where \mathfrak{b} is an alcove in \mathcal{A}_{der} and $c \in \Lambda_G$. Clearly each extended alcove determines a unique alcove in the usual sense, but not conversely. However, in the sequel we will often use the terms interchangeably, leaving context to determine what is meant. In particular, we often write \mathfrak{a} in place of $\mathfrak{a} \times 0$.

We denote by \widetilde{W} the extended affine Weyl group $X_*(A) \rtimes W$ of G . Then \widetilde{W} acts transitively on the set of all alcoves in \mathfrak{a} , and simply transitively on the set of all extended alcoves. Let $\Omega = \Omega_{\mathfrak{a}}$ denote the stabilizer of \mathfrak{a} when it is viewed as an alcove in the usual (non-extended) sense. We can write an extended (resp. non-extended) alcove in the form $x\mathfrak{a}$ for a unique element $x \in \widetilde{W}$ (resp. $x \in \widetilde{W}/\Omega$). Of course, this is just another way of saying that we can think of extended alcoves simply as elements of \widetilde{W} .

As usual a standard parabolic subgroup is one containing B , and a semistandard parabolic subgroup is one containing A . Similarly, a semistandard Levi subgroup is one containing A , and a standard Levi subgroup is the unique semistandard Levi component of a standard parabolic subgroup. Given a semistandard Levi subgroup M of G we write $\mathcal{P}(M)$ for the set of parabolic subgroups of G admitting M as Levi component. For $P \in \mathcal{P}(M)$ we denote by $\overline{P} = M\overline{N} \in \mathcal{P}(M)$ the parabolic subgroup opposite to P . We write R_N for the set of roots of A in N . We denote by $I_M, I_N, I_{\overline{N}}$ the intersections of I with M, N, \overline{N} respectively; one then has the Iwahori decomposition $I = I_N I_M I_{\overline{N}}$.

Recall that for $x \in \widetilde{W}$ and $b \in G(L)$ the affine Deligne-Lusztig variety $X_x(b)$ is defined by

$$X_x(b) := \{g \in G(L)/I : g^{-1}b\sigma(g) \in IxI\}.$$

In the sequel we often abuse notation and use the symbols G, P, M, N to denote the corresponding objects over L .

Let $b \in G(L)$. We denote by $[b]$ the σ -conjugacy class of b inside $G(L)$:

$$[b] = \{g^{-1}b\sigma(g); g \in G(L)\},$$

and for a subgroup $H \subseteq G(L)$ we write

$$[b]_H := \{h^{-1}b\sigma(h); h \in H\} \subseteq G(L)$$

for the σ -conjugacy class of b under H . Further notation relevant to $B(G)$ such as η_G will be explained in section 7.

Finally we note that ${}^x I$ will be used as an abbreviation for xIx^{-1} . We use the symbols \subset and \subseteq interchangeably with the meaning “not necessarily strict inclusion”.

2. STATEMENT OF THE MAIN THEOREM

2.1. Let $\alpha \in R$. We identify the root group U_α with the additive group \mathbb{G}_a over k , which then allows us to identify $U_\alpha(L) \cap K$ with \mathfrak{o} . The root α induces a partial order \geq_α on the set of (extended) alcoves in the standard apartment as follows: given an alcove \mathbf{b} , write it as $x\mathbf{a}$ for $x \in \widetilde{W}$. Let $k(\alpha, \mathbf{b}) \in \mathbb{Z}$ such that $U_\alpha(L) \cap {}^x I = \epsilon^{k(\alpha, \mathbf{b})} \mathfrak{o}$. In other words, $k(\alpha, \mathbf{b})$ is the unique integer k such that \mathbf{b} lies in the region between the affine root hyperplanes $H_{\alpha, k} = \{x \in X_*(A)_\mathbb{R}; \langle \alpha, x \rangle = k\}$ and $H_{\alpha, k-1}$. This description shows immediately that $k(\alpha, \mathbf{b}) + k(-\alpha, \mathbf{b}) = 1$. (For instance, we have $k(\alpha, \mathbf{a}) = 1$ if $\alpha > 0$ and $k(\alpha, \mathbf{a}) = 0$ if $\alpha < 0$. This reflects the fact that the fixer I of \mathbf{a} is the inverse image of the opposite Borel \overline{B} under the projection $G(\mathfrak{o}) \rightarrow G(\overline{k})$.) We define

$$\mathbf{b}_1 \geq_\alpha \mathbf{b}_2 : \iff k(\alpha, \mathbf{b}_1) \geq k(\alpha, \mathbf{b}_2).$$

This is a partial order in the weak sense: $\mathbf{b}_1 \geq_\alpha \mathbf{b}_2$ and $\mathbf{b}_2 \geq_\alpha \mathbf{b}_1$ does not imply that $\mathbf{b}_1 = \mathbf{b}_2$. We also define

$$\mathbf{b}_1 >_\alpha \mathbf{b}_2 : \iff k(\alpha, \mathbf{b}_1) > k(\alpha, \mathbf{b}_2).$$

Definition 2.1.1. Let $P = MN$ be a semistandard parabolic subgroup. Let $x \in \widetilde{W}$. We say $x\mathbf{a}$ is a P -alcove, if

- (1) $x \in \widetilde{W}_M$, and
- (2) $\forall \alpha \in R_N, x\mathbf{a} \geq_\alpha \mathbf{a}$.

We say $x\mathbf{a}$ is a strict P -alcove if instead of (2) we have

- (2') $\forall \alpha \in R_N, x\mathbf{a} >_\alpha \mathbf{a}$.

Note that condition (2) depends only on the image of x in \widetilde{W}/Ω ; however, condition (1) depends on x itself.

By the definition of the partial order \geq_α , the condition (2) is equivalent to

$$(2.1.1) \quad \forall \alpha \in R_N, \quad U_\alpha \cap {}^x I \subseteq U_\alpha \cap I,$$

or, likewise, to

$$(2.1.2) \quad \forall \alpha \in R_N, \quad U_{-\alpha} \cap {}^x I \supseteq U_{-\alpha} \cap I$$

and under our assumption that $x \in \widetilde{W}_M$, these in turn are equivalent to the conditions

$$(2.1.3) \quad {}^x(N \cap I) \subseteq N \cap I \iff {}^x(\overline{N} \cap I) \supseteq \overline{N} \cap I.$$

(And condition (2') is equivalent to (2.1.1) with the inclusions replaced by strict inclusions.) Indeed, noting that conjugation by $x = \epsilon^\lambda w$ permutes the subgroups U_α with $\alpha \in R_N$, it is easy to see from the (Iwahori) factorization

$$(2.1.4) \quad N \cap I = \prod_{\alpha \in R_N} U_\alpha \cap I,$$

that (2.1.1) is equivalent to (2.1.3). For a fixed semistandard parabolic subgroup $P = MN$, the set of alcoves $x\mathbf{a}$ which satisfy (2.1.1) forms a union of ‘‘acute cones of alcoves’’ in the sense of [HN]. We shall explain this in section 3 below.

Our main theorem concerns the map

$$\begin{aligned} \phi : I \times I_M x I_M &\rightarrow I x I \\ (i, m) &\mapsto i m \sigma(i)^{-1}. \end{aligned}$$

There is a left action of I_M on $I \times I_M x I_M$ given by $i_M(i, m) = (i i_M^{-1}, i_M m \sigma(i_M)^{-1})$, for $i_M \in I_M$, $i \in I$ and $m \in I_M x I_M$. Let us denote by $I \times^{I_M} I_M x I_M$ the quotient of $I \times I_M x I_M$ by this action of I_M . Denote by $[i, m]$ the equivalence class of $(i, m) \in I \times I_M x I_M$. The map ϕ obviously factors through $I \times^{I_M} I_M x I_M$. We can now state the main theorem.

Theorem 2.1.2. *Suppose $P = MN$ is a semistandard parabolic subgroup, and $\mathbf{x}a$ is a P -alcove. Then the map*

$$\phi : I \times^{I_M} I_M x I_M \rightarrow IxI$$

induced by $(i, m) \mapsto i m \sigma(i)^{-1}$, is surjective. If $\mathbf{x}a$ is a strict P -alcove, then ϕ is injective. In general, ϕ is not injective, but if $[i, m]$ and $[i', m']$ belong to the same fiber of ϕ , the elements m and m' are σ -conjugate by an element of I_M .

This theorem was partially inspired by Labesse's study of the "elementary functions" he introduced in [La].

Let us mention a few consequences. First, consider the quotient $IxI/\sigma I$, where the action of I on IxI is given by σ -conjugation. We also can form in a parallel manner the quotient $I_M x I_M / \sigma I_M$. Further, let $B(G)_x$ denote the set of σ -conjugacy classes $[b]$ in $G(L)$ which meet IxI . We note that for $G = SL_3$ all of the sets $B(G)_x$ have been determined explicitly by Beazley [Be].

Corollary 2.1.3. *Suppose $P = MN$ is semistandard, and $\mathbf{x}a$ is a P -alcove. Then the following statements hold.*

(a) *The inclusion $I_M x I_M \hookrightarrow IxI$ induces a bijection*

$$I_M x I_M / \sigma I_M \xrightarrow{\sim} IxI / \sigma I.$$

(b) *The canonical map $\iota : B(M)_x \rightarrow B(G)_x$ is bijective.*

Part (a) follows directly from Theorem 2.1.2. Indeed, the surjectivity of ϕ implies the surjectivity of $I_M x I_M / \sigma I_M \rightarrow IxI / \sigma I$. As for the injectivity of the latter, note that if $i \in I$ and $m, m' \in I_M x I_M$ satisfy $i m \sigma(i)^{-1} = m'$, then $[i, m]$ and $[1, m']$ belong to the same fiber of ϕ . As for part (b), we will derive it from part (a) in section 8. (In fact the surjectivity in part (b) follows easily from the surjectivity in Theorem 2.1.2.)

Another consequence is a version of the Hodge-Newton decomposition, given in Theorem 2.1.4 below. For affine Deligne-Lusztig varieties in the affine Grassmannian of a split group, the analogous Hodge-Newton decomposition was proved under unnecessarily strict hypotheses in [K3] and in the general case by Viehmann [V2, Theorem 1] (see also Mantovan-Viehmann [MV] for the case of unramified groups). To state this we need to fix a standard parabolic subgroup $P = MN$ and an element $b \in M(L)$. Let $K_M = M \cap K$, where K , as usual, denotes $G(\mathfrak{o})$. For a G -dominant coweight $\mu \in X_*(A)$, the σ -centralizer $J_b^G := \{g \in G(L) : g^{-1} b \sigma(g) = b\}$ of b acts naturally on the affine Deligne-Lusztig variety $X_\mu^G(b) \subset G(L)/K$ defined to be

$$X_\mu^G(b) := \{gK \in G(L)/K \mid g^{-1} b \sigma(g) \in K e^\mu K\}.$$

Also, J_b^M acts on $X_\mu^M(b) \subset M(L)/K_M$. Now the Hodge-Newton decomposition under discussion asserts the following: suppose that the Newton point $\bar{\nu}_b^M \in X_*(A)_\mathbb{R}$ is G -dominant, and that $\eta_M(b) = \mu$ in Λ_M . Then the canonical closed immersion $X_\mu^M(b) \hookrightarrow X_\mu^G(b)$ induces a bijection

$$J_b^M \backslash X_\mu^M(b) \xrightarrow{\sim} J_b^G \backslash X_\mu^G(b).$$

Of course if we impose the stricter condition that $\langle \alpha, \overline{\nu}_b^M \rangle > 0$ for all $\alpha \in R_N$, then $J_b^M = J_b^G$ and so we get the stronger conclusion $X_\mu^M(b) \cong X_\mu^G(b)$, yielding what is normally known as the Hodge-Newton decomposition in this context. The version with the weaker condition is essentially a result of Viehmann, who formulates it somewhat differently [V2, Theorem 2], in a way that brings out a dichotomy occurring when G is simple.

In the affine flag variety, it still makes sense to ask how $X_x^G(b)$ and $X_x^M(b)$ are related, for $x \in \widetilde{W}_M$ and $b \in M(L)$. Our Hodge-Newton decomposition below provides some information in this direction.

Theorem 2.1.4. *Suppose $P = MN$ is semistandard and $x\mathbf{a}$ is a P -alcove.*

- (a) *If $X_x^G(b) \neq \emptyset$, then $[b]$ meets $M(L)$.*
- (b) *Suppose $b \in M(L)$. Then the canonical closed immersion $X_x^M(b) \hookrightarrow X_x^G(b)$ induces a bijection*

$$J_b^M \setminus X_x^M(b) \xrightarrow{\cong} J_b^G \setminus X_x^G(b).$$

Note that part (b) implies that if $x\mathbf{a}$ is a P -alcove, then for every $b \in M(L)$, we have $X_x^G(b) = \emptyset$ if and only if $X_x^M(b) = \emptyset$. We will prove Theorem 2.1.4 in section 8 and then derive some further consequences relating to emptiness/non-emptiness of $X_x^G(b)$, in section 9.

3. P -ALCOVES AND ACUTE CONES OF ALCOVES

Let $P = MN$ be a fixed semistandard parabolic subgroup. The aim of this section is to help the reader visualize the set of P -alcoves. Let \mathfrak{A} denote the set of alcoves $x\mathbf{a}$ which satisfy the inequalities $x\mathbf{a} \geq_\alpha \mathbf{a}$ for all $\alpha \in R_N$.

For each element $w \in W$, we recall the notion of *acute cone* of alcoves $C(\mathbf{a}, w)$, following [HN]. Given an affine hyperplane $H = H_{\alpha, k} = H_{-\alpha, -k}$, we assume α has the sign such that $\alpha \in w(R^+)$, i. e. such that α is a positive root with respect to wB . Then define the *w-positive* half space

$$H^{w+} = \{v \in X_*(A)_\mathbb{R} : \langle \alpha, v \rangle > k\}.$$

Let H^{w-} denote the other half-space.

Then the acute cone of alcoves $C(\mathbf{a}, w)$ is defined to be the set of alcoves $x\mathbf{a}$ such that some (equivalently, every) minimal gallery joining \mathbf{a} to $x\mathbf{a}$ is in the w -direction. By definition, a gallery $\mathbf{a}_1, \dots, \mathbf{a}_l$ is *in the w -direction* if for each crossing $\mathbf{a}_{i-1} |_H \mathbf{a}_i$, the alcove \mathbf{a}_{i-1} belongs to H^{w-} and \mathbf{a}_i belongs to H^{w+} . By loc. cit. Lemma 5.8, the acute cone $C(\mathbf{a}, w)$ is an intersection of half-spaces:

$$C(\mathbf{a}, w) = \bigcap_{\mathbf{a} \subset H^{w+}} H^{w+}.$$

Proposition 3.0.5. *The set of alcoves \mathfrak{A} is the following union of acute cones of alcoves*

$$(3.0.5) \quad \mathfrak{A} = \bigcup_{w: P \supseteq {}^wB} C(\mathbf{a}, w).$$

Proof. For any root $\alpha \in R$ and $k \in \mathbb{Z}$, let $H_{\alpha, k}^+$ denote the unique half-space for $H_{\alpha, k}$ which contains the base alcove \mathbf{a} . Note that for any $\alpha \in R$ and $w \in W$, we




FIGURE 1. The figure illustrates the notion of P -alcove for G of type C_2 . On the left, $P = w_0 B$, where w_0 is the longest element in W . On the right, $P = s_1 s_2 s_1 P'$ where P' is the standard parabolic $B \cup B s_2 B$. In both cases, the black alcove is the base alcove, the region \mathfrak{A} is in light grey, and the P -alcoves are shown in dark grey.




FIGURE 2. This figure shows P -alcoves for G of type G_2 . On the left, $P = s_2 s_1 s_2 (B \cup B s_1 B)$, on the right, $P = s_2 s_1 s_2 s_1 B$.

have

$$(3.0.6) \quad H_{\alpha, k(\alpha, \mathbf{a})-1}^+ = \begin{cases} H_{\alpha, k(\alpha, \mathbf{a})-1}^{w^+}, & \text{if } \alpha \in w(R^+) \\ H_{\alpha, k(\alpha, \mathbf{a})-1}^{w^-}, & \text{if } \alpha \in w(R^-). \end{cases}$$

Now suppose $w \in W$ satisfies $P \supseteq {}^w B$, or in other words $N \subseteq {}^w U$, or equivalently, $R_N \subseteq w(R^+)$. Then we see using (3.0.6) that

$$C(\mathbf{a}, w) = \bigcap_{\alpha \in w(R^+)} H_{\alpha, k(\alpha, \mathbf{a})-1}^{w+} = \bigcap_{\alpha \in w(R^+)} H_{\alpha, k(\alpha, \mathbf{a})-1}^+,$$

so the union on the right hand side of (3.0.5) is

$$(3.0.7) \quad \bigcup_{w : R_N \subseteq w(R^+)} \bigcap_{\alpha \in w(R^+)} H_{\alpha, k(\alpha, \mathbf{a})-1}^+$$

and in particular is contained in $\bigcap_{\alpha \in R_N} H_{\alpha, k(\alpha, \mathbf{a})-1}^+ = \mathfrak{P}$.

For the opposite inclusion, we set

$$\mathcal{U} = \bigcup_{w : R_N \subseteq w(R^+)} C(\mathbf{a}, w).$$

We will prove the implication

$$(3.0.8) \quad x\mathbf{a} \notin \mathcal{U} \implies x\mathbf{a} \notin \mathfrak{P}$$

by induction on the length ℓ of a minimal gallery $\mathbf{a} = \mathbf{a}_0, \mathbf{a}_1, \dots, \mathbf{a}_\ell = x\mathbf{a}$. If $\ell = 0$, there is nothing to show, so we assume that $\ell > 0$ and that the implication holds for $y\mathbf{a} := \mathbf{a}_{\ell-1}$.

Assume $x\mathbf{a} \notin \mathcal{U}$. There are two cases to consider. If $y\mathbf{a} \notin \mathcal{U}$, then by induction $y\mathbf{a} \notin \mathfrak{P}$. This means that $y\mathbf{a}$ and \mathbf{a} are on opposite sides of a hyperplane $H_{\alpha, k(\alpha, \mathbf{a})-1}$ for some $\alpha \in R_N$. The same then holds for $x\mathbf{a}$, which shows that $x\mathbf{a} \notin \mathfrak{P}$.

Otherwise, $y\mathbf{a} \in \mathcal{U}$, so that $y\mathbf{a}$ belongs to some $C(\mathbf{a}, w)$ with $R_N \subseteq w(R^+)$. Let $H = H_{\beta, m}$ be the wall separating $y\mathbf{a}$ and $x\mathbf{a}$. Since $x\mathbf{a} \notin C(\mathbf{a}, w)$ and $s_{\beta, m}x\mathbf{a} \in C(\mathbf{a}, w)$, we have that $m \in \{0, \pm 1\}$, and $x\mathbf{a} \in C(\mathbf{a}, s_{\beta}w)$. Now, if $s_{\beta} \in W_M$, then $R_N \subseteq s_{\beta}w(R^+)$ and $x\mathbf{a} \in \mathcal{U}$, a contradiction. Thus $\beta \in \pm R_N$, and without loss of generality we may assume $\beta \in R_N$. Now in passing from $y\mathbf{a}$ to $x\mathbf{a}$, we crossed H in the β -opposite direction, where by definition this means for any point a in the interior of \mathbf{a} , $x(a) - y(a) \in \mathbb{R}_{<0}\beta^\vee$. Indeed, if not then since $\beta \in w(R^+)$ the crossing $y\mathbf{a}|_H x\mathbf{a}$ is in the w -direction; in that case $x\mathbf{a}$ belongs to $C(\mathbf{a}, w)$ (since $y\mathbf{a}$ does), a contradiction.

To conclude, we observe that if $\mathbf{a} = \mathbf{a}_0, \dots, \mathbf{a}_\ell$ is a minimal gallery and crosses some $H_{\beta, m}$ with $\beta \in R_N$ in the β -opposite direction, then the terminal alcove \mathbf{a}_ℓ must actually lie outside of \mathfrak{P} (since such a gallery must cross the hyperplane $H_{\beta, k(\beta, \mathbf{a})-1}$). \square

4. REFORMULATION OF THEOREM 2.1.2

In the following reformulation of Theorem 2.1.2, we assume $P = MN$ is semistandard and $x\mathbf{a}$ is a P -alcove. As in Beazley's work [Be], it is easier to work with single cosets xI than with double cosets IxI , and the next result allows us to do just that.

Lemma 4.0.6. *Theorem 2.1.2 is equivalent to the following statement: the map*

$$\phi : ({}^x I \cap I) \times {}^x I_M \cap I_M \rightarrow xI$$

given by $(i, m) \mapsto im\sigma(i)^{-1}$ is surjective. Moreover, it is bijective if $x\mathbf{a}$ is a strict P -alcove. In general, if $[i, xj]$ and $[i', xj']$ belong to the same fiber of ϕ , then xj and xj' are σ -conjugate by an element of ${}^x I_M \cap I_M$.

Proof. It is straightforward to verify that the following diagram with vertical inclusion maps is Cartesian:

$$\begin{array}{ccc} ({}^xI \cap I) \times {}^xI_M \cap I_M & \xrightarrow{\phi} & {}^xI \\ \downarrow & & \downarrow \\ I \times {}^xI_M & \xrightarrow{\phi} & IxI. \end{array}$$

The lemma is now clear by appealing to I -equivariance: each element of IxI is σ -conjugate under I to an element of xI , and ϕ is I -equivariant with respect to the action by σ -conjugation on IxI and the action on $I \times {}^xI_M$ given by $i'[i, m] := [i'i, m]$ for $i' \in I$ and $[i, m] \in I \times {}^xI_M$. \square

We can now prove the portion of Theorem 2.1.2 relating to the fibers of ϕ . Suppose that $[i_1, xj_1], [i_2, xj_2] \in ({}^xI \cap I) \times {}^xI_M \cap I_M$ satisfy $i_1xj_1\sigma(i_1)^{-1} = i_2xj_2\sigma(i_2)^{-1}$. Letting $i := i_2^{-1}i_1$, we see that

$$(4.0.9) \quad x^{-1}ix = j_2\sigma(i)j_1^{-1}.$$

We have the Iwahori decompositions $I = I_{\overline{N}}I_M I_N$ and ${}^xI = {}^xI_{\overline{N}} {}^xI_M {}^xI_N$, where $I_N := N \cap I$ and $I_{\overline{N}} := \overline{N} \cap I$. Using our assumption that $x\mathbf{a}$ is a P -alcove, we deduce

$$(4.0.10) \quad {}^xI \cap I = I_{\overline{N}} ({}^xI_M \cap I_M) {}^xI_N.$$

Write $i = i_- i_0 i_+$, with $i_- \in I_{\overline{N}}$, $i_0 \in {}^xI_M \cap I_M$, and $i_+ \in {}^xI_N$. Using (4.0.9) we get

$$(4.0.11) \quad x^{-1}i_- \cdot x^{-1}i_0 \cdot x^{-1}i_+ = j_2\sigma(i_-) \cdot j_2\sigma(i_0)j_1^{-1} \cdot j_1\sigma(i_+).$$

By the uniqueness of the factorization of elements in $\overline{N} \cdot M \cdot N$, we get

$$(4.0.12) \quad x^{-1}i_- = j_2\sigma(i_-)$$

$$(4.0.13) \quad x^{-1}i_0 = j_2\sigma(i_0)j_1^{-1}$$

$$(4.0.14) \quad x^{-1}i_+ = j_1\sigma(i_+).$$

From (4.0.13), we deduce that xj_1 is σ -conjugate to xj_2 by an element in ${}^xI_M \cap I_M$. This proves the main assertion regarding the fibers of ϕ .

It remains to prove that ϕ is injective when $x\mathbf{a}$ is a strict P -alcove. In that case conjugation by x is strictly expanding (resp. contracting) on $I_{\overline{N}}$ (resp. on I_N). In other words, the condition (2.1.1) hence also (2.1.3) holds with the inclusions replaced by strict inclusions. But then (4.0.12) (resp. (4.0.14)) can hold only if $i_- = 1$ (resp. $i_+ = 1$). Thus, in that case we have $i = i_0 \in {}^xI_M \cap I_M$, and it follows that $[i_1, xj_1] = [i_2, xj_2]$. This proves the desired injectivity of ϕ . \square

5. A VARIANT OF LANG'S THEOREM FOR VECTOR GROUPS

As before, let k denote a finite field with q elements, and let \overline{k} denote an algebraic closure of k . We write σ for the Frobenius automorphism $x \mapsto x^q$ of \overline{k} . In this section we will be concerned with an automorphism τ of \overline{k} , which is required to be either σ or σ^{-1} . By a τ -space (V, Φ) we mean a finite dimensional vector space V over \overline{k} together with a τ -linear map $\Phi : V \rightarrow V$. We do *not* require that Φ be bijective. The category of τ -spaces is abelian and every object in it has finite length.

Let (V, Φ) be a simple object in this category. We claim that V is 1-dimensional (cf. the proof of Lemma 1.3 in [KR]). Since $\ker \Phi$ is a subobject of V , we must have either $\ker \Phi = V$ or $\ker \Phi = 0$. In the first case $\Phi = 0$, every subspace is a subobject, and therefore simplicity forces V to be 1-dimensional. In the second case Φ is bijective, and a subspace W is a subobject $\iff \Phi W = W \iff \Phi^{-1}W = W$. Therefore we may as well assume that $\tau = \sigma$ (since Φ^{-1} is σ -linear if Φ is σ^{-1} -linear). Then by Lang's theorem for general linear groups over k , our τ -space is a direct sum of copies of (\bar{k}, σ) , hence due to simplicity is 1-dimensional.

Lemma 5.0.7. *Let (V, Φ) be a τ -space. Then the k -linear map $v \mapsto v - \Phi(v)$ from V to V is surjective.*

Proof. Filter (V, Φ) so that each successive quotient is 1-dimensional. Since the desired surjectivity follows from surjectivity of the induced map on the associated graded object, we just need to prove surjectivity when V is 1-dimensional. This amounts to the solvability of the equations $x - ax^q = b$ and $x - ax^{1/q} = b$. Solvability of the first equation is obvious, and so too is that of the second after the change of variables $x = y^q$, which leads to the equivalent equation $y^q - ay = b$. \square

Corollary 5.0.8. *Let V_0 be a finite dimensional k -vector space, let $V = V_0 \otimes_k \bar{k}$, and let $M : V \rightarrow V$ be a linear map. Then*

- (1) *for every $w \in V$ there exists $v \in V$ such that $\sigma v - Mv = w$, and*
- (2) *for every $w \in V$ there exists $v \in V$ such that $v - M\sigma v = w$.*

Proof. The second statement follows from the lemma (with $\tau = \sigma$), and the first follows from the lemma (with $\tau = \sigma^{-1}$) after making the change of variables $v = \sigma^{-1}v'$. \square

Remark 5.0.9. *We note that the second statement of the corollary can also be proved in the same way as Lang's theorem. However this method does not handle the first statement of the corollary in the case when M is not bijective.*

6. PROOF OF SURJECTIVITY IN THEOREM 2.1.2

6.1. The method of successive approximations. Again assume that $x\mathfrak{a}$ is a P -alcove. Recall that by Lemma 4.0.6, we need to prove the surjectivity of the map

$$({}^x I \cap I) \times xI_M \rightarrow xI$$

given by $(i, m) \mapsto im\sigma(i)^{-1}$. In other words, given an element of xI , we can σ -conjugate it by an element of ${}^x I \cap I$ into the set xI_M .

Define the normal subgroup $I_n \subset I$, $n = 0, 1, 2, \dots$, to be the n -th principal congruence subgroup of I . More precisely, let \mathcal{G} denote the Bruhat-Tits parahoric \mathfrak{o} -group scheme corresponding to I , so that $\mathcal{G}(\mathfrak{o}) = I$. For $n \geq 0$, let I_n denote the kernel of $\mathcal{G}(\mathfrak{o}) \rightarrow \mathcal{G}(\mathfrak{o}/\epsilon^n \mathfrak{o})$.

Define the normal subgroups $N_n \subset N(\mathfrak{o}) \cap I$, $\bar{N}_n \subset \bar{N}(\mathfrak{o}) \cap I$ and $M_n \subset M(\mathfrak{o}) \cap I$ to be the intersections $I_n \cap N$ resp. $I_n \cap \bar{N}$ resp. $I_n \cap M$. For each $n \geq 0$, we have the Iwahori factorization

$$I_n = M_n N_n \bar{N}_n = \bar{N}_n N_n M_n.$$

Conjugating by x the decomposition $I = I_M I_N I_{\bar{N}}$ yields ${}^x I = {}^x I_M {}^x I_N {}^x I_{\bar{N}}$. By our assumptions on x , we have

$${}^x I \cap I = ({}^x I_M \cap I_M) {}^x I_N I_{\bar{N}}.$$

Similarly, for each $n \geq 0$, we have

$${}^x I_n \cap I_n = ({}^x M_n \cap M_n) {}^x N_n \overline{N}_n.$$

Here we have used the relations

$$(6.1.1) \quad \begin{aligned} {}^x N_n &\subseteq N_n \\ {}^x \overline{N}_n &\supseteq \overline{N}_n. \end{aligned}$$

For the case $n = 0$ each of these relations follows from our assumption that $x\mathbf{a}$ is a P -alcove. In case $n > 0$, the relations can be proved the same way, again using the assumption that $x\mathbf{a}$ is a P -alcove.

The next lemma is a key ingredient in the proof of Theorem 2.1.2. Here and in the remainder of this section we use the following notation: for $h \in G(L)$, a superscript $h-$ stands for conjugation by h , and a superscript $\sigma-$ means application of σ , so in particular, for $g, h \in G(L)$, the symbol ${}^{h\sigma}g$ will stand for $h\sigma(g)h^{-1}$, and ${}^{\sigma h}g$ will stand for $\sigma(h)\sigma(g)\sigma(h^{-1})$.

Lemma 6.1.1. *Fix an element $m \in I_M$ and an integer $n \geq 0$.*

- (i) *Given $i_- \in \overline{N}_n$, there exists $b_- \in \overline{N}_n$ such that $({}^{xm})^{-1}b_-i_- \sigma b_-^{-1} \in \overline{N}_{n+1}$.*
- (ii) *Given $i_+ \in N_n$, there exists $b_+ \in N_n$ such that $b_+i_+ {}^{mx\sigma}b_+^{-1} \in N_{n+1}$.*

Proof. Borrowing the notation of [GHKR], §5.3, the group N possesses a finite separating filtration by normal subgroups

$$N = N[1] \supset N[2] \supset \dots$$

defined as follows. Choose a Borel subgroup B' containing A and contained in P ; use B' to determine a notion of (simple) positive root for A acting on $\text{Lie}(G)$. Let δ'_N be the cocharacter in $X_*(A/Z)$ (where Z denotes the center of G) which is the sum of the B' -fundamental coweights ϖ_α , where α ranges over the simple B' -positive roots for A appearing in $\text{Lie}(N)$. Then let $N[i]$ be the product of the root groups $U_\beta \subset N$ for β satisfying $\langle \beta, \delta'_N \rangle \geq i$. The subgroups $N[i]$ are stable under conjugation by any element in M (as one can check using the Bruhat decomposition of M with respect to the Borel subgroup $B' \cap M$). The successive quotients $N\langle i \rangle := N[i]/N[i+1]$ are abelian (see loc. cit.).

We define $N_n[i] := N_n \cap N[i]$, and $N_n\langle i \rangle := N_n[i]/N_n[i+1]$. We define the groups $\overline{N}[i]$, $\overline{N}\langle i \rangle$, $\overline{N}_n[i]$, and $\overline{N}_n\langle i \rangle$ in an analogous manner.

Now we are ready to prove statement (i). Note that the successive quotients $\overline{N}_n\langle i \rangle$ are abelian, and moreover $\overline{N}_{n+1}\langle i \rangle$ is a subgroup of $\overline{N}_n\langle i \rangle$, and the quotient

$$\overline{N}_n\langle i \rangle / \overline{N}_{n+1}\langle i \rangle$$

is a vector group over the residue field of \mathfrak{o} . Conjugation by $m^{-1} \in I_M$ or x^{-1} preserves \overline{N}_n as well as each $\overline{N}_n[i]$ and $\overline{N}_n\langle i \rangle$ (for x^{-1} , we use (6.1.1) above). Hence the map $b_- \mapsto ({}^{xm})^{-1}b_- \sigma b_-^{-1}$ induces on each vector group $\overline{N}_n\langle i \rangle / \overline{N}_{n+1}\langle i \rangle$ a map like that considered in Corollary 5.0.8 (1). Using that lemma repeatedly on these quotients in a suitable order, we may find $b_- \in \overline{N}_n$ such that

$$({}^{xm})^{-1}b_-i_- \sigma b_-^{-1} \in \overline{N}_{n+1},$$

thus verifying part (i).

Now for part (ii) we use a very similar argument. Conjugation by mx preserves N_n (for x we use (6.1.1) above), as well as each $N_n[i]$ and $N_n\langle i \rangle$. Hence the map $b_+ \mapsto b_+ {}^{mx\sigma}b_+^{-1}$ induces on each vector group $N_n\langle i \rangle / N_{n+1}\langle i \rangle$ a map like that

considered in Corollary 5.0.8 (2). We conclude as in part (i) above. This completes the proof of the lemma. \square

Now we continue with the proof of Theorem 2.1.2. The Iwahori subgroup I has the filtration $I \supset I_1 \supset I_2 \supset I_3 \supset \dots$ by principal congruence subgroups. We want to refine this filtration to a filtration $I = I[0] \supset I[1] \supset I[2] \supset I[3] \supset \dots$ satisfying the following conditions:

- (1) Each $I[r]$ is normal in I .
- (2) Each $I[r]$ is a semidirect product $I\langle r \rangle I[r+1]$, where $I\langle r \rangle$ is either an affine root subgroup (hence one-dimensional over our ground field k) or else contained in $A(\mathfrak{o})$.

One can construct such filtrations directly by inserting suitable terms into the filtration by principal congruence subgroups, or one can just take a generic Moy-Prasad filtration (see below for a discussion of these). In any case, we fix one such filtration (which need not have any special properties relative to our chosen $P = MN$).

We start with a P -alcove $x\mathfrak{a}$ and an element $y \in xI$. We want to find an element $g \in {}^xI \cap I$ such that $gy\sigma(g)^{-1} \in xI_M$. As usual we do this by successive approximations, first σ -conjugating y into $xI_M I[1]$, then into $xI_M I[2]$, and so on. We have to take care that the elements doing the σ -conjugating approach 1 as $r \rightarrow \infty$. Assuming we can do this, if $h^{(r)} \in {}^xI \cap I$ is used to σ -conjugate the appropriate element of $xI_M I[r]$ into $xI_M I[r+1]$, then the convergent product

$$g := \dots h^{(2)} h^{(1)} h^{(0)}$$

has the desired property.

So we need to show that any element $xi_M i[r] \in xI_M I[r]$ is σ -conjugate under ${}^xI \cap I$ to an element of $xI_M I[r+1]$ (and that the σ -conjugators can be taken to be small when r is large). Use item (2) to decompose $i[r]$ as $i\langle r \rangle i[r+1]$. There are two cases. If $I\langle r \rangle \subset A(\mathfrak{o})$, then we can absorb $i\langle r \rangle$ into i_M , showing that our element already lies in $xI_M I[r+1]$.

Otherwise $i\langle r \rangle$ lies in one of the affine root subgroups of I ; write α for the ordinary root obtained as the vector part of our affine root. If α is a root in M , then again we absorb $i\langle r \rangle$ into i_M and do not need to σ -conjugate. Otherwise α is a root in N or \overline{N} , and in either case we may use the Lang theorem variant (i.e. the appropriate statement in Lemma 6.1.1) to produce an element $h \in {}^xI \cap I$ (suitably small when r is large) such that

$$hxi_M i\langle r \rangle \sigma(h)^{-1} = xi_M i',$$

for some $i' \in I[r+1]$. (For example, if $i\langle r \rangle \in N_n$ take $h := {}^x b_+$, where b_+ is the element produced in Lemma 6.1.1 (ii) for $m := i_M$ and $i_+ := mi\langle r \rangle m^{-1}$.) Then

$$hxi_M i\langle r \rangle i[r+1] \sigma(h)^{-1} = xi_M i' (\sigma(h) i[r+1] \sigma(h)^{-1}) \in xI_M I[r+1],$$

as desired. (We used here that $I[r+1]$ is normal in I .) Lemma 6.1.1 produces elements h which are suitably small when r is large, so that we are done, modulo the information on Moy-Prasad filtrations which follows.

6.2. Moy-Prasad filtrations. Our reference for Moy-Prasad filtrations is [MP]. Recall that Moy-Prasad filtrations on I are obtained from points x in the base alcove \mathbf{a} . On the Lie algebra this works as follows. The vector space $\mathfrak{g} \otimes_k k[\epsilon, \epsilon^{-1}]$ is graded by the group $X^*(A) \oplus \mathbb{Z}$ (since \mathfrak{g} is graded by $X^*(A)$ and $k[\epsilon, \epsilon^{-1}]$ is graded by \mathbb{Z}). (For the moment k is any field.) The pair $(x, 1)$ gives a homomorphism $X^*(A) \oplus \mathbb{Z} \rightarrow \mathbb{R}$, which we use to obtain an \mathbb{R} -grading on $\mathfrak{g} \otimes_k k[\epsilon, \epsilon^{-1}]$, as well as an associated \mathbb{R} -filtration. We also obtain an \mathbb{R} -filtration on the completion $\mathfrak{g}(F)$ of $\mathfrak{g} \otimes_k k[\epsilon, \epsilon^{-1}]$. Thus, for $r \in \mathbb{R}$ the subspace $\mathfrak{g}(F)_{\geq r}$ is the completion of the direct sum of the affine weight spaces of weight (with respect to $(x, 1)$) greater than or equal to r , which for the affine weight space $\epsilon^n \mathbf{a}$ means that $n \geq r$, and for an affine weight space $\epsilon^n \mathfrak{g}_\alpha$ (α being an ordinary root) means that $\alpha(x) + n \geq r$. Of course $\mathfrak{g}(F)_{\geq 0}$ is the Iwahori subalgebra obtained as the Lie algebra of I ¹. It is clear that $[\mathfrak{g}(F)_{\geq r}, \mathfrak{g}(F)_{\geq s}] \subset \mathfrak{g}(F)_{\geq r+s}$, from which it follows that $\mathfrak{g}(F)_{\geq r}$ is an ideal in $\mathfrak{g}(F)_{\geq 0}$ whenever r is non-negative.

When r is non-negative, the Moy-Prasad subgroups $G(F)_{\geq r}$ of $G(F)$ are by definition the subgroups generated by suitable subgroups of $A(\mathfrak{o})$ and of the various root subgroups, in such a way that the Lie algebra of $G(F)_{\geq r}$ ends up being $\mathfrak{g}(F)_{\geq r}$. In characteristic 0 the fact that $\mathfrak{g}(F)_{\geq r}$ is an ideal in $\mathfrak{g}(F)_{\geq 0}$ implies that $G(F)_{\geq r}$ is normal in $I = G(F)_{\geq 0}$. Moy and Prasad prove normality in the general case from other considerations. In our present situation, where G is split, it is straightforward to prove the normality using commutator relations for the various affine root groups $U_{\alpha+n}$ in $G(F)$.

What does it mean for x to be a *generic* element in the base alcove? For an arbitrary point x in the standard apartment it may accidentally happen that the homomorphism $(x, 1) : X^*(A) \oplus \mathbb{Z} \rightarrow \mathbb{R}$ sends two distinct affine weights occurring in $\mathfrak{g} \otimes_k k[\epsilon, \epsilon^{-1}]$ to the same real number. When such an accident never occurs, we say that x is generic. The set of non-generic points in the standard apartment is a locally finite union of affine hyperplanes, including all the affine root hyperplanes, but also those obtained by setting any difference of roots equal to an integer. In the case of $\mathrm{SL}(2)$, all points in the base alcove but its midpoint are generic. In general one can at least say that the set of generic points in the base alcove is non-empty and open. When x is generic, then going down the Moy-Prasad filtration strips away affine weight spaces, one-by-one, just as we want.

6.3. A refinement. It is clear that in case ${}^x I_M = I_M$, we can do better: we can σ -conjugate ${}^x I_M$ to x using an element of I_M . To see this we adapt the proof of Lang's theorem to prove the surjectivity of the map $I_M \rightarrow I_M$ given by $h \mapsto h^{-1} {}^x \sigma h$. Indeed, I_M has a filtration by normal subgroups which are stabilized by $\mathrm{Ad}(x)$, such that our map induces on the successive quotients a finite étale surjective map (take the Moy-Prasad filtration on I_M corresponding to the barycenter of the alcove in the reduced building for $M(L)$ corresponding to I_M). Using the surjectivity just proved, given $i \in I_M$ we find an $h \in I_M$ solving the equation $xi x^{-1} = h^{-1} {}^x \sigma h$. We then have $h(xi)\sigma(h)^{-1} = x$. Thus, we have proved the following proposition.

¹Warning: This description is incompatible with the normalization of the correspondence between alcoves and Iwahori subgroups we are using in this paper: it turns out $G(F)_{\geq 0}$ is really "opposite" to our Iwahori I . To get our I , we should instead define $\mathfrak{g}(F)_{\geq r}$ to be the completion of the sum of the affine weight spaces of weight (with respect to $(x, -1)$) less than or equal to $-r$.

Proposition 6.3.1. *Suppose $x \in \widetilde{W}_M$ is such that there exists a semistandard parabolic subgroup $P = MN$ having the property that ${}^x I_N \subseteq I_N$, i. e. such that $x\mathbf{a}$ is a P -alcove. Then any element of xI is σ -conjugate to an element of xI_M using an element of ${}^x I \cap I$. If moreover, ${}^x I_M = I_M$, then we may σ -conjugate any element of xI to x , using an element of ${}^x I \cap I$.*

Given an element $x \in \widetilde{W}_M$ such that ${}^x I_M = I_M$, in general there is no parabolic $P = MN$ such that ${}^x I_N \subseteq I_N$ and ${}^{x^{-1}} I_{\overline{N}} \subseteq I_{\overline{N}}$ (see also the discussion after Definition 7.2.3 below). However, when M is adapted to I in the sense of Definition 13.2.1, such P does exist, as is shown in Proposition 13.2.2.

7. REVIEW OF σ -CONJUGACY CLASSES

7.1. Classification of σ -conjugacy classes. We recall the description of the set $B(G)$ of σ -conjugacy classes in $G(L)$; for details see [K1], [K2] 5.1, and [K4] 1.3. We denote by Λ_G the quotient of $X_*(A)$ by the coroot lattice; this is the algebraic fundamental group of G . We can identify Λ_G with the group of connected components of the loop group $G(L)$. Let $\eta_G: G(L) \rightarrow \Lambda_G$ be the natural surjective homomorphism, as constructed in [K2], §7 and denoted there by ω_G ; it is sometimes called the *Kottwitz map*. Analogously, we denote by Λ_M the quotient of $X_*(A)$ by the coroot lattice for M , and by η_M the corresponding homomorphism.

If $P = MN$ is a standard parabolic subgroup of G with unipotent radical N and M the unique Levi containing A , then the set Δ of simple roots for G decomposes as the disjoint union of Δ_M and Δ_N , where Δ_M is the set of simple roots of M , and Δ_N is the set of those simple roots for G which occur in the Lie algebra of N . We write A_P (or A_M) for the connected component of the center of M , and we let \mathfrak{a}_P denote the real vector space $X_*(A_P) \otimes \mathbb{R}$. As usual, P determines an open chamber \mathfrak{a}_P^+ in \mathfrak{a}_P defined by

$$\mathfrak{a}_P^+ = \{v \in \mathfrak{a}_P : \langle \alpha, v \rangle > 0, \text{ for all } \alpha \in \Delta_N\}.$$

The composition $X_*(A_P) \hookrightarrow X_*(A) \rightarrow \Lambda_M$, when tensored with \mathbb{R} , yields a canonical isomorphism $\mathfrak{a}_P \cong \Lambda_M \otimes \mathbb{R}$. Let Λ_M^+ denote the subset of elements in Λ_M whose image under $\Lambda_M \otimes \mathbb{R} \cong \mathfrak{a}_P$ lies in \mathfrak{a}_P^+ .

Let \mathbb{D} be the diagonalizable group over F with character group \mathbb{Q} . As in [K1], an element $b \in G(L)$ determines a homomorphism $\nu_b: \mathbb{D} \rightarrow G$ over L , whose $G(L)$ -conjugacy class depends only on the σ -conjugacy class $[b] \in B(G)$. We can assume this homomorphism factors through our torus A , and that the corresponding element $\overline{\nu}_b \in X_*(A)_{\mathbb{Q}}$ is dominant. Then $b \mapsto \overline{\nu}_b$ is called the *Newton map* (relative to the group G). Recall that $b \in G(L)$ is called *basic* if ν_b factors through the center $Z(G)$ of G .

We shall use some properties of the Newton map. We can identify the quotient $X_*(A)_{\mathbb{Q}}/W$ with the closed dominant chamber $X_*(A)_{\mathbb{Q}}^+$. The map

$$(7.1.1) \quad \begin{aligned} B(G) &\rightarrow X_*(A)_{\mathbb{Q}}^+ \times \Lambda_G \\ b &\mapsto (\overline{\nu}_b, \eta_G(b)) \end{aligned}$$

is injective ([K2], 4.13).

The Newton map is functorial, such that we have a commutative diagram

$$(7.1.2) \quad \begin{array}{ccc} B(M) & \longrightarrow & B(G) \\ \downarrow & & \downarrow \\ X_*(A)_{\mathbb{Q}}/W_M \times \Lambda_G & \longrightarrow & X_*(A)_{\mathbb{Q}}/W \times \Lambda_G \end{array}$$

and moreover the vertical arrows, given by “(Newton point, Kottwitz point)”, are *injections*. Indeed, the right vertical arrow is the injection (7.1.1). To show the left vertical arrow is injective, it is enough to prove that if $b_1, b_2 \in M(L)$ have the same Newton point and the same image under η_G , then they have the same image under η_M . We may assume that $b_1, b_2 \in \widetilde{W}_M$ (see Corollary 7.2.2 below); for $i = 1, 2$ write $b_i = \epsilon^{\lambda_i} w_i$ for $\lambda_i \in X_*(A)$ and $w_i \in W_M$. Let Q^\vee (resp. Q_M^\vee) denote the lattice generated by the coroots of G (resp. M) in $X_*(A)$. The equality $\eta_G(b_1) = \eta_G(b_2)$ means that $\lambda_1 - \lambda_2 \in Q^\vee$. The equality $\bar{v}_{b_1} = \bar{v}_{b_2}$ implies that $\lambda_1 - \lambda_2 \in Q_M^\vee \otimes \mathbb{R}$. It follows that $\lambda_1 - \lambda_2 \in Q_M^\vee$, and this is what we wanted to prove.

The following lemma is a direct consequence of the commutativity of the diagram above.

Lemma 7.1.1. *Let $M \subset G$ be a Levi subgroup containing A . If $[b']_M \subset [b]$ for some $b' \in M(L)$, then $\bar{v}_b = \bar{v}_{b', G\text{-dom}}$ as elements of $X_*(A)_{\mathbb{Q}}^+$.*

Here $\bar{v}_{b'}$ is the Newton point of b' (viewed as an element of $M(L)$) and $\bar{v}_{b', G\text{-dom}}$ denotes the unique G -dominant element of $X_*(A)_{\mathbb{Q}}$ in its W -orbit.

Next we define the following subsets of $X_*(A)_{\mathbb{Q}}^+$: the subset \mathcal{N}_G consists of all Newton points \bar{v}_b for $b \in B(G)$, and \mathcal{N}_M^+ consists of the images of elements of Λ_M^+ , under the map $\Lambda_M \rightarrow X_*(A_M)_{\mathbb{Q}} \hookrightarrow X_*(A)_{\mathbb{Q}}$. We have the equality

$$(7.1.3) \quad \mathcal{N}_G = \coprod_{P=MN} \mathcal{N}_M^+,$$

the union ranging over all standard parabolic subgroups of G .

This equality results from two facts. First, we are taking the Newton points associated to elements of $B(G)$ and making use of the decomposition of $B(G)$

$$B(G) = \coprod_P B(G)_P,$$

where P ranges over standard parabolic subgroups and $B(G)_P$ is the set of elements $[b] \in B(G)$ such that $\bar{v}_b \in \mathfrak{a}_P^+$ (see [K1, K2]); note that elements in $B(G)_P$ can be represented by basic elements in $M(L)$ ([K2], 5.1.2). Second, for b a basic element in $M(L)$ (representing e. g. an element in $B(G)_P$) its Newton point \bar{v}_b is the image of $\eta_M(b) \in \Lambda_M$ under the canonical map

$$(7.1.4) \quad \Lambda_M = X^*(Z(\widehat{M})) \rightarrow X^*(Z(\widehat{M}))_{\mathbb{R}} = X_*(Z(M))_{\mathbb{R}} \hookrightarrow X_*(A)_{\mathbb{R}}.$$

This follows from the characterization of \bar{v}_b in [K1], 4.3 (applied to M in place of G), together with (7.1.2).

Remark 7.1.2. *The right hand side in (7.1.3) is easy to enumerate for any given group (with the aid of a computer). This fact makes feasible our computer-aided verifications of our conjectures relating to the non-emptiness of $X_x(b)$, see section 9. Moreover, the injectivity of (7.1.1) together with (7.1.3) gives a concrete way to check whether two elements in $G(L)$ are σ -conjugate.*

7.2. Construction of standard representatives for $B(G)$. Here we will define the *standard representatives* of σ -conjugacy classes in the extended affine Weyl group. First note that the map $G(L) \rightarrow B(G)$ induces a map $\widetilde{W} \rightarrow B(G)$. Our goal is to find special elements in \widetilde{W} which parametrize the elements of $B(G)$.

Denote by $\Omega_G \subset \widetilde{W}(G)$ the subgroup of elements of length 0. Let $G(L)_b$ resp. $B(G)_b$ denote the set of basic elements resp. basic σ -conjugacy classes in $G(L)$. In the following lemma we recollect some standard facts relating the Newton map to the homomorphism $\eta_G : G(L) \rightarrow \Lambda_G$. The connection between the two stems from fact that if $b \in G(L)$ is basic, then the Newton point $\bar{\nu}_b \in X_*(Z(G))_{\mathbb{R}}$ is the image of $\eta_G(b) \in \Lambda_G$ under the canonical map $\Lambda_G = X^*(Z(\widehat{G})) \rightarrow X_*(Z(G))_{\mathbb{R}} \hookrightarrow X_*(A)_{\mathbb{R}}$ (see (7.1.4)).

Lemma 7.2.1. (i) *The map η_G induces a bijection $B(G)_b \xrightarrow{\sim} \Lambda_G$.*
 (ii) *Elements in $\Omega_G \subset G(L)$ are basic, and the map η_G induces a bijection $\Omega_G \xrightarrow{\sim} \Lambda_G$.*
 (iii) *The canonical map $\Omega_G \rightarrow B(G)_b$ is a bijection.*

Proof. First suppose $b \in \Omega_G$. For sufficiently divisible $N > 1$, the element b^N is a translation element which preserves the base alcove, hence belongs to $X_*(Z(G))$. The characterization of ν_b in [K1], 4.3, then shows that b is basic, proving the first statement in (ii). For part (i), recall that an isomorphism is constructed in loc. cit. 5.6, and this is shown to be induced by η_G in [K2], 7.5. Since η_G is trivial on I and $W_{\text{aff}} \subset G_{\text{sc}}(L)$, (i) and the Bruhat-Tits decomposition

$$G(L) = \coprod_{w\tau \in W_{\text{aff}} \rtimes \Omega_G} Iw\tau I$$

imply that the composition

$$\Omega_G \longrightarrow G(L)_b \xrightarrow{\eta_G} \Lambda_G$$

is surjective. Since this composition is easily seen to be injective, (ii) holds. Part (iii) follows using (i-ii). \square

Here is a slightly different point of view of the lemma: The basic conjugacy classes are in bijection with Λ_G , the group of connected components of the ind-scheme $G(L)$ (or the affine flag variety), and the bijection is given by just mapping each basic σ -conjugacy class to the connected component it lies in. The key point here is that the Kottwitz homomorphism agrees with the natural map $G(L) \rightarrow \pi_0(G(L)) = \Lambda_G$; see [K1], [PR] §5.

As a consequence of the lemma (applied to G and its standard Levi subgroups), we have the following corollary.

Corollary 7.2.2. *The map $\widetilde{W} \rightarrow B(G)$ is surjective.*

Definition 7.2.3. *For $[b] \in B(G)_P \subset B(G)$, we call the representative in $\Omega_M \subset \widetilde{W}$ which we get from Lemma 7.2.1 (iii) the standard representative of $[b]$. Here standard refers back to our particular choice B of Borel subgroup. If we made a different choice of Borel subgroup containing A , we would get a different standard representative; all such representatives will be referred to as semistandard.*

The standard representative $b = \epsilon^\nu v$ hence satisfies

- (1) $b \in \widetilde{W}_M$, i. e. $v \in W_M$,

$$(2) bI_M b^{-1} = I_M.$$

Remark 7.2.4. Let $x \in \Omega_G$ and write $x = \epsilon^\lambda w$ with $\lambda \in X_*(A)$ and $w \in W$. Then λ is the (unique) dominant minuscule coweight whose image in Λ_G coincides with that of x . Indeed, since x preserves the base alcove \mathfrak{a} , the transform of the origin by x , namely λ , lies in the closure of the base alcove. This is what it means to be dominant and minuscule.

Now consider standard (semistandard is not enough) $P = MN$ and $x \in \Omega_M$. Write $x = \epsilon^\lambda w_M$ with $\lambda \in X_*(A)$ and $w_M \in W_M$. We know that λ is M -dominant and M -minuscule. We claim that $x\mathfrak{a}$ is a P -alcove iff λ is dominant. Indeed, $x\mathfrak{a}$ is a P -alcove iff $xI_N x^{-1} \subset I_N$. Now $w_M I_N w_M^{-1} = I_N$, because P was assumed standard. So $x\mathfrak{a}$ is a P -alcove iff $\epsilon^\lambda I_N \epsilon^{-\lambda} \subset I_N$ iff $\alpha(\lambda) \geq 0$ for all $\alpha \in R_N$ iff $\alpha(\lambda) \geq 0$ for all $\alpha > 0$.

Example 7.2.5. Let $G = GL_n$, let A be the diagonal torus, and let B be the Borel group of upper triangular matrices. In this case, the Newton map is injective. See [K4], in particular the last paragraph of section 1.3. We can view the Newton vector ν of a σ -conjugacy class $[b]$ as a descending sequence $a_1 \geq \dots \geq a_n$ of rational numbers, satisfying an integrality condition. The standard parabolic subgroup $P = MN$ is given by the partition $n = n_1 + \dots + n_r$ of n such that the a_i in each corresponding batch are equal to each other, and such that the a_i in different batches are different. The standard representative is (represented by) the block diagonal matrix with r blocks, one for each batch of entries, where the i -th block is

$$\begin{pmatrix} 0 & \epsilon^{k_i+1} I_{k'_i} \\ \epsilon^{k_i} I_{n_i-k'_i} & 0 \end{pmatrix} \in GL_{n_i}(F).$$

Here we write the entry $a_{n_1+\dots+n_{i-1}+1} = \dots = a_{n_1+\dots+n_i}$ of the i -th batch as $k_i + \frac{k'_i}{n_i}$ with $k_i, k'_i \in \mathbb{Z}$, $0 \leq k'_i < n_i$, which is possible by the integrality condition, and I_ℓ denotes the $\ell \times \ell$ unit matrix. It follows from the definitions that $k_i \geq k_{i+1}$ for all $i = 1, \dots, r-1$. We see that the standard representative x of $[b]$ has dominant translation part if and only if for all i with $k'_{i+1} \neq 0$ we have $k_i > k_{i+1}$. Furthermore, this is equivalent to $x\mathfrak{a}$ being a P -alcove. If these conditions are satisfied, then $x\mathfrak{a}$ is a fundamental P -alcove in the sense of Definition 13.1.2.

8. PROOFS OF COROLLARY 2.1.3(B) AND THEOREM 2.1.4

Assume $P = MN$ is semistandard and $x\mathfrak{a}$ is a P -alcove. There is a commutative diagram

$$(8.0.1) \quad \begin{array}{ccc} I_M x I_M /_\sigma I_M & \xrightarrow{\sim} & I x I /_\sigma I \\ \cong \downarrow & & \cong \downarrow \\ \coprod_{[b'] \in B(M)_x} J_{b'}^M \backslash X_x^M(b') & \longrightarrow & \coprod_{[b] \in B(G)_x} J_b^G \backslash X_x^G(b). \end{array}$$

Here, for $[b'] \in B(M)_x$ we choose once and for all a representative $b' \in M(L)$; for $[b] \in B(G)_x$ we also choose once and for all a representative $b \in G(L)$. If under $B(M)_x \rightarrow B(G)_x$, $[b'] \mapsto [b]$, then choose once and for all $c \in G(L)$ such that

$c^{-1}b\sigma(c) = b'$. In that case our choices yield the map

$$\begin{aligned} J_{b'}^M \backslash X_x^M(b') &\rightarrow J_b^G \backslash X_x^G(b) \\ m &\mapsto cm. \end{aligned}$$

We have now defined the bottom horizontal arrow.

Next we define the right vertical arrow. Let an element of $IxI/\sigma I$ be represented by $y \in IxI$. There is a unique $[b] \in B(G)_x$ such that $y \in [b]$. Write $y = g^{-1}b\sigma(g)$ for some $g \in G(L)$. Then the right vertical map associates to $[y] = [g^{-1}b\sigma(g)]$ the J_b^G -orbit of $gI \in X_x^G(b)$. The left vertical arrow is defined similarly. It is easy to check that both vertical arrows are bijective. It is also clear that the diagram commutes. The bijectivity of the top horizontal arrow (Corollary 2.1.3(a)) thus implies the surjectivity of the map $B(M)_x \rightarrow B(G)_x$ (in Corollary 2.1.3(b)).

We now prove that $B(M)_x \rightarrow B(G)_x$ is also injective. Given $b \in M(L)$, regard its Newton point \bar{v}_b^M as an element in $X_*(A)_{\mathbb{Q}}^+$, which denotes here the set of M -dominant elements of $X_*(A)_{\mathbb{Q}}$. The map

$$\begin{aligned} B(M) &\rightarrow X_*(A)_{\mathbb{Q}}^+ \times \Lambda_M \\ b &\mapsto (\bar{v}_b^M, \eta_M(b)) \end{aligned}$$

is injective, see (7.1.1). Now suppose $b_1, b_2 \in B(M)_x$ have the same image in $B(G)_x$. Since $\eta_M(b_1) = \eta_M(x) = \eta_M(b_2)$, by the preceding remark it is enough to show that $\bar{v}_{b_1}^M = \bar{v}_{b_2}^M$. We claim that our assumption on x forces each $\bar{v}_{b_i}^M$ to be not only M -dominant, but G -dominant. Indeed, b_i is σ -conjugate in $M(L)$ to an element in $I_M x I_M$, and since ${}^x(N \cap I) \subseteq N \cap I$, it follows that the isocrystal

$$(\mathrm{Lie} N(L), \mathrm{Ad}(b_i) \circ \sigma)$$

comes from a crystal (i.e., there is some \mathfrak{o} -lattice in $\mathrm{Lie} N(L)$ carried into itself by the σ -linear map $\mathrm{Ad}(b_i) \circ \sigma$; in fact, when b_i itself lies in $I_M x I_M$, the lattice $\mathrm{Lie} N(L) \cap I$ does the job). The slopes of any crystal are non-negative, which means in this situation that $\langle \alpha, \bar{v}_{b_i}^M \rangle \geq 0$ for all $\alpha \in R_N$. This proves our claim. Now since $\bar{v}_{b_1}^M$ and $\bar{v}_{b_2}^M$ are conjugate under W (cf. (7.1.2)) they are in fact equal. This completes the proof of Corollary 2.1.3(b).

In light of the diagram (8.0.1), Theorem 2.1.4 follows from Corollary 2.1.3. \square

9. CONSEQUENCES FOR AFFINE DELIGNE-LUSZTIG VARIETIES

In this section we present various consequences of Theorem 2.1.4, and also some conjectures, relating to the non-emptiness and dimension of $X_x^G(b)$. We prove some parts of our conjectures. Our conjectures have been corroborated by ample computer evidence. The computer calculations were done using the ‘‘generalized superset method’’, that is, the algorithm implicit in Theorem 11.2.1. This will be discussed in section 11.

9.1. Translation elements $x = \epsilon^\lambda$. Let us examine the non-emptiness of $X_x(b)$ in a very special case.

Corollary 9.1.1. *Suppose $x = \epsilon^\lambda$. Then $X_x(b) \neq \emptyset$ if and only if $[b] = [\epsilon^\lambda]$ in $B(G)$.*

Proof. There is a choice of Borel $B' = AU'$ such that $x\mathbf{a}$ is a B' -alcove (λ is B' -dominant for an appropriate choice of B'). Thus, by Theorem 2.1.4 with $M = A$,

we see $X_x^G(b) \neq \emptyset$ if and only if b is σ -conjugate to a translation ϵ^ν for $\nu \in X_*(A)$, and $X_x^A(\epsilon^\nu) \neq \emptyset$. But the latter inequation holds if and only if $\lambda = \nu$. \square

Remark 9.1.2. *As G. Lusztig pointed out, the Corollary has a simple direct proof in the special case where G is simply-connected and $b = 1$. Let $x = \epsilon^\lambda$ and suppose λ belongs to the coroot lattice. Suppose $g^{-1}\sigma(g) \in IxI$. Since the affine flag variety is of ind-finite type, the Iwahori subgroup gI is fixed by σ^r for some $r > 0$. Thus, $g^{-1}\sigma^r(g) \in I$. On the other hand, $g^{-1}\sigma^r(g) \in IxI \cdots IxI$ (product of r copies of IxI), which since the lengths add is just $I\epsilon^{r\lambda}I$. This intersects I only if $\lambda = 0$.*

9.2. A necessary condition for the non-emptiness of $X_x(b)$.

Proposition 9.2.1. *Fix a σ -conjugacy class $[b]$ in G with Newton vector $\bar{\nu}_b$, and an element $x \in \widetilde{W}$. If $X_x^G(b) \neq \emptyset$, then the following holds: if $P = MN$ is a semistandard parabolic subgroup such that $x\mathbf{a}$ is a P -alcove, then $\eta_G(x) = \eta_G(b)$ and*

$$(9.2.1) \quad \eta_M(x) \in \eta_M(W\bar{\nu}_b \cap \mathcal{N}_M),$$

where \mathcal{N}_M denotes the image of $B(M)$ in $X_*(A)_{\mathbb{Q}}^+$ under the Newton map.

Proof. It is clear that $X_x^G(b)$ can be non-empty only if $\eta_G(x) = \eta_G(b)$. What is the meaning of our second condition (9.2.1)? The set $W\bar{\nu}_b \cap \mathcal{N}_M$ consists of the finite set of Newton points $\bar{\nu}_{b'}^M$, for $b' \in M(L)$, which are W -conjugate to $\bar{\nu}_b$. Our condition (9.2.1) means that x has the same value under η_M as an element $b' \in M(L)$ with $\bar{\nu}_{b'}^M \in W\bar{\nu}_b$. By the injectivity of the left vertical arrow of (7.1.2), for a fixed $[b]$ there are only *finitely many* σ -conjugacy classes $[b'] \in B(M)$ such that $\bar{\nu}_{b'}^M \in W\bar{\nu}_b$ and $\eta_G(b') = \eta_G(b)$. Thus the condition that $\eta_M(x) = \eta_M(b')$ for some such b' is a condition which we can check with a computer.

That said, the condition (9.2.1) is a direct consequence of Theorem 2.1.4. Indeed, we know from part (a) of that theorem that $[b] = [b']$ for some $b' \in M(L)$, and that $X_x^M(b') \neq \emptyset$, which implies in turn that $\eta_M(x) = \eta_M(b')$. Lemma 7.1.1 then shows that $\bar{\nu}_{b'}^M \in W\bar{\nu}_b$, as desired. \square

Note that Proposition 9.2.1 implies that for fixed b and *proper* parabolic subgroup P , there are only finitely many x such that $x\mathbf{a}$ is a P -alcove and for which $X_x(b)$ can be non-empty.

If b is basic, then the statement of Proposition 9.2.1 simplifies. We will consider the basic case in the next subsection.

Proposition 9.2.1 provides an obstruction to the non-emptiness of affine Deligne-Lusztig varieties: (9.2.1) must hold whenever $x\mathbf{a}$ is a P -alcove. In the case where $[b]$ is basic, it seems reasonable to expect that this is the only obstruction; see Conjecture 9.3.2 below. In the general case, it is clear that there are additional obstructions. If b is a translation element, then from Theorem 6.3.1 in [GHKR] we see that whenever $X_x(b) \neq \emptyset$, there exists $w \in W$ such that $x \geq {}^w b$ in the Bruhat order. (For general b , one can obtain a similar criterion by passing to a totally ramified extension of L where b splits.) This condition implies in particular that for all projections to affine Grassmannians, the corresponding affine Deligne-Lusztig variety is non-empty, but is stronger than that. However, as the following example shows, there are still more elements x which give rise to an empty affine Deligne-Lusztig variety.

Example 9.2.2. Let $G = SL_3$, $b = \epsilon^\lambda$ where $\lambda = (2, 0, -2)$. Let $x = s_{01210120120} = \epsilon^{(3,1,-4)} s_{121}$ (we write s_{12} for $s_1 s_2$ etc.). Then $x \geq b$ (a reduced expression for b is $s_{01210121}$), and $x\mathbf{a}$ is not a P -alcove for any proper parabolic subgroup P . However, $X_x(b) = \emptyset$. (Cf. Figure 3.24 in [Re1] which shows the situation for this b .)

9.3. Non-emptiness of $X_x(b)$ for b basic. In this subsection, let b be basic in $G(L)$. In that case Lemma 7.1.1 and the injectivity of the left vertical arrow of (7.1.2) imply the following: if $[b] \cap M(L) \neq \emptyset$ for some semistandard Levi subgroup $M \subseteq G$, then Lemma 7.1.1 shows that $[b] \cap M(L)$ is a single σ -conjugacy class inside M with the same Newton vector as the Newton vector of $[b]$ with respect to G . (On the other hand, the standard representative of $[b]$ with respect to G is not necessarily an element of M , and in particular is in general different from the standard representative with respect to M .)

Applying Proposition 9.2.1 to the basic case, we get

Corollary 9.3.1. Let $[b]$ be basic. Suppose $P = MN$ is a semistandard parabolic subgroup such that $x\mathbf{a}$ is a P -alcove. Then $X_x(b) = \emptyset$, unless $[b]$ meets $M(L)$ and $\eta_M(x) = \eta_M(\bar{v}_b)$.

Let us once again emphasize that $\eta_M(\bar{v}_b)$ is really an abbreviation; here it stands for the value under η_M for the unique σ -conjugacy class $[b'] \in B(M)$ which satisfies $\eta_G(b') = \eta_G(b)$ and $\bar{v}_{b'}^M = \bar{v}_b$.

Conjecture 9.3.2. In the corollary, the opposite implication holds as well. In other words, when b is basic, $X_x(b)$ is empty if and only if there exists a semistandard $P = MN$ such that $x\mathbf{a}$ is a P -alcove, and $\eta_M(x) \neq \eta_M(\bar{v}_b)$.

This conjecture can be checked in the rank 2 cases “by hand”, and in higher rank cases, computer experiments provide further support for the conjecture: it has been confirmed for the simply connected groups (i. e. for $b = 1$) of type A_3 and x of length ≤ 27 , of type A_4 and x of length ≤ 17 and of type C_3 and x of length ≤ 23 , and in several cases with b basic, but different from 1.

In the remainder of this subsection we discuss some sufficient conditions for the non-emptiness of $X_x(b)$, when b is basic.

Lemma 9.3.3. Let $x = \epsilon^\lambda w \in \widetilde{W}$ be an element which is not contained in any Levi subgroup. Then

$$X_x(b) \neq \emptyset \iff \eta_G(x) = \eta_G(b).$$

Here by *not contained in any Levi subgroup*, we mean that no representative of x in $N_G(A)(L)$ is contained in a Levi subgroup of G associated with a proper semistandard parabolic subgroup of G . Since we consider only Levi subgroups containing the fixed maximal torus A , their (extended affine) Weyl groups are subgroups of the (extended affine) Weyl group of G . In terms of Weyl groups we can state the condition as: the finite part w of x is not contained in any conjugate of a proper parabolic subgroup of W .

If w belongs to the Coxeter conjugacy class of W , then the condition is satisfied. For the symmetric groups, i. e. if G is of type A_n , the converse is also true, as one sees using disjoint cycle decompositions. For all other types, however, there exist other conjugacy classes which do not meet any (standard) parabolic subgroup of W (see for instance [GP], where these conjugacy classes are called cuspidal; some authors call them elliptic).

Before beginning the proof we note that similar considerations can be found in [KR, Proposition 4.1] and [Re1, §3.3.4].

Proof. As before, it is clear that $X_x(b) \neq \emptyset$ implies $\eta_G(x) = \eta_G(b)$. On the other hand, given the latter condition, we will show that x is itself σ -conjugate to b , in other words that the Newton vector of x is $\bar{\nu}_b$. Our assumption ensures that x is in the right connected component of $G(L)$, so that we only need to prove that x is basic.

In order to show that x is basic, we prove that the Newton vector of x , $\bar{\nu}_x = \frac{1}{N} \sum_{i=0}^{N-1} w^i \lambda \in X_*(A)_{\mathbb{Q}}$ is W -invariant. (Here N denotes the order of w in W .) The point $\bar{\nu}_x$ lies in (the closure of) some Weyl chamber, and hence its stabilizer is generated by a subset of the set of simple reflections for this chamber, and hence is the Weyl group of some Levi subgroup (or of all of G). On the other hand, w is contained in this stabilizer, and so our assumption gives us that the stabilizer of $\bar{\nu}_x$ is in fact W . \square

As the proof shows, if G is semi-simple the elements $x \in \widetilde{W}$ which are not contained in any Levi have finite order in \widetilde{W} . Cf. [GHKR] Prop. 7.3.1.

Now let $x \in \widetilde{W}$. If x is not contained in any Levi, then we understand whether $X_x(b) = \emptyset$ by the lemma. In general, there is a minimal semistandard Levi subgroup M_- containing x , and a minimal semistandard Levi subgroup $M_+ \supseteq M_-$ such that $x\mathbf{a}$ is a P_+ -alcove for some semistandard parabolic subgroup P_+ with Levi part M_+ . Both of these statements follow from [Bo], Prop. 14.22, which says that for (semistandard) parabolic subgroups P_1, P_2 , the subgroup $(P_1 \cap P_2)R_u P_1$ is again a (semistandard) parabolic subgroup; it has Levi part $M_1 \cap M_2$. There may be more than one parabolic P_+ with Levi part M_+ for which $x\mathbf{a}$ is a P_+ -alcove, and of course, we may have $M_+ = P_+ = G$.

We then have, by Theorem 2.1.4, (and assuming that $[b]$ meets M_+ , because otherwise $X_x^G(b) = \emptyset$, again by Theorem 2.1.4),

$$X_x^G(b) \neq \emptyset \iff X_x^{M_+}(b) \neq \emptyset \implies \eta_{M_+}(x) = \eta_{M_+}(\bar{\nu}_b).$$

Further, the lemma gives us (assuming that $[b]$ meets M_-)

$$(X_x^{M_+}(b) \neq \emptyset \iff) X_x^{M_-}(b) \neq \emptyset \iff \eta_{M_-}(x) = \eta_{M_-}(\bar{\nu}_b).$$

The condition $\eta_{M_-}(x) = \eta_{M_-}(\bar{\nu}_b)$ is quite restrictive; and it becomes more restrictive the smaller M_- is.

So, in terms of proving Conjecture 9.3.2, the case which remains to consider is the case of x which satisfy the following two conditions: (i) either $[b]$ does not meet M_- or it does and $X_x^{M_-}(b) = \emptyset$, and (ii) $[b]$ meets M_+ and $\eta_{M_+}(x) = \eta_{M_+}(\bar{\nu}_b)$. The conjecture predicts that in this case $X_x^{M_+}(b) \neq \emptyset$.

9.4. Relation with Reuman's conjecture. In this section, we will formulate a generalization of Reuman's conjecture, and prove part of it, as a consequence of the results obtained above. To formulate the conjecture, we consider the following maps from \widetilde{W} to W . The map η_1 is just the projection from $\widetilde{W} = W \times X_*(A)$ to W . It is a group homomorphism. To describe the second map, we identify W with the set of Weyl chambers. The map $\eta_2: \widetilde{W} \rightarrow W$ keeps track of the finite Weyl chamber whose closure contains the alcove $x\mathbf{a}$. We define $\eta_2(x) = w$, where w is

the unique element in W such that $w^{-1}x\mathbf{a}$ is contained in the dominant chamber (so that the identity element of \widetilde{W} maps to the identity element of W).

We say that $x \in \widetilde{W}$ lies in the shrunken Weyl chambers, if $k(\alpha, x\mathbf{a}) \neq k(\alpha, \mathbf{a})$ for all roots α , or equivalently, if $U_\alpha \cap xI \neq U_\alpha \cap I$ for all α . For T a subset of the set S of simple reflections in W , let $W_T \subset W$ denote the subgroup generated by T . Let $\ell(w)$ denote the length of an element $w \in \widetilde{W}$. Finally, recall that we define the defect $\text{def}_G(b)$ of an element $b \in G(L)$ to be the F -rank of G minus the F -rank of J_b (cf. [GHKR]).

Conjecture 9.4.1. *a) Let $[b]$ be a basic σ -conjugacy class. Suppose $x \in \widetilde{W}$ lies in the shrunken Weyl chambers. Then $X_x(b) \neq \emptyset$ if and only if*

$$\eta_2(x)^{-1}\eta_1(x)\eta_2(x) \in W \setminus \bigcup_{T \subsetneq S} W_T,$$

and in this case

$$\dim X_x(b) = \frac{1}{2} (\ell(x) + \ell(\eta_2(x)^{-1}\eta_1(x)\eta_2(x)) - \text{def}_G(b)).$$

b) Let $[b]$ be an arbitrary σ -conjugacy class, and let $[b_b]$ be the unique basic σ -conjugacy class with $\eta_G(b) = \eta_G(b_b)$. Then there exists $N_b \in \mathbb{Z}_{\geq 0}$, such that for all $x \in \widetilde{W}$ of length $\ell(x) \geq N_b$, we have

$$X_x(b) \neq \emptyset \iff X_x(b_b) \neq \emptyset,$$

and in this case

$$\dim X_x(b) = \dim X_x(b_b) - \frac{1}{2} (\langle 2\rho, \nu \rangle + \text{def}_G(b) - \text{def}_G(b_b)),$$

where ν denotes the Newton point of b .

Part (b) of this conjecture generalizes Conjecture 7.5.1 of [GHKR]. It fits well with Beazley's Conjecture 1.0.1 and the qualitative picture of $B(G)_x$ that is suggested by her results on $SL(3)$ (see [Be]). The term $\langle 2\rho, \nu \rangle$ appearing here can also be interpreted (see section 13) as the length of a suitable semistandard representative of $[b]$ in \widetilde{W} .

Using the algorithms discussed in [GHKR] and in this article, we obtained ample numerical evidence for this conjecture. We made computations for root systems of type $A_2, A_3, A_4, C_2, C_3, G_2$, and for a number of choices of b , including cases where b is split, basic, or neither of the two, and both cases where $\eta_G(b) = 0$ and $\neq 0$.

The following remark shows that this conjecture is compatible with what we already know about affine Deligne-Lusztig varieties in the affine Grassmannian (cf. [GHKR],[V2]).

Remark 9.4.2. *Conjecture 9.4.1 implies Rapoport's dimension formula for affine Deligne-Lusztig varieties $X_\mu(b)$ in the affine Grassmannian for b basic (and $\mu \in X_*(A)$ dominant). Indeed, if $w_0 \in W$ is the longest element, then we have*

$$\dim X_\mu(b) + \ell(w_0) = \sup\{\dim X_x(b); x \in W\epsilon^\mu W\}.$$

Now for the longest element $x \in W\epsilon^\mu W$, we have $\eta_1(x) = \eta_2(x) = w_0$, so

$$\eta_2(x)^{-1}\eta_1(x)\eta_2(x) = w_0 \in W \setminus \bigcup_{T \subsetneq S} W_T,$$

and by the dimension formula given in the conjecture, the supremum above is equal to

$$\frac{1}{2} (\sup\{\ell(x); x \in W\epsilon^\mu W\} + \ell(w_0) - \text{def}_G(b)).$$

Let X^μ denote the $G(\mathfrak{o})$ -orbit of $\epsilon^\mu G(\mathfrak{o})$ in the affine Grassmannian. Since

$$\sup\{\ell(x); x \in W\epsilon^\mu W\} = \dim X^\mu + \ell(w_0) = \langle 2\rho, \mu \rangle + \ell(w_0),$$

altogether we obtain

$$\dim X_\mu(b) = \langle \rho, \mu \rangle - \frac{1}{2} \text{def}_G(b),$$

which is the desired result.

Let us relate this conjecture to the results of the previous subsection. The relation relies on the following lemma (which also follows easily from Proposition 3.0.5).

Lemma 9.4.3. *Let $x \in \widetilde{W}$, and write $w = \eta_2(x) \in W$.*

- a) *If $P = MN \supset {}^w B$ is a parabolic subgroup with $x \in \widetilde{W}_M$, then $x\mathbf{a}$ is a P -alcove.*
- b) *If x is an element of the shrunken Weyl chambers which is a P -alcove for a semistandard parabolic subgroup P , then $P \supset {}^w B$.*

Proof. First note that by assumption $w^{-1}x\mathbf{a}$ lies in the dominant chamber. This means precisely that $w^{-1}xI \cap U \subseteq I \cap U$ (where U denotes the unipotent radical of our Borel B), so we obtain

$${}^x I \cap N \subseteq {}^x I \cap {}^w U \subseteq {}^w (I \cap U) \subseteq I.$$

This inclusion is what we needed to show for part a).

Now let us prove b). Assume $x\mathbf{a}$ is a P -alcove and write $P = MN$ for the Levi decomposition of P . We need to show that $N \subseteq {}^w U$. Let $\alpha \in R_N$. Then we have

$${}^x I \cap U_\alpha \subsetneq I \cap U_\alpha.$$

(We get \subsetneq rather than just \subseteq because x is in the shrunken Weyl chambers.) This implies however that

$${}^x I \cap U_{-\alpha} \supsetneq I \cap U_{-\alpha}.$$

On the other hand, by what we have seen above,

$${}^x I \cap {}^w U \subseteq {}^w I \cap {}^w U \subseteq {}^w U(\epsilon\mathfrak{o}).$$

This shows that $U_{-\alpha} \not\subseteq {}^w U$, hence $U_\alpha \subseteq {}^w U$, as we wanted to show. \square

From this lemma, we obtain the following strengthening of the “only if” direction of part a) of Conjecture 9.4.1 above.

Proposition 9.4.4. *Let b be basic. Let $x \in \widetilde{W}$, and write $x = \epsilon^\lambda v$, $v \in W$. Assume that $\lambda \neq \overline{\nu}_b$ and that $\eta_2(x)^{-1}\eta_1(x)\eta_2(x) \in \bigcup_{T \subsetneq S} W_T$. Then $X_x(b) = \emptyset$.*

Proof. Write $w := \eta_2(x) \in W$. By the lemma and our hypothesis, $x\mathbf{a}$ is a P -alcove for a parabolic subgroup $P = MN \supset {}^w B$ of G . The only thing we need to check in order to apply Corollary 9.3.1 is that $\eta_{M'}(w^{-1}x) \neq \eta_{M'}(\overline{\nu}_b)$, where $M' = w^{-1}M$. (Recall that the precise meaning of $\eta_{M'}(\overline{\nu}_b)$ is described after Cor. 9.3.1.) But if we had equality here, then $w^{-1}\lambda - \overline{\nu}_b$ would be a linear combination of coroots of M' . On the other hand, $w^{-1}\lambda$ is dominant, and since M' is the Levi component of a proper standard parabolic subgroup, we obtain $\lambda = \overline{\nu}_b$, which is excluded by assumption. \square

Why does this imply the “only if” direction of part a) of Conjecture 9.4.1? Write $x = \epsilon^\lambda v$. We claim that if $x\mathbf{a}$ belongs to the shrunken Weyl chambers and $\eta_2(x)^{-1}\eta_1(x)\eta_2(x)$ belongs to a proper parabolic subgroup of W , then $\lambda \neq \bar{\nu}_b$. Suppose instead that $\lambda = \bar{\nu}_b$. Then ϵ^λ belongs to the center of G and $x\mathbf{a} = v\mathbf{a}$. This alcove belongs to the shrunken Weyl chambers only if $\eta_1(x) = v = w_0$. But in that case $\eta_2(x)^{-1}\eta_1(x)\eta_2(x)$ cannot belong to a proper parabolic subgroup of W .

We conclude this subsection by showing that our Conjecture 9.3.2 implies the validity of the “if” direction of part a) of Conjecture 9.4.1.

Proposition 9.4.5. *Assume that Conjecture 9.3.2 holds. Let $x \in \widetilde{W}$ be an element of the shrunken Weyl chambers with $\eta_G(x) = \eta_G(b)$ and*

$$\eta_2(x)^{-1}\eta_1(x)\eta_2(x) \in W \setminus \bigcup_{T \subsetneq S} W_T.$$

Then $X_x(b) \neq \emptyset$.

Proof. It is enough to show that $x\mathbf{a}$ is not a P -alcove for any *proper* parabolic subgroup $P = MN \subset G$. By the lemma above, if it were we would have $P \supset \eta_2(x)B$. But the assumption says precisely that x does not lie in \widetilde{W}_M for such P . \square

10. DIMENSION THEORY FOR THE GROUPS $I_M N$

In this section we lay some conceptual foundations for studying the dimensions of affine Deligne-Lusztig varieties $X_x(b)$, where $[b] \in B(G)$ is an arbitrary σ -conjugacy class. In the case where $b = \epsilon^\nu$ for some $\nu \in X_*(A)$, such a study was carried out in [GHKR], section 6. The result was a finite algorithm to compute dimensions (a special case of our Theorem 11.2.1 below). Underlying this study was the notion of (*ind-*)*admissible* subset of $U(L)$. In this paragraph, we introduce a suitable framework that works for general elements b .

Let J be an Iwahori subgroup which is the fixer of an alcove in the standard apartment, and let $P = MN \supset A$ be any parabolic subgroup of G . Let $J_P = J_M N$ (where $J_M := J \cap M$). We want to establish a “dimension theory” for ind-admissible subsets of J_P , similar to the theory in [GHKR].

Fix any semistandard Borel subgroup contained in P and use it to define the sets of simple roots Δ_M and Δ_N . We fix a coweight λ_0 with $\langle \alpha, \lambda_0 \rangle = 0$ for $\alpha \in \Delta_M$, and $\langle \alpha, \lambda_0 \rangle > 0$ for $\alpha \in \Delta_N$, and consider the subgroups

$$N(m) := \epsilon^{m\lambda_0}(N \cap J)\epsilon^{-m\lambda_0}, \quad m \in \mathbb{Z},$$

cf. loc. cit. 5.2; our choice of λ_0 is a little different, but this clearly does not affect the validity of the dimension theory for N as in loc. cit. Furthermore, we choose a separated descending filtration $(J_M(m))_{m \in \mathbb{Z}}$ of J_M by normal subgroups, such that $J_M(m) = J_M$ for $m \leq 0$, and such that all the quotients $J_M(m)/J_M(m')$ are finite-dimensional over \bar{k} . (For example, we could use a Moy-Prasad filtration.) Finally, we set $J_P(m) := J_M(m)N(m)$, and we obtain a separated and exhaustive filtration

$$J_P \supset \cdots J_P(-1) \supset J_P(0) \supset J_P(1) \supset J_P(2) \supset \cdots.$$

The quotients $J_P(m)/J_P(m')$, $m \leq m'$ are finite-dimensional varieties over k in a natural way (more precisely, they coincide, in a natural way, with the set of \bar{k} -valued points of a k -variety). Since J_M normalizes each $N(m)$, $J_P(m)/J_P(m')$ is a fiber bundle over $J_M(m)/J_M(m')$ with fibers $N(m)/N(m')$. We say that a subset

$Y \subseteq J_P$ is *admissible*, if there are $m \leq m'$ such that it is contained in $J_P(m)$ and is the full inverse image under the projection $J_P(m) \rightarrow J_P(m)/J_P(m')$ of a locally closed subset of $J_P(m)/J_P(m')$. We say that $Y \subseteq J_P$ is *ind-admissible*, if for all m , $Y \cap J_P(m)$ is an admissible subset of J_P . Obviously, admissible subsets are in particular ind-admissible.

As in [GHKR], for an admissible subset $Y \subset J_P(m)$, we can define a notion of dimension

$$\dim Y := \dim(Y/J_P(m)) - \dim(J_P(0)/J_P(m));$$

note this is always an element of \mathbb{Z} , unless Y is empty. For an ind-admissible subset $Y \subset J_P$, we define

$$\dim Y := \sup\{\dim(Y \cap J_P(-m)) : m \geq 0\}.$$

We may sometimes have $\dim Y = +\infty$ (for example for $Y = J_P$). Of course in making these definitions we made a choice, namely we normalized things so that $\dim(J_P(0)) = 0$. But as before differences

$$\dim Y_1 - \dim Y_2$$

for admissible subsets Y_1, Y_2 are independent of any such choice.

11. THE GENERALIZED SUPERSET METHOD

11.1. The retractions ρ_P . Fix a standard parabolic $P = MN$. Write $I_P = I_M N = (I \cap M(L))N(L)$.

Lemma 11.1.1. *Let $w \in \widetilde{W}$, and $J_P = w^{-1}I_P$. The projection $N_G A(L) \rightarrow J_P \backslash G(L)/I$ induces a bijection*

$$\widetilde{W} \cong J_P \backslash G(L)/I.$$

Proof. Because we can conjugate the situation by w^{-1} , we may as well assume that $w = 1$. Since the set $P \backslash G(L)/K$ has only one element, we can identify the double quotient $P \backslash G(L)/I$ with $W_M \backslash W \cong \widetilde{W}_M \backslash \widetilde{W}$. We obtain a commutative diagram

$$\begin{array}{ccc} \widetilde{W} & \longrightarrow & I_P \backslash G(L)/I \\ \downarrow q & & \downarrow p \\ \widetilde{W}_M \backslash \widetilde{W} & \xrightarrow{\cong} & P \backslash G(L)/I. \end{array}$$

Now for $v \in \widetilde{W}$, we have

$$q^{-1}(\widetilde{W}_M v) = \widetilde{W}_M v \cong I_M \backslash M / ({}^v I)_M \cong I_P \backslash P / ({}^v I \cap P) \cong p^{-1}(PvI).$$

This proves the lemma. \square

Denote by ${}^M W$ the set of minimal length representatives in W of the cosets in $W_M \backslash W$.

Lemma 11.1.2. *Let $\lambda \in X_*(A)$ be such that $\langle \alpha, \lambda \rangle = 0$ for all roots α in M , and let $v \in {}^M W$.*

- (1) *All elements of I_M fix the alcove $\epsilon^\lambda v \mathbf{a}$.*
- (2) *If $n \in N$, and if λ satisfies $\epsilon^{-\lambda} n \epsilon^\lambda \in {}^v I \cap N$ (which is true whenever λ is sufficiently antidominant with respect to the roots in $\text{Lie } N$), then n fixes the alcove $\epsilon^\lambda v \mathbf{a}$.*

Proof. To prove (1), we first note that $({}^v I)_M = I_M$, because v is the minimal length representative in its W_M -coset. This shows that

$$I_M = \epsilon^\lambda v(I \cap v^{-1} M) \subseteq \epsilon^\lambda v I.$$

Similarly, under the assumption on n made in (2), we obtain that $n \in \epsilon^\lambda v I$. \square

Denote by \mathcal{A} the standard apartment of G with respect to our fixed torus A . Let ρ_P be the retraction from the Bruhat-Tits building of $G(L)$ to \mathcal{A} , defined as follows. For each alcove \mathbf{b} in the building, all retractions of \mathbf{b} with respect to an alcove of the form $\epsilon^\lambda v \mathbf{a}$, λ, v as in part (2) of the lemma, have the same image, say \mathbf{c} . Here we must stipulate that λ is sufficiently anti-dominant (depending on \mathbf{b}) with respect to the roots in $\text{Lie } N$. We set

$$\rho_P(\mathbf{b}) = \mathbf{c}.$$

(In fact, we get the same retraction if we retract with respect to any alcove which lies between the root hyperplanes H_α and $H_{\alpha,1}$ for all roots α of M , and is sufficiently antidominant for all roots of G lying in N . Compare also Rousseau's notion of *cheminée*, [Ro] §9.)

Lemma 11.1.3. *For $g \in I_P$, $\rho_P|_{g\mathcal{A}} = g^{-1}$.*

Proof. Clearly, g^{-1} maps $g\mathcal{A}$ to \mathcal{A} , and g^{-1} fixes the alcoves $t_\lambda v \mathbf{a}$ for λ sufficiently anti-dominant. This implies the lemma. \square

Proposition 11.1.4. *Let $y \in \widetilde{W}$.*

(1) *We have*

$$I_P y I / I = \rho_P^{-1}(y \mathbf{a}).$$

In other words: we can identify ρ_P (as a map from the set of alcoves in the building to the set of alcoves in the standard apartment) with the map $G(L)/I \rightarrow I_P \backslash G(L)/I \cong \widetilde{W}$ obtained from Lemma 11.1.1.

(2) *More generally, let $w \in \widetilde{W}$, and let $J_P = w^{-1} I_P$. Consider the map*

$$\rho_{P,w}: G(L)/I \rightarrow \widetilde{W}, \quad g \mapsto w^{-1} \rho_P(wg).$$

Then

$$J_P y I = \rho_{P,w}^{-1}(y \mathbf{a}).$$

Proof. Part (1) follows from the previous lemma, cf. [BT], Remarque 7.4.22 which deals with the case $P = G$. To prove part (2), combine part (1) with the following commutative diagram:

$$\begin{array}{ccccc} & & \rho_P & & \\ & & \curvearrowright & & \\ G(L) & \xrightarrow{\text{proj}} & I_P \backslash G(L)/I & \xrightarrow{\cong} & \widetilde{W} \\ \downarrow w^{-1} \cdot & & \downarrow w^{-1} \cdot & & \downarrow w^{-1} \cdot \\ G(L) & \xrightarrow{\text{proj}} & J_P \backslash G(L)/I & \xrightarrow{\cong} & \widetilde{W} \end{array}$$

\square

In the extreme cases, we get the following: If $P = G$, then ρ_G is just the usual retraction $\rho_{\mathbf{a}}$ with respect to the base alcove. If $P = B$, then we get as ρ_B the retraction with respect to “a point at infinity in the B -antidominant chamber”. Note that the maps $\rho_{P,w}$ are retractions to the standard apartment just like the ρ_P , but for a different choice of base alcove.

11.2. An algorithm for computing $\dim X_x(b)$. In this subsection, we give a formula for the dimensions

$$\dim X_x(b) \cap I_P w \mathbf{a},$$

for any $w \in \widetilde{W}$. This gives us the dimension of $X_x(b)$, because we have

$$(11.2.1) \quad \dim X_x(b) = \sup_{w \in \widetilde{W}} \dim(X_x(b) \cap I_P w \mathbf{a}).$$

To show this, observe that

$$\dim X_x(b) = \sup_{v \in \widetilde{W}} \dim(X_x(b) \cap \overline{Iv \mathbf{a}}),$$

where $\overline{\cdot}$ indicates the closure. Now every $\overline{Iv \mathbf{a}}$ is contained in a finite union of I_P -orbits, in fact

$$\overline{Iv \mathbf{a}} \subseteq \bigcup_{w \in S_v} I_P w \mathbf{a}$$

where $S_v := \{w \in \widetilde{W} : w \leq v\}$. Thus

$$\dim(X_x(b) \cap \overline{Iv \mathbf{a}}) = \sup_{w \in S_v} \dim(X_x(b) \cap \overline{Iv \mathbf{a}} \cap I_P w \mathbf{a}) \leq \sup_{w \in \widetilde{W}} \dim(X_x(b) \cap I_P w \mathbf{a})$$

which shows that in (11.2.1), \leq holds. Since the inequality \geq is obviously true, the desired equality follows. Also note that we know a priori that $\dim X_x(b)$ is finite, for example by using the finite-dimensionality of affine Deligne-Lusztig varieties in the affine Grassmannian, established in [GHKR] and [V1].

Our result in Theorem 11.2.1 is not a “closed formula”, even for fixed w , because it involves the dimensions of intersections of I - and $w^{-1}I_P$ -orbits. However, these dimensions can be computed (at least by a computer) for fixed w . (Here we make use of the interpretation of I_P -orbits in terms of “foldings”, see Proposition 11.1.4.)

Throughout this subsection, we fix a σ -conjugacy class, say $[b] \in B(G)_P \subset B(G)$, letting M denote the Levi component of a standard parabolic $P = MN$. Denote by $b \in \widetilde{W}_M$ the standard representative of $[b]$ (see Definition 7.2.3). Write $I_P = I_M N$. We have $b I_P b^{-1} = I_P$. Denote by $\nu \in X_*(A)_{\mathbb{Q}}$ the Newton vector for b (where b is considered as an element of $M(L)$). Since b is M -basic, ν is “central in M ” (and in particular M -dominant). Let ν_{dom} denote the unique G -dominant element in the W -orbit of ν .

For any $y \in \widetilde{W}$, we write $\mathbf{a}_y := y \mathbf{a}$. Let $\rho \in X^*(A)_{\mathbb{Q}}$ denote the half-sum of the positive roots of A in G .

Theorem 11.2.1. *Let $w \in \widetilde{W}$. Then writing $\tilde{b} = w^{-1} b w$, and denoting by ν the Newton vector of b , we have*

$$\dim(X_x(b) \cap I_P w \mathbf{a}) = \dim(I \mathbf{a}_x \cap w^{-1} I_P \mathbf{a}_{\tilde{b}}) - \langle \rho, \nu + \nu_{\text{dom}} \rangle.$$

Proof. Fix a representative of w in $N_G A(L)$ fixed by σ , and again denote it by w . Then multiplication by w^{-1} defines a bijection

$$X_x(b) \cap I_P w \mathbf{a} \cong X_x(w^{-1} b w) \cap w^{-1} I_P \mathbf{a},$$

which preserves the dimensions. Note that $w^{-1}I_P := w^{-1}(I_P)$ here.

We write $\tilde{b} = w^{-1}bw$, and consider the map

$$\begin{aligned} f_{\tilde{b}}: w^{-1}I_P &\longrightarrow w^{-1}I_P, \\ g &\longmapsto g^{-1}\tilde{b}\sigma(g)\tilde{b}^{-1}. \end{aligned}$$

Let

$$\widetilde{X_x(\tilde{b})} = \{g \in G(L); g^{-1}\tilde{b}\sigma g \in IxI\}.$$

Then $\widetilde{X_x(\tilde{b})} \cap w^{-1}I_P = f_{\tilde{b}}^{-1}(IxI\tilde{b}^{-1} \cap w^{-1}I_P)$, so

$$X_x(\tilde{b}) \cap w^{-1}I_P \mathbf{a} = f_{\tilde{b}}^{-1}(IxI\tilde{b}^{-1} \cap w^{-1}I_P) / (I \cap w^{-1}I_P).$$

Lemma 11.2.2. *We have the equality*

$$\dim f_{\tilde{b}}^{-1}(IxI\tilde{b}^{-1} \cap w^{-1}I_P) - \dim(IxI\tilde{b}^{-1} \cap w^{-1}I_P) = \langle \rho, \nu - \nu_{\text{dom}} \rangle.$$

Proof of Lemma. To ease the notation, let us write $J_P := w^{-1}(I_P) = (w^{-1}I)_{w^{-1}P}$, and $J_M := (w^{-1}I)_{wM}$. It is easy to see that $IxI\tilde{b}^{-1} \cap J_P$ is an admissible subset of J_P . It will follow from our proof below that its preimage under $f_{\tilde{b}}$ is ind-admissible, so that we can define the dimensions of these subsets using the theory from section 10. The left hand side of the equality is therefore well-defined. We can even make a very convenient choice of filtration on J_M , one which is stable under $\text{Ad}(\tilde{b})$: take the Moy-Prasad filtration $J_M(\bullet)$ on J_M associated to the barycenter of the alcove in the reduced building of $M(L)$ which corresponds to J_M .

A straightforward calculation shows that we can write the map $f_{\tilde{b}}$ as follows (here $i \in J_M$, $n \in w^{-1}N$):

$$g = in \mapsto g^{-1}\tilde{b}\sigma(g)\tilde{b}^{-1} = i^{-1}\tilde{b}\sigma(i) \cdot \tilde{i}n^{-1}\tilde{b}\sigma(n),$$

with $\tilde{i} := \tilde{b}\sigma(i)^{-1}i$.

The projection $J_P \rightarrow J_M$ is an ‘‘ind-admissible fiber bundle’’, in a sense which the reader will have no trouble making precise (see section 10). The above description of $f_{\tilde{b}}$ indicates how it behaves on the base and on the fibers. Let us analyze the relative dimension of $f_{\tilde{b}}$ by studying the base and the fibers in turn.

First, we consider the base J_M . Since \tilde{b} normalizes J_M , the map $J_M \rightarrow J_M$, $i \mapsto i^{-1}\tilde{b}\sigma(i)$ is surjective, and has relative dimension zero. The proof is an adaptation of the proof of Lang’s theorem. Indeed, J_M has a filtration by normal subgroups (the $J_M(m)$ for $m \geq 0$ in the Moy-Prasad filtration described above) which are stabilized by $\text{Ad}(\tilde{b})$, such that on the finite-dimensional quotients our map $J_M \rightarrow J_M$ induces a Lang map, which is finite étale and surjective.

Second, we study the relative dimension of $f_{\tilde{b}}$ ‘‘on the fibers’’ of $J_P \rightarrow J_M$. That is, we fix $\tilde{i} \in J_M$ as above, and study the fibers of the map $w^{-1}N(L) \rightarrow w^{-1}N(L)$ given by $n \mapsto \tilde{i}n^{-1}\tilde{b}\sigma(n)$. Fortunately, most of the necessary work was already done in [GHKR], Prop. 5.3.2. In fact, that proposition implies that the fiber dimension is (using the notation of loc. cit.)

$$d(\tilde{i}, \tilde{b}) := d(\mathfrak{n}(L), \text{Ad}_{\mathfrak{n}}(\tilde{i})^{-1} \text{Ad}_{\mathfrak{n}}(\tilde{b})\sigma) + \text{val det Ad}_{\mathfrak{n}}(\tilde{i}).$$

Here \mathfrak{n} denotes the Lie algebra of $w^{-1}N$. Since $\tilde{i} \in J_M$, the second summand vanishes. Moreover, $\text{Ad}_{\mathfrak{n}}(\tilde{i})^{-1} \text{Ad}_{\mathfrak{n}}(\tilde{b}) = \text{Ad}_{\mathfrak{n}}(i^{-1}\tilde{b}\sigma(i))$. Since σ -conjugation induces

an isomorphism of F -spaces, we obtain

$$d(\tilde{i}, \tilde{b}) = d(1, \tilde{b}) = \langle \rho, \nu - \nu_{\text{dom}} \rangle,$$

cf. loc. cit. Prop. 5.3.1.

It is clear that we should be able to put these two pieces of information together (and obtain the stated result that the relative dimension of $f_{\tilde{b}}$ is $\langle \rho, \nu - \nu_{\text{dom}} \rangle$) by looking at the corresponding finite-dimensional situation. However, to make this vague idea convincing it seems easiest to follow the argument of loc. cit. Prop. 5.3.1. First, we correct for the inconvenient fact that $f_{\tilde{b}}$ need not preserve $J_P(0)$. Let $P' := w^{-1}P$, $M' := w^{-1}M$, $N' := w^{-1}N$, and $I' := w^{-1}I$. For any $m_1, m_2 \in M'(L)$ which normalize $J_P = I'_{P'}$, define

$$\begin{aligned} f_{m_1, m_2} : J_P &\longrightarrow J_P, \\ g &\longmapsto m_1 g^{-1} m_1^{-1} \cdot m_2 \sigma(g) m_2^{-1}. \end{aligned}$$

Note that $f_{\tilde{b}} = f_{1, \tilde{b}}$. Fix $\lambda_0 \in X_*(Z(M'))$ such that $\langle \alpha, \lambda_0 \rangle > 0$ for all $\alpha \in R_{N'}$. Then we may replace $f_{\tilde{b}} = f_{1, \tilde{b}}$ with $f := f_{\epsilon^{t\lambda_0}, \epsilon^{t\lambda_0} \tilde{b}}$ for a suitably large integer t , chosen such that f preserves $J_P(0) = I'_{M'} \cdot N' \cap I'$. Note that f then automatically preserves $J_P(m)$ for each integer $m \geq 0$ (we shall not need this fact). Denote by $f_0 : J_P(0) \rightarrow J_P(0)$ the restriction of f to $J_P(0)$. As in loc. cit., our goal is now to prove the following

Claim: Let $m_1 = \epsilon^{t\lambda_0}$ and $m_2 = \epsilon^{t\lambda_0} \tilde{b}$ and set $f := f_{m_1, m_2}$. If $Y \subset J_P$ is admissible, then $f^{-1}Y$ is ind-admissible and

$$\dim f^{-1}Y - \dim Y = d(m_1, m_2).$$

Continuing to follow the strategy of the proof of Prop. 5.3.2 of loc. cit., we can use *the proof of* loc. cit. Claim 1 to find an $a := \epsilon^{t_1 \lambda_0}$ for a large integer t_1 such that

$$c_a J_P(0) \subseteq f J_P(0),$$

where c_a denotes the conjugation map $g \mapsto aga^{-1}$ for $g \in J_P$. Fix this element a once and for all. Next we prove the following

Subclaim: Suppose that Y is an admissible subset of $c_a J_P(0)$. Then $f_0^{-1}(Y)$ is admissible, and

$$\dim f_0^{-1}Y - \dim Y = d(m_1, m_2).$$

Proof of Subclaim: At this point we have to replace the filtration $\{J_P(m)\}_{m \geq 0}$ of $J_P(0)$ with one which is better behaved with respect to the morphism f_0 . So, for $m \geq 0$ let $I'_m \subset I'$ denote the m -th principal congruence subgroup of the Iwahori subgroup I' ; by convention $I'_0 = I'$. Let $J_{M, m} := I'_m \cap M'$ and $N'_m := I'_m \cap N'$. Let $J_{P, m} = J_{M, m} N'_m = I'_m \cap P'$. It is clear that J_M normalizes each N'_m , so that we have a fiber bundle for each $0 \leq m_1 \leq m_2$

$$\pi : J_{P, m_1} / J_{P, m_2} \rightarrow J_{M, m_1} / J_{M, m_2}$$

with fiber N_{m_1} / N_{m_2} . Also, using our specific choices of m_1, m_2 above, it is clear that f_0 preserves $J_{P, m}$ and in fact f_0 induces a well-defined map on the quotients

$$\bar{f} : J_{P, 0} / J_{P, m} \rightarrow J_{P, 0} / J_{P, m}$$

for any $m \geq 0$. Here, we used that m_1 and m_2 and $J_{P, 0}$ each normalize $J_{P, m}$, for all $m \geq 0$. (See (6.1.1).)

Now choose a large positive integer m such that Y comes from a locally closed subset \bar{Y} of $J_{P,0}/J_{P,m}$. Consider the following commutative diagram

$$\begin{array}{ccc} J_{P,0} & \xrightarrow{f_0} & J_{P,0} \\ p \downarrow & & p \downarrow \\ J_{P,0}/J_{P,m} & \xrightarrow{\bar{f}} & J_{P,0}/J_{P,m} \\ \pi \downarrow & & \pi \downarrow \\ J_{M,0}/J_{M,m} & \xrightarrow{\bar{f}_M} & J_{M,0}/J_{M,m}, \end{array}$$

where p is the canonical projection, π is the fiber bundle described above, and \bar{f} and \bar{f}_M are the morphisms induced by f_0 . Note that $f_0^{-1}Y = p^{-1}\bar{f}^{-1}\bar{Y}$, showing that $f_0^{-1}Y$ is admissible. Note also that since $Y \subseteq c_a J_P(0) \subseteq f J_P(0)$, the subset \bar{Y} is contained in the image of \bar{f} , and our dimension formula is a consequence of the identity

$$\dim \bar{f}^{-1}\bar{Y} - \dim \bar{Y} = d(m_1, m_2).$$

But the latter equality now follows easily from our earlier considerations of the base and fiber of the fiber bundle π : the map \bar{f}_M is surjective of relative dimension zero, and the relative dimension of \bar{f} on locally closed subsets of the fibers of π over $\pi(\bar{Y})$ is given by $d(m_1, m_2)$; see the proof of loc. cit. Claim 3. This proves our subclaim.

As in loc. cit., our claim follows from the subclaim. Write $d(m_1, m_2) =: d$. If $Y \subset J_P$ is any admissible subset, then we have proved that $f^{-1}Y \cap a_1^{-1}J_P(0)a_1$ is admissible of dimension $\dim Y + d$ for any $a_1 \in Z(M')(F)$ such that $a_1 Y a_1^{-1} \subseteq a J_P(0) a^{-1}$. Let t_0 be sufficiently large so that $a_t := \epsilon^{t\lambda_0}$ satisfies $a_t Y a_t^{-1} \subseteq a J_P(0) a^{-1}$ for all $t \geq t_0$. For all such t we have proved that $f^{-1}Y \cap a_t^{-1}J_P(0)a_t$ is admissible of dimension $\dim Y + d$. This is enough to prove the claim, hence also the lemma. \square

Remark 11.2.3. *The proof of Lemma 11.2.2 shows that $f_{\tilde{b}} : J_P \rightarrow J_P$ is surjective.*

Now let

$$d(x, \tilde{b}, {}^{w^{-1}}I_P) := \dim(I\mathbf{a}_x \cap {}^{w^{-1}}I_P \mathbf{a}_{\tilde{b}}).$$

We have a dimension-preserving bijection

$$I\mathbf{a}_x \cap {}^{w^{-1}}I_P \mathbf{a}_{\tilde{b}} \cong (IxI\tilde{b}^{-1} \cap {}^{w^{-1}}I_P) / ({}^{w^{-1}}I_P \cap \tilde{b}I)$$

given by right multiplication by \tilde{b}^{-1} , so that

$$d(x, \tilde{b}, {}^{w^{-1}}I_P) = \dim IxI\tilde{b}^{-1} \cap {}^{w^{-1}}I_P - \dim {}^{w^{-1}}I_P \cap \tilde{b}I.$$

Let $\rho_N \in X^*(A)_{\mathbb{Q}}$ denote the half-sum of the roots in R_N .

Lemma 11.2.4. *Consider $c_{\tilde{b}} : {}^{w^{-1}}I_P \rightarrow {}^{w^{-1}}I_P$, $g \mapsto \tilde{b}g\tilde{b}^{-1}$. Then*

$${}^{w^{-1}}I_P \cap \tilde{b}I = c_{\tilde{b}}({}^{w^{-1}}I_P \cap I),$$

hence

$$\dim({}^{w^{-1}}I_P \cap I) - \dim({}^{w^{-1}}I_P \cap \tilde{b}I) = \langle 2\rho_N, \nu \rangle.$$

Proof. As the previous lemma, this can be proved by looking at the projection $J_P \rightarrow J_M$ and then separately computing the contribution from the base J_M (which is 0) and that from the fibers (which is $\langle 2\rho_N, \nu \rangle$, see [GHKR]). \square

Altogether we have now

$$\begin{aligned}
& \dim X_x(b) \cap I_P \mathbf{a}_w \\
&= \dim f_{\tilde{b}}^{-1}(IxI\tilde{b}^{-1} \cap w^{-1}I_P) - \dim I \cap w^{-1}I_P \\
&= \dim IxI\tilde{b}^{-1} \cap w^{-1}I_P - \dim I \cap w^{-1}I_P + \langle \rho, \nu - \nu_{\text{dom}} \rangle \\
&= d(x, \tilde{b}, w^{-1}I_P) + \dim w^{-1}I_P \cap \tilde{b}I - \dim I \cap w^{-1}I_P + \langle \rho, \nu - \nu_{\text{dom}} \rangle \\
&= d(x, \tilde{b}, w^{-1}I_P) + \langle \rho, \nu - \nu_{\text{dom}} \rangle - \langle 2\rho_N, \nu \rangle \\
&= d(x, \tilde{b}, w^{-1}I_P) - \langle \rho, \nu + \nu_{\text{dom}} \rangle,
\end{aligned}$$

where in the final step we have used the equality $\langle \rho, \nu \rangle = \langle \rho_N, \nu \rangle$. This is what we wanted to show. \square

Together with the description (Proposition 11.1.4) of $w^{-1}I_P$ -orbits in $G(L)/I$ as fibers of a certain retraction of the building, Theorem 11.2.1 gives us an algorithm to compute whether for a given w the intersection $X_x(b) \cap I_P w \mathbf{a}$ is empty or non-empty. If this information were available for all w , we could conclude whether $X_x(b)$ is non-empty (and compute its dimension from the dimensions of all these intersections). As noted above, it is clear that all affine Deligne-Lusztig varieties are finite-dimensional, so that the supremum of $\dim(X_x(b) \cap I_P w \mathbf{a})$ is attained for some w . It does not seem easy to give a bound for the length of w depending on x and b .

As examples, let us consider the extreme cases:

- (1) $P = B$. Then $I_P = A(\mathfrak{o})U$, and $b = \epsilon^\nu \in B(G)_B$ where $\nu \in X_*(A)$ is a regular dominant translation element. This case was considered in [GHKR]. The above formula is the same as in loc. cit., equations (6.3.3), (6.3.4).
- (2) $P = G$. Then $I_P = I$, and $b \in \Omega_G$ is a basic σ -conjugacy class. In this case, the dimension formula reads

$$\dim X_x(b) \cap I w \mathbf{a} = \dim I \mathbf{a}_x \cap w^{-1} I \mathbf{a}_{w^{-1}bw}$$

(since ν is central in G). This case is the case analyzed by Reuman in [Re2] for the case $b = 1$, and low-rank groups. So let $b = 1$ (the case of other basic b 's is analogous). We have that

$$\begin{aligned}
X_x(1) \neq \emptyset &\iff \exists w \in \widetilde{W} : IxI \cap w^{-1}I \neq \emptyset \\
&\iff \exists w \in \widetilde{W} : \rho_G^{-1}(x) \cap \rho_{G,w}^{-1}(1) \neq \emptyset.
\end{aligned}$$

There are two ways to reformulate this. The algorithmic description in the spirit of the above amounts to

$$X_x(1) \neq \emptyset \iff \exists w \in \widetilde{W} : 1 \in \rho_{G,w}(IxI)$$

On the other hand, we also obtain

$$X_x(1) \neq \emptyset \iff \exists w \in \widetilde{W} : x \in \rho_G(Iw^{-1}IwI).$$

which leads to the ‘‘folding method’’ used by Reuman, since $Iw^{-1}IwI/I$, as a set of alcoves in the building, is exactly the set of alcoves which can be reached by a gallery of type $i_r, \dots, i_1, i_1, \dots, i_r$ (for a fixed reduced expression $w = s_{i_1} \cdots s_{i_r}$). See also section 13.

Remark 11.2.5. *The dimension formula in example (2) can be interpreted in terms of structure constants for the affine Hecke algebra. Let H denote the affine Hecke algebra over $\mathbb{Z}[v, v^{-1}]$ corresponding to the extended affine Weyl group \widetilde{W} and let $T_x \in H$ denote the standard basis element corresponding to $x \in \widetilde{W}$. Define the parameter $q := v^2$, and consider the structure constants $C(x, y, z) \in \mathbb{Z}[q]$ for $x, y, z \in \widetilde{W}$ defined by the equality in H*

$$T_x T_y = \sum_z C(x, y, z) T_z.$$

Then it is straightforward to check that

$$\dim I\mathbf{a}_x \cap {}^{w^{-1}}I\mathbf{a}_{w^{-1}bw} = \deg_q C(x, w^{-1}b^{-1}, w^{-1}).$$

(By convention, we set $\deg_q 0 := -\infty = \dim \emptyset$.) Determining the structure constants $C(x, w^{-1}b^{-1}, w^{-1})$ is also a “folding algorithm”, so this does not give an essentially different way to compute dimensions of affine Deligne-Lusztig varieties. But it does give some insight on the inherent complexity of the algorithm.

12. ON REDUCTION TO THE BASIC CASE AND A FINITE ALGORITHM

One drawback of Theorem 11.2.1 is that it does not produce a *finite* algorithm to compute the non-emptiness or dimension of $X_x^G(b)$. In this section, we explain how we can at least find a finite algorithm which reduces the non-emptiness and dimension of $X_x^G(b)$ to that of a finite number of related varieties $X_y^{M'}(\tilde{b})$, where for all the latter \tilde{b} is basic in M' .

Using Theorem 11.2.1, we will usually have to check an infinite number of orbit intersections to determine whether a given $X_x(b)$ is empty or not. However, for b basic, we have proved the emptiness predicted by Conjecture 9.3.2 in Corollary 9.3.1. Why are we confident that Conjecture 9.3.2 also correctly predicts non-emptiness? In order to confirm the non-emptiness of $X_x(b)$ in a case it is expected, it is sufficient for the computer to detect a single non-empty intersection $I\mathbf{a}_x \cap {}^{w^{-1}}I\mathbf{a}_{w^{-1}bw}$ for some w , and in practice the computer also detects one (as far as we have checked). In other words, concerning the non-emptiness question for b basic, in practice the algorithm always terminates in finitely many steps, and in this way we are able to generate a complete emptiness/non-emptiness picture, at least when $\ell(x)$ is small enough for the computer to handle.

Let $P = MN$ denote a standard parabolic subgroup. Suppose $b \in \Omega_M \subset M(L)$ is the standard representative of a basic σ -conjugacy class in $M(L)$, and let $\nu = \bar{\nu}_b^M$ denote its Newton vector.

Recall that ${}^M W$ denotes the set of minimal length representatives of the cosets in $W_M \backslash W$. Note that $P \backslash G(L) / I \cong {}^M W$.

From now on, we fix an element $w \in {}^M W$. Write $M' = w^{-1}M$, $N' = w^{-1}N$, and $P' = w^{-1}P$. Let us denote $\tilde{b} := w^{-1}b \in \Omega_{M'}$. Note that $I_{M'} = w^{-1}(M \cap wI) = w^{-1}(M \cap I)$ is an Iwahori subgroup of M' . Let e_0 denote the base point of the affine flag variety $G(L) / I$ and let e'_0 denote the base point in $M'(L) / I_{M'}$.

We consider the map

$$\begin{aligned} \alpha_w : Pwe_0 &\rightarrow M'(L) / I_{M'} \\ mnwe_0 &\mapsto {}^{w^{-1}}me'_0, \end{aligned}$$

which is easily seen to be well-defined and surjective. Fix $m \in M(L)$ and write $m' := w^{-1}m \in M'(L)$. The map $mnwe_0 \mapsto w^{-1}n$ determines a bijection

$$(12.0.2) \quad \alpha_w^{-1}(m'e'_0) = N'/N' \cap I.$$

We warn the reader that α_w is not a morphism of ind-schemes; however its restriction to the inverse image of any connected component of $M'(L)/I_{M'}$ is a morphism of ind-schemes.

Now for $x \in \widetilde{W}$, and w, b as above, define the finite set

$$S_P(x, w) := \{y \in \widetilde{W}_{M'} : N'\mathbf{a}_y \cap I\mathbf{a}_x \neq \emptyset\}.$$

Note that $N'\mathbf{a}_y \cap I\mathbf{a}_x \neq \emptyset \Leftrightarrow I_{P'}\mathbf{a}_y \cap I\mathbf{a}_x \neq \emptyset$. For a given x , there are only finitely many y such that the latter holds; see Proposition 11.1.4.

The following proposition is an analogue of part of [GHKR], Prop. 5.6.1.

Proposition 12.0.6. (1) *The map α_w restricts to give a surjective map*

$$(12.0.3) \quad \beta_w : X_x^G(b) \cap Pwe_0 \longrightarrow \bigcup_{y \in S_P(x, w)} X_y^{M'}(\tilde{b}).$$

(2) *Assume $X_x^G(b) \cap Pwe_0 \neq \emptyset$. For a fixed $m' \in M'(L)$ such that $m'e'_0 \in X_y^{M'}(\tilde{b})$, set $b' := m'^{-1}\tilde{b}\sigma(m') \in I_{M'}yI_{M'}$. Then the fiber $\beta_w^{-1}(m'e'_0)$ is a locally finite-type algebraic variety having dimension*

$$\dim \beta_w^{-1}(m'e'_0) = \dim(I\mathbf{a}_x \cap N'\mathbf{a}_y) - \langle \rho, \nu + \nu_{\text{dom}} \rangle,$$

a number which depends on y but not on $m'e'_0$.

(3) *We have*

$$\dim X_x^G(b) = \sup_{w, y : y \in S_P(x, w)} \{\dim(I\mathbf{a}_x \cap w^{-1}N\mathbf{a}_y) + \dim(X_y^{w^{-1}M}(w^{-1}b))\} - \langle \rho, \nu + \nu_{\text{dom}} \rangle.$$

The proposition implies that, modulo knowledge of certain basic cases (i.e., the $X_y^{M'}(\tilde{b})$), there is a finite algorithm to determine the non-emptiness and dimension of $X_x^G(b)$. Conjecture 9.3.2 predicts a finite algorithm to determine the non-emptiness of each $X_y^{M'}(\tilde{b})$. Thus, in effect it predicts a finite algorithm for the non-emptiness of $X_x^G(b)$ itself.

Corollary 12.0.7. *We have $X_x^G(b) \neq \emptyset$ if and only if there exist $w \in {}^M W$ and $y \in S_P(x, w)$ with $X_y^{M'}(\tilde{b}) \neq \emptyset$.*

Proof of Proposition: It is clear that α_w sends the left hand side of (12.0.3) into the right hand side. If $m'e'_0 \in X_y^{M'}(\tilde{b})$, then the isomorphism (12.0.2) restricts to give an isomorphism

$$(12.0.4) \quad \beta_w^{-1}(m'e'_0) = f_{b'}^{-1}(IxIb'^{-1} \cap N')/N' \cap I,$$

where $b' := m'^{-1}\tilde{b}\sigma(m')$ and where we define

$$\begin{aligned} f_{b'} : N' &\longrightarrow N' \\ n' &\longrightarrow n'^{-1}b'\sigma(n')b'^{-1}. \end{aligned}$$

Since $f_{b'}$ is surjective (see Remark 11.2.3) and $IxI \cap N'b' \neq \emptyset$, we see that β_w is surjective, proving (1). Also, the fibers of β_w are algebraic varieties locally of finite type, and their dimension can be computed from (12.0.4) using the method of the proof of Theorem 11.2.1. This proves (2). Finally, (3) follows from (1) and (2). \square

Remark 12.0.8. For affine Deligne-Lusztig varieties in the affine Grassmannian, it is known that $X_\mu^G(b) \neq \emptyset$ if and only if $[b] \in B(G, \mu)$ (cf. [KR],[K3],[Lu],[Ga]). The condition $[b] \in B(G, \mu)$ means that $\eta_G(b) = \mu$ in Λ_G and $\bar{\nu}_b \leq \mu$ (“Mazur’s inequality”). For $X_x^G(b)$, where as before we take $b \in \Omega_M$, one might ask for the analogues of “Mazur’s inequalities,” where by this we mean a family of congruence conditions and inequalities imposed on x, b and $\bar{\nu}_b$ which hold if and only if $X_x^G(b)$ is non-empty. In light of the above proposition, we see that, whatever Mazur’s inequalities end up being, they should hold if and only if there exists $w \in {}^M W$ such that for some $y \in \widetilde{W}_{w^{-1}M}$, we have

- $w^{-1}Ny \cap IxI \neq \emptyset$ and
- $X_y^{w^{-1}M}(w^{-1}b) \neq \emptyset$.

In view of Conjecture 9.3.2, the second item should be understood as a family of congruence conditions. The first item should correspond to a family of inequalities and congruence conditions between $x, y \in \widetilde{W}$. Taken together the inequalities will be somewhat stronger than the condition $y \leq x$ in the Bruhat order on \widetilde{W} .

13. FUNDAMENTAL ALCOVES AND THE SUPERSET METHOD

13.1. Fundamental alcoves. We now single out some alcoves that will be used to generalize Reuman’s superset method [Re2] to all σ -conjugacy classes in $G(L)$.

Definition 13.1.1. For $x \in \widetilde{W}$ we say that $x\mathfrak{a}$ is a fundamental alcove if every element of IxI is σ -conjugate under I to x .

Equivalently, the alcove $x\mathfrak{a}$ is fundamental if every element of xI is σ -conjugate under $xI \cap I$ to x .

Now let $P = MN$ be a semistandard parabolic subgroup of G . There is then an Iwahori decomposition $I = I_N I_M I_{\overline{N}}$. We use the Iwahori subgroup I_M of $M(L)$ to form the subgroup $\Omega_M \subset \widetilde{W}_M$; note that the canonical surjective homomorphism $\widetilde{W}_M \rightarrow \Lambda_M$ restricts to an isomorphism $\Omega_M \cong \Lambda_M$. We compose this isomorphism with the canonical homomorphism $\Lambda_M \rightarrow \mathfrak{a}_M$, obtaining a homomorphism $\Omega_M \rightarrow \mathfrak{a}_M$; for $x \in \Omega_M$ we will denote by $\nu_x \in \mathfrak{a}_M$ the image of x under this homomorphism. Note that $x \mapsto \nu_x$ is intrinsic to M and has nothing to do with P .

Definition 13.1.2. For $x \in \widetilde{W}_M$ we say that $x\mathfrak{a}$ is a fundamental P -alcove if it is a P -alcove for which $x \in \Omega_M$, or, in other words, if $xI_M x^{-1} = I_M$, $xI_N x^{-1} \subset I_N$, and $x^{-1}I_{\overline{N}}x \subset I_{\overline{N}}$.

Proposition 6.3.1 implies that any fundamental P -alcove is a fundamental alcove, just as the terminology suggests. An obvious question (that we have not tried to answer) is whether any fundamental alcove arises as a fundamental P -alcove for some semistandard P .

The next result gives some insight into P -alcoves, although we will make only incidental use of it. We write $\rho_N \in \mathfrak{a}^*$ for the half-sum of the elements in R_N .

Proposition 13.1.3. Write Ω_P for the set of $x \in \Omega_M$ such that $x\mathfrak{a}$ is a fundamental P -alcove.

- (1) Ω_P is a submonoid of Ω_M .
- (2) Let $x, y \in \Omega_P$. Then $IxIyI = IxyI$ and $\ell(x) + \ell(y) = \ell(xy)$. Here ℓ is the usual length function on \widetilde{W} .

(3) Let $x \in \Omega_P$. Then $\ell(x) = \langle 2\rho_N, \nu_x \rangle$.

Proof. (1) This is clear from the definitions.

(2) For the first statement just note that

$$xIy = (xI_Nx^{-1})xy(y^{-1}I_My)(y^{-1}I_{\overline{N}}y) \subset I_NxyI_MI_{\overline{N}} \subset IxyI.$$

The second statement follows from the first (easy, and presumably well-known).

(3) Since both the left and right sides of the equality to be proved are additive functions on the monoid Ω_P , we may replace x by x^m for any positive integer m . Taking m to be the order of the image of x in \widetilde{W}_M , we are reduced to the case in which x is a translation element lying in Ω_P . Such an element is of the form ϵ^μ for some cocharacter $\mu \in X_*(A)$ whose image is central in M and dominant with respect to any Borel subgroup of P containing A . It is easy to see that ν_x is simply the image of μ under the canonical inclusion of $X_*(A)$ in \mathfrak{a} . Thus the equality to be proved is a consequence of the equality $\ell(\epsilon^\mu) = \langle 2\rho_N, \mu \rangle$, which in turn follows from the usual formula for the length of translation elements in \widetilde{W} , in view of the fact that all roots of M vanish on μ . \square

13.2. Levi subgroups adapted to I . Let M be a Levi subgroup of G containing A . Once again we put $I_M = M(L) \cap I$ and form $\Omega_M \subset \widetilde{W}_M$ relative to I_M . We will also make use of the homomorphism $x \mapsto \nu_x$ from Ω_M to \mathfrak{a}_M that was explained in the previous subsection.

We write $\mathcal{P}(M)$ for the set of parabolic subgroups of G having M as Levi component. For $P \in \mathcal{P}(M)$ we define $\Omega_M^{\geq 0}$ (respectively, $\Omega_M^{> 0}$) to be the set of elements $x \in \Omega_M$ such that $\langle \alpha, \nu_x \rangle \geq 0$ (respectively, $\langle \alpha, \nu_x \rangle > 0$) for all $\alpha \in R_N$. It is clear that most elements of $\Omega_M^{\geq 0}$ lie in Ω_P ; however, we are going to give a condition on M which will guarantee that every element of $\Omega_M^{\geq 0}$ lies in Ω_P . (Compare this with Remark 7.2.4, which shows that when $P = MN$ is standard, an element $\epsilon^\lambda \in \Omega_M$ lies in Ω_P if and only if λ is G -dominant.)

As usual the group \widetilde{W}_M acts by affine linear transformations on both \mathfrak{a} and its quotient $\mathfrak{a}/\mathfrak{a}_M$, the natural surjection $\mathfrak{a} \rightarrow \mathfrak{a}/\mathfrak{a}_M$ being \widetilde{W}_M -equivariant. The subgroup Ω_M then inherits an action on \mathfrak{a} and $\mathfrak{a}/\mathfrak{a}_M$.

Definition 13.2.1. *We say that M is adapted to I (respectively, weakly adapted to I) if there exists $\lambda \in \mathfrak{a}$ (respectively, in the closure of \mathfrak{a}) whose image in $\mathfrak{a}/\mathfrak{a}_M$ is fixed by the action of Ω_M .*

For any such λ it is easy to see that $x\lambda = \lambda + \nu_x$ for all $x \in \Omega_M$.

Proposition 13.2.2. *If M is adapted to I , then $\Omega_M^{\geq 0} \subset \Omega_P$, and consequently for every $x \in \Omega_M$ there exists $P \in \mathcal{P}(M)$ for which $x\mathfrak{a}$ is a fundamental P -alcove. Similarly, if M is weakly adapted to I , then $\Omega_M^{> 0} \subset \Omega_P$.*

Proof. We begin by proving the first statement. For $\alpha \in R_N$ we must show that $x\mathfrak{a} \geq_\alpha \mathfrak{a}$, which is to say that $k(\alpha, x\mathfrak{a}) \geq k(\alpha, \mathfrak{a})$. For any $\lambda \in \mathfrak{a}$ we have $k(\alpha, x\mathfrak{a}) = \lceil \alpha(x\lambda) \rceil$ and $k(\alpha, \mathfrak{a}) = \lceil \alpha(\lambda) \rceil$. Now pick λ as in the definition of being adapted to I . Since $x \in \Omega_M^{\geq 0}$, we see from the equality $x\lambda = \lambda + \nu_x$ that $\alpha(x\lambda) \geq \alpha(\lambda)$; it is then clear that $\lceil \alpha(x\lambda) \rceil \geq \lceil \alpha(\lambda) \rceil$.

Now we prove the second statement. For $\alpha \in R_N$ we now have

$$k(\alpha, \mathfrak{a}) - 1 \leq \alpha(\lambda) < \alpha(x\lambda) \leq k(\alpha, x\mathfrak{a})$$

and hence $k(\alpha, \mathfrak{a}) \leq k(\alpha, x\mathfrak{a})$, as desired. \square

Proposition 13.2.3. *Let M be any Levi subgroup containing A . Then there exists $w \in W$ such that wM is adapted to I .*

Proof. There exist fixed points of Ω_M on $\mathfrak{a}/\mathfrak{a}_M$ lying on no affine root hyperplane for M (for example, when M is simple, one can take the barycenter of the base alcove for \widetilde{W}_M). We choose such a fixed point $\bar{\lambda}$ and then choose $\lambda \in \mathfrak{a}$ mapping to $\bar{\lambda}$. We are free to add any element of \mathfrak{a}_M to λ , so we may assume that λ lies on no affine root hyperplane for G . If λ happens to lie in \mathfrak{a} , then M is adapted to I . In any case there exists a unique alcove $x'\mathfrak{a}$ containing λ . The Levi subgroup is then adapted to $I' = x'Ix'^{-1}$. Taking w to be the inverse of the image of x' in W , we find that wM is adapted to I . \square

Being adapted to I is quite a strong condition on M . It is important to realize that standard Levi subgroups are often not adapted to our standard Iwahori subgroup I , even though both notions of standard are tied to the same Borel subgroup.

Corollary 13.2.4. *For every $[b] \in B(G)$ there exists a semistandard representative $x \in \widetilde{W}$ of $[b]$ such that $x\mathfrak{a}$ is a fundamental alcove and hence $IxI \subset [b]$.*

Proof. This follows from the previous two propositions and Definition 7.2.3. \square

13.3. Superset method. Let $b \in G(L)$. The *superset* $\widetilde{W}(b)$ associated to b is the set of $x \in \widetilde{W}$ such that IxI is contained in $Iy^{-1}IbIyI$ for some $y \in \widetilde{W}$. The reason for the name superset is that the set of $x \in \widetilde{W}$ such that $X_x(b) \neq \emptyset$ is contained in $\widetilde{W}(b)$. Indeed, if $X_x(b) \neq \emptyset$, then there exists $g \in G(L)$ such that $g^{-1}b\sigma(g) \in IxI$. There also exists $y \in \widetilde{W}$ such that $g \in IyI$, and then

$$IxI = Ig^{-1}b\sigma(g)I \subset Iy^{-1}IbIyI.$$

Proposition 13.3.1. *Suppose that $x_0\mathfrak{a}$ is a fundamental alcove, and let b_0 be any element of Ix_0I . Then*

$$\{x \in \widetilde{W} : X_x(b_0) \neq \emptyset\} = \widetilde{W}(b_0).$$

Proof. We already know the inclusion \subset . To establish \supset we consider $x \in \widetilde{W}(b_0)$ and choose $y \in \widetilde{W}$ such that $IxI \subset Iy^{-1}Ib_0IyI$. Then IxI meets $y^{-1}Ib_0Iy$, and since (by our hypothesis on x_0) every element of Ib_0I has the form $i^{-1}b_0\sigma(i)$ for suitable $i \in I$, there is some element in IxI of the form $\dot{y}^{-1}i^{-1}b_0\sigma(i)\dot{y}$, where \dot{y} is a representative of y in the F -points of the normalizer of A in G . Since $\dot{y} = \sigma(\dot{y})$, this shows that IxI meets $[b_0]$, as desired. \square

Corollary 13.3.2. *For every $[b] \in B(G)$ there is a semistandard representative $b_0 \in [b]$ for which the superset method applies, yielding*

$$\{x \in \widetilde{W} : X_x(b_0) \neq \emptyset\} = \widetilde{W}(b_0).$$

Proof. Combine Corollary 13.2.4 with Proposition 13.3.1. \square

REFERENCES

- [Be] E. T. Beazley, *Codimensions of Newton strata for $SL_3(F)$ in the Iwahori case*, arXiv:0711.3820v1
- [Bo] A. Borel, *Linear Algebraic Groups*, Springer Graduate Text in Mathematics 126 (1991)
- [BT] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. I*, Inst. Hautes Études Sci. Publ. Math. **41** (1972), 5–251.

- [Ch] C.-L. Chai, *Newton polygons as lattice points*, Amer. J. Math. **122** (2000), no. 5, 967–990.
- [Ga] Q. Gashi, *A conjecture of Kottwitz and Rapoport for split groups*, preprint in preparation.
- [GP] M. Geck, G. Pfeiffer, *Characters of finite Coxeter groups and Iwahori-Hecke algebras*, London Math. Soc. Monographs, New Series **21** (2000)
- [GHKR] U. Görtz, T. Haines, R. Kottwitz, D. Reuman, *Dimensions of some affine Deligne-Lusztig varieties*, Ann. sci. École Norm. Sup. 4^e série, t. **39** (2006), 467–511.
- [HN] T. Haines, B.C. Ngô, *Alcoves associated to special fibers of local models*, Amer. J. Math. **124** (2002), 1125–1152.
- [K1] R. Kottwitz, *Isocrystals with additional structure*, Compositio Math. **56** (1985), 201–220.
- [K2] R. Kottwitz, *Isocrystals with additional structure. II*, Compositio Math. **109** (1997), 255–339.
- [K3] R. Kottwitz, *On the Hodge-Newton decomposition for split groups*, Int. Math. Res. Not. **26** (2003), 1433–1447.
- [K4] R. Kottwitz, *Dimensions of Newton strata in the adjoint quotient of reductive groups*, Pure Appl. Math. Q. **2** (2006), no. 3, 817–836.
- [KR] R. Kottwitz, M. Rapoport, *On the existence of F -crystals*, Comment. Math. Helv. **78** (2003), no.1, 153–184.
- [MP] A. Moy, G. Prasad, *Unrefined minimal K -types for p -adic groups*, Invent. Math. **116** (1994), 393–408.
- [La] J.-P. Labesse, *Fonctions élémentaires et lemme fondamental pour la changement de base stable*, Duke Math J. **61**, no. 2 (1990), 519–530.
- [Lu] C. Lucarelli, *A converse to Mazur’s inequality for split classical groups*, J. Inst. Math. Jussieu **3** (2004), no. 2, 165–183.
- [MV] E. Mantovan, E. Viehmann, *On the Hodge-Newton filtration for p -divisible O -modules*, preprint. arXiv:0701.4194.
- [PR] G. Pappas, M. Rapoport, *Twisted loop groups and their affine flag varieties*, arXiv:math/0607130v1.
- [Ra] M. Rapoport, *A guide to the reduction modulo p of Shimura varieties*, Astérisque (2005), no. 298, 271–318.
- [Re1] D. Reuman, *Determining whether certain affine Deligne-Lusztig sets are non-empty*, Thesis Chicago 2002, math.NT/0211434.
- [Re2] D. Reuman, *Formulas for the dimensions of some affine Deligne-Lusztig varieties*, Michigan Math. J. **52** (2004), no. 2, 435–451.
- [Ro] G. Rousseau, *Euclidean buildings*, preprint Nancy, 27 pp. In “Géométries à courbure négative ou nulle, groupes discrets et rigidité”, école d’été Grenoble, 14 juin-2 juillet 2004, to appear in Séminaires et congrès **18**, Société Math. de France. <http://hal.archives-ouvertes.fr/hal-00094363>
- [V1] E. Viehmann. *The dimension of affine Deligne-Lusztig varieties*, Ann. sci. École Norm. Sup. 4^e série, t. **39** (2006), 513–526.
- [V2] E. Viehmann, *Connected components of closed affine Deligne-Lusztig varieties*, Math. Ann. **340**, No. 2 (2008), 315–333.

ULRICH GÖRTZ, MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, BERINGSTR. 1, 53115 BONN, GERMANY

E-mail address: ugoertz@math.uni-bonn.de

THOMAS J. HAINES, MATHEMATICS DEPARTMENT, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742-4015

E-mail address: tjh@math.umd.edu

ROBERT E. KOTTWITZ, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 UNIVERSITY AVENUE, CHICAGO, ILLINOIS 60637

E-mail address: kottwitz@math.uchicago.edu

DANIEL C. REUMAN, IMPERIAL COLLEGE LONDON, SILWOOD PARK CAMPUS, BUCKHURST ROAD, ASCOT, BERKSHIRE, SL5 7PY, UNITED KINGDOM

E-mail address: d.reuman@imperial.ac.uk