THE TOPOLOGY OF THE RELATIVE CHARACTER VARIETIES OF A QUADRUPLY-PUNCTURED SPHERE

ROBERT L. BENEDETTO AND WILLIAM M. GOLDMAN

ABSTRACT. Let M be a quadruply-punctured sphere with boundary components A, B, C, D. The $\mathrm{SL}(2,\mathbb{C})$-character variety of M consists of equivalence classes of homomorphisms ρ of $\pi_1(M) \longrightarrow \mathrm{SL}(2,\mathbb{C})$ and can be identified with a quartic hypersurface in $\mathbb{C}^7$. For fixed $a, b, c, d \in \mathbb{C}$, the subset $V_{a,b,c,d}$ corresponding to representations ρ with $\mathrm{tr}(\rho(A)) = a$, $\mathrm{tr}(\rho(B)) = b$, $\mathrm{tr}(\rho(C)) = c$, $\mathrm{tr}(\rho(D)) = d$ is a cubic surface in $\mathbb{C}^3$. We determine the singular points of $V_{a,b,c,d}$, classify its set $V_{a,b,c,d}(\mathbb{R})$ of $\mathbb{R}$-points into six topological types, at least when $V_{a,b,c,d}(\mathbb{R})$ is nonsingular. $V_{a,b,c,d}(\mathbb{R})$ contains a compact component if and only if $-2 < a, b, c, d < 2$. For certain (a, b, c, d), this component corresponds to representations in $\mathrm{SL}(2,\mathbb{R})$.

CONTENTS

1. Introduction 1
2. Character varieties for surfaces 4
2.1. Cyclic groups 4
2.2. The triply-punctured sphere 7
2.3. Properties of $\kappa(x, y, z)$ 8
2.4. The quadruply-punctured sphere 9
2.5. Amalgamating representations 9
2.6. Defining equations 10
2.7. Symmetries 11
3. Singular points and reducible characters 13
3.1. A criterion for singularity 13

Date: October 26, 1997.

1991 *Mathematics Subject Classification.* 57M05 (Low-dimensional topology).

Key words and phrases. Surface, Fundamental group, Character variety, Representation variety.

Goldman gratefully acknowledges partial support from National Science Foundation grants DMS-8613576, DMS-8902619, DMS-9205139, DMS-9504764, the Alfred P. Sloan Foundation, and the University of Maryland Institute for Advanced Computer Studies.

3.2. Solutions of quadratic equations 13
3.3. Singularities of $V_{a,b,c,d}$ 15
3.4. A discriminant polynomial 17
4. Topology of the set of $\mathbb{R}$-points 18
4.1. Singular points 18
4.2. Topological classification of the smooth real varieties 20
4.3. Reduction to a plane quartic curve 20
4.4. The six cases 26
5. A compact component of $\mathrm{SL}(2,\mathbb{R})$-representations 32
References 36
List of Figures 36

1. Introduction

Moduli spaces of representations of fundamental groups of surfaces arise in many natural algebraic, geometric, and analytic problems. In particular, the classification of geometric structures on manifolds and solutions of the Yang-Mills equations from gauge theory lead to moduli spaces closely related to representations of surface groups. Furthermore these moduli spaces enjoy a rich symmetry due to the large topological symmetry groups of surfaces. The present paper describes in detail the topology of some of these moduli spaces, in the case of surfaces homeomorphic to a quadruply-punctured sphere.

Decompositions of surfaces into subsurfaces provide a technique for analyzing the moduli spaces in terms of moduli spaces for simpler surfaces. For example, every compact connected surface M of genus g with b boundary components decomposes along $3g+b-3$ simple closed curves into $2g+b-2$ "pants", provided that

$$0 > \chi(M) = 2 - 2g - b.$$

A "pants" is a triply-punctured sphere, that is, a surface of genus 0 with three boundary components. In general, we shall refer to a connected surface of genus g with b boundary components as a *b-punctured surface of genus g,* although, strictly speaking the ends of a "punctured surface" are noncompact and do not have boundary. Similarly, a *b-punctured torus* is a surface of genus 1 with b boundary components. Surfaces of small negative Euler characteristic are the building blocks for all surfaces of negative Euler characteristic and their moduli spaces are the building blocks for moduli spaces of more complicated surfaces.

This decomposition technique has been used in [7] to study the automorphism group of surfaces and in [5] to analyze moduli spaces of geometric structures on surfaces.

This paper concerns the moduli space of representations of a quadruply-punctured sphere into $\mathrm{SL}(2,\mathbb{C})$. This moduli space is naturally parametrized by the moduli of the images of the boundary components, which are described by four scalar parameters (a,b,c,d). For fixed values of (a,b,c,d), the moduli space is a surface $V_{a,b,c,d}$. We determine for which (a,b,c,d) is $V_{a,b,c,d}$ nonsingular.

Theorem. *For fixed $a,b,c,d \in \mathbb{C}$, the surface $V_{a,b,c,d}$ singular if and only if either at least one of a,b,c,d equals ± 2 or there is a reducible representation ρ with boundary traces a,b,c,d. The latter case occurs when $\Delta(a,b,c,d)=0$ where $\Delta(a,b,c,d)$ is the polynomial*

$$\begin{aligned}\Delta(a,b,c,d) = &\left(2(a^2+b^2+c^2+d^2)-abcd-16\right)^2 \\ &-(4-a^2)(4-b^2)(4-c^2)(4-d^2)\end{aligned}$$

In this case $V_{a,b,c,d}(\mathbb{C})$ is a cubic surface in $\mathbb{C}^3$ whose projective completion $\bar{V}_{a,b,c,d}(\mathbb{C})$ is smooth. Furthermore $\bar{V}_{a,b,c,d}(\mathbb{C}) \setminus V_{a,b,c,d}$ consists of three lines in general position, and using the classification of cubic surfaces, one proves that the set of $\mathbb{C}$-points of $V_{a,b,c,d}$ deformation retracts to a bouquet of five 2-spheres.

However, the topologies of the real varieties are much more varied. For $(a,b,c,d) \in \mathbb{R}$, the set $V_{a,b,c,d}(\mathbb{R})$ of $\mathbb{R}$-points of $V_{a,b,c,d}$, at least when $V_{a,b,c,d}(\mathbb{R})$ is nonsingular, one of six topological types, depending on the boundary parameters a,b,c,d. These six topological types are depicted in Figures 11, 12, 13, 14, 15, and 16. Let n denote the number of boundary traces a,b,c,d which lie in the interval $(-2,2)$:

Theorem. *Let $a,b,c,d \in \mathbb{R}$ be such that $V_{a,b,c,d}$ is nonsingular. Then the set $V_{a,b,c,d}(\mathbb{R})$ of $\mathbb{R}$-points is homeomorphic to one of the following:*

1. *A quadruply-punctured sphere if $n=0$ and $abcd<0$;*
2. *A disjoint union of a triply-punctured torus and a disc if $n=0$ and $abcd>0$;*
3. *A disjoint union of a triply-punctured sphere and a disc if $n=1$;*
4. *A disjoint union of an annulus and two discs if $n=2$;*
5. *A disjoint union of four discs if $n=3$;*
6. *A disjoint union of four discs and a sphere if $n=4$.*

Corollary. *For a,b,c,d and n as above, the Euler characteristic of $V_{a,b,c,d}(\mathbb{R})$ is given by the following formula:*

$$\chi(V_{a,b,c,d}(\mathbb{R})) = 2n-2$$

The points of $V_{a,b,c,d}(\mathbb{R})$ correspond to representations into either $\mathrm{SU}(2)$ or $\mathrm{SL}(2,\mathbb{R})$. Components of spaces of representations into $\mathrm{SU}(2)$ are always compact, and in the analogous case where M is a once-punctured torus, the only compact components of relative real character varieties of M correspond to $\mathrm{SU}(2)$-representations. When $-2 < a, b, c, d < 2$, the variety $V_{a,b,c,d}(\mathbb{R})$ contains a compact component, and we determine conditions when this compact component corresponds to representations in $\mathrm{SU}(2)$ or $\mathrm{SL}(2,\mathbb{R})$.

Proposition. *Suppose $a, b, c, d \in (-2, 2)$ and $V_{a,b,c,d}$ is smooth. Then $V_{a,b,c,d}(\mathbb{R})$ contains a compact component, which consists of characters of $\mathrm{SL}(2,\mathbb{R})$-representations if and only if*

$$\Delta(a,b,c,d) > 0$$
$$16 - abcd - 2(a^2 + b^2 + c^2 + d^2) > 0.$$

Otherwise, each element in the component is the character of an $\mathrm{SU}(2)$-representation.

The component of $V_{a,b,c,d}(\mathbb{R})$ homeomorphic to a disc corresponds to the "Teichmüller space" of M and consists of characters of holonomy representations of hyperbolic structures on M with geodesic boundary. In all of the last five cases, the component which is homeomorphic to a disc admits a similar interpretation in terms of hyperbolic structures with conical singularities; see [5].

This investigation began as "Research Experiences for Undergraduates" project at the University of Maryland in the summers of 1992–1993, under Goldman's direction. During this project, Benedetto used Geomview [4] and Mathematica [10] software on Silicon Graphics workstations to visualize the components of the real moduli spaces. Many of the results in this paper were suggested by computer experiments before they were rigorously proved. (Compare Benedetto's senior thesis [1]). The illustrations in this paper were drawn using Geomview and Mathematica software on Silicon Graphics and NeXT workstations.

2. Character varieties for surfaces

Let $\mathbf{k}$ denote a field (usually either $\mathbb{R}$ or $\mathbb{C}$) and let $G = \mathrm{SL}(2,\mathbf{k})$. For the moment let π denote an arbitrary finitely generated group. We denote the set of homomorphisms $\pi \longrightarrow G$ by $\mathrm{Hom}(\pi, G)$; since G is an affine algebraic group defined over $\mathbf{k}$, $\mathrm{Hom}(\pi, G)$ naturally enjoys the structure of the set of $\mathbf{k}$-points of an affine algebraic set defined over $\mathbb{Z}$. (Compare Lubotzky-Magid [8],§1.) A central problem is to determine the global topology and geometric structure on the space $\mathrm{Hom}(\pi, G)$.

We would like to consider the space of equivalence classes of such representations under the natural action of $\mathrm{SL}(2,\mathbb{C})$ by conjugation, but the quotient space is generally not well-behaved. By enlarging the equivalence relation slightly, one can define a *categorical quotient* $\mathrm{Hom}(\pi, G)//G$ (in the sense of algebraic geometry) which does have the structure of an affine algebraic set. For brevity denote $\mathrm{Hom}(\pi, G)$ by X. The guiding principle is that the regular functions on the quotient $X//G$ should be the G-invariant regular functions on X, and thus the quotient $X//G$ is *defined* as $\mathrm{Spec}(\mathbf{k}[X])$. The points of $X//G$ correspond to equivalence classes, where two orbits are considered equivalent if they cannot by distinguished by G-invariant regular functions on X. In general there is a G-invariant Zariski-dense subset X^- such that the restriction of the quotient map $X \longrightarrow X//G$ to X^- defines a bijection of the orbit space X^-/G to $X//G$. In practice, we can take (assuming $\mathbf{k}$ is algebraically closed) X^- to be the set of *completely reducible* (or *semisimple*) representations $\pi \longrightarrow G$. For a careful treatment of this material, we refer the reader to Lubotzky-Magid [8],§1.

2.1. **Cyclic groups.** When π is a cyclic group, then a homomorphism $\pi \longrightarrow G$ is completely determined by the image of a generator of π, and every regular function on G which is invariant under conjugation is a polynomial in the trace function

$$\begin{aligned} G &\longrightarrow \mathbf{k} \\ A &\longmapsto \mathrm{tr}(A) \end{aligned}$$

Thus $X \cong \mathbf{k}$ under the trace mapping. Note that the trace function has critical values ± 2, with critical points at the center $\pm I$.

More generally, suppose that π is a free group freely generated by $x_1, \ldots, x_n$. Then $\mathrm{Hom}(\pi, G)$ identifies with the Cartesian product G^n and Procesi [12] proved that every G-invariant function on G^n is a polynomial in finitely many functions of the form

$$\begin{aligned} G^n &\longrightarrow \mathbf{k} \\ (A_1, \ldots, A_n) &\longmapsto \mathrm{tr}\, w(A_1, \ldots, A_n) \end{aligned}$$

where $w(x_1, \ldots, x_n)$ is a word in $x_1, \ldots, x_n$, that is, an element of π.

For example, for $n = 2$ and $\mathbf{k} = \mathbb{C}$, classical work (dating at least as far back as Fricke and Klein [3]) implies the following fundamental result:

Theorem 2.1.1. *The invariant function*

$$\chi : \mathrm{Hom}(\pi, G) \longrightarrow \mathbb{C}^3$$
$$\rho \longmapsto \begin{bmatrix} \mathrm{tr}(\rho(x_1)) \\ \mathrm{tr}(\rho(x_2)) \\ \mathrm{tr}(\rho(x_1 x_2)) \end{bmatrix}$$

defines an isomorphism $X \longrightarrow \mathbb{C}^3$. *That is, every function of pairs* $(A_1, A_2) \in \mathrm{SL}(2,\mathbb{C}) \times \mathrm{SL}(2,\mathbb{C})$ *invariant under* simultaneous conjugation

$$(A_1, A_2) \mapsto (TA_1T^{-1}, TA_2T^{-1})$$

can be expressed is a function of $\mathrm{tr}(A_1)$, $\mathrm{tr}(A_2)$, *and* $\mathrm{tr}(A_1A_2)$. *Furthermore, if* A_1, A_2 *generate an irreducible representation on* $\mathbb{C}^2$, *then for any other pair* A_1', A_2' *with*

$$\mathrm{tr}(A_1) = \mathrm{tr}(A_1')$$
$$\mathrm{tr}(A_2) = \mathrm{tr}(A_2')$$
$$\mathrm{tr}(A_1A_2) = \mathrm{tr}(A_1'A_2')$$

there exists $T \in \mathrm{SL}(2,\mathbb{C})$ *with* $A_1' = TA_1T^{-1}$, $A_2' = TA_2T^{-1}$.

We call the triple $\chi(\rho)$ the *character* of ρ, since from it one can deduce the trace $\mathrm{tr}(\rho(A))$ for any $A \in \pi$. (Compare Magnus [9].)

The criterion for reducible representations is slightly more complicated. If ρ is reducible, then it is conjugate to a representation in upper-triangular matrices

$$\rho(\gamma) = \begin{bmatrix} a(\gamma) & b(\gamma) \\ 0 & a(\gamma)^{-1} \end{bmatrix}$$

It and its *semisimplification*

$$\rho^s(\gamma) = \begin{bmatrix} a(\gamma) & 0 \\ 0 & a(\gamma)^{-1} \end{bmatrix}$$

have the same character, but in general will not be conjugate. Indeed, ρ^s lies in the closure of the orbit of ρ. Two reducible representations have the same character if and only if their semisimplifications are conjugate, or equivalently, if their orbit closures are equal.

For example, the G-invariant function on $G \times G$ defined by

$$f : (A_1, A_2) \mapsto \mathrm{tr}(A_1A_2A_1^{-1}A_2^{-1})$$

is given by the expression

$$f(A_1, A_2) = \kappa(\mathrm{tr}(A_1), \mathrm{tr}(A_2), \mathrm{tr}(A_1A_2))$$

where $\kappa \in \mathbb{C}[x,y,z]$ is the cubic polynomial defined by:

$$\kappa(x,y,z) = x^2 + y^2 + z^2 - xyz - 2 \tag{2.1.1}$$

A slight variant of this polynomial will be useful later on. Let $a, b \in \mathbf{k}$. Then $\kappa_{a,b}(x) \in \mathbf{k}[x]$ is the *quadratic* polynomial defined by:

$$\kappa_{a,b}(x) = \kappa(a,b,x) - 2 = x^2 - abx + (a^2 + b^2 - 4) \tag{2.1.2}$$

The reducible representations occur for characters (x,y,z) with

$$\kappa(x,y,z) = 2$$

where $\kappa : \mathbb{C}^3 \longrightarrow \mathbb{C}$ is defined by (2.1.1). We refer to such a character as a *reducible character.*

The set of $\mathbb{R}$-points in $\mathfrak{X}(M)$ equals $\mathfrak{X}(M)(\mathbb{R}) = \mathbb{R}^3$. By Prop. III.1.1 of Morgan-Shalen [11] (p.458), real characters correspond to representations taking values in the real forms SU(2) and $\mathrm{SL}(2,\mathbb{R})$ of $\mathrm{SL}(2,\mathbb{C})$. A point $(x,y,z) \in \mathbb{R}^3$ corresponds to an SU(2)-representation if and only if

$$-2 \leq x,y,z \leq 2 \qquad \text{and} \qquad \kappa(x,y,z) \leq 2.$$

Otherwise $(x,y,z) \in \mathbb{R}^3$ corresponds to an $\mathrm{SL}(2,\mathbb{R})$-representation. A character $(x,y,z) \in \mathbb{R}^3$ corresponds to both an SU(2)-representation and an $\mathrm{SL}(2,\mathbb{R})$-representation if and only if it corresponds to a representation in

$$\mathrm{SO}(2) = \mathrm{SU}(2) \cap \mathrm{SL}(2,\mathbb{R});$$

such a representation is necessarily reducible. See Goldman [6] for details.

2.2. **The triply-punctured sphere.** We interpret Theorem 2.1.1 in terms of the the triply-punctured sphere. For any compact manifold, denote

$$\mathfrak{X}(M) = \prod_i \mathrm{Hom}(\pi_1(M_i), G)//G$$

where

$$M = \coprod_i M_i$$

is the decomposition of M into connected components. When M is a compact surface with boundary

$$\partial M = \coprod_{i=1}^{m} \partial_i M$$

each component $\partial_i M$ has fundamental group $\pi_1(\partial_i M) \cong \mathbb{Z}$. Thus each $\mathfrak{X}(\partial_i M) \cong \mathbb{C}$ and $\mathfrak{X}(\partial M) \cong \mathbb{C}^m$. The homomorphism of fundamental groups

$$\pi_1(\partial_i M) \longrightarrow \pi_1(M)$$

induces a restriction map

$$\mathfrak{X}(M) \longrightarrow \mathfrak{X}(\partial_i M) \cong \mathbb{C}.$$

Taking the product, we obtain

$$\partial_* : \mathfrak{X}(M) \longrightarrow \mathbb{C}^m$$

Let M denote a triply-punctured sphere. Its fundamental group admits the redundant geometric presentation

$$\pi = \langle X, Y, Z \mid XYZ = I \rangle$$

where X, Y, Z are simple loops correponding to the three boundary components. In this case the boundary map ∂_* defines an isomorphism $\mathfrak{X}(M) \approx \mathbb{C}^3$, where

$$\partial_*([\rho]) = (\mathrm{tr}(\rho(X)), \mathrm{tr}(\rho(Y)), \mathrm{tr}(\rho(Z))).$$

2.3. **Properties of $\kappa(x, y, z)$.** The polynomial κ and its variant $\kappa_{a,b}$ satisfy many remarkable properties related to their natural occurence as invariants of elements of $\mathrm{SL}(2, \mathbb{C})$.

We use the following lemma concerning $\kappa_{a,b}$ and $\mathbb{R}$-points extensively in later sections. Its proof is straightforward.



FIGURE 1. A triply-punctured sphere and its three boundary components

Lemma 2.3.1. *Fix $a, b \in \mathbb{R}$. Then $\kappa_{a,b}$ has:*

- *One real double root if either of a or b is ± 2;*
- *No real roots if $|a| > 2$ and $|b| < 2$ or vice versa;*
- *Two real roots, both greater than or equal to 2, if $|a|, |b| > 2$ and $ab > 0$;*
- *Two real roots, both less than or equal to -2, if $|a|, |b| > 2$ and $ab < 0$;*
- *Two real roots, both in $[-2, 2]$, if $|a|, |b| < 2$.*

This lemma illustrates the properties of reducible characters, since x is a root of $\kappa_{a,b}$ if and only if (a, b, x) is a reducible character. Indeed, we may assume that (a, b, x) is the character of a representation with abelian image. For example, the second case occurs when a is the trace of a hyperbolic element and b is the trace of an elliptic element, and in $\mathrm{SL}(2, \mathbb{R})$ hyperbolic elements and elliptic elements cannot commute. The other cases can be discussed similarly.

An important property of κ is that its restriction to every coordinate plane and coordinate line is quadratic. In particular its values at ± 2 are very special.

Lemma 2.3.2. $\kappa_{a,b}(\pm 2) = (a \pm b)^2$.

The discriminant of the quadratic polynomial $\kappa_{a,b}(x)$ equals

$$(a^2 - 4)(b^2 - 4).$$

Furthermore the critical points of $\kappa : \mathbb{C}^3 \longrightarrow \mathbb{C}$ are the four points

$$(2, 2, 2), \qquad (2, -2, -2), \qquad (-2, 2, -2), \qquad (-2, -2, 2)$$

with critical value $+2$ and the point $(0, 0, 0)$ with critical value -2.

2.4. **The quadruply-punctured sphere.** Let M be a quadruply-punctured sphere with boundary components A, B, C, D. Denoting the corresponding elements of $\pi = \pi_1(M)$ by the same name, the fundamental group $\pi = \pi_1(M)$ has the redundant geometric presentation

$$\pi = \langle A, B, C, D \mid ABCD = I \rangle$$

(Compare Figure 2.) However, in the present context there is an even more concrete description. A representation $\rho : \pi \longrightarrow \mathrm{SL}(2, \mathbb{C})$ corresponds to a quadruple

$$\alpha = \rho(A), \quad \beta = \rho(B),$$
$$\gamma = \rho(C), \quad \delta = \rho(D)$$

in $\mathrm{SL}(2, \mathbb{C})^4$ where

$$\alpha\beta\gamma\delta = I,$$



FIGURE 2. A quadruply-punctured sphere

Using this equation to eliminate δ, such a quadruple merely refers to an arbitrary triple $(\alpha, \beta, \gamma) \in \mathrm{SL}(2, \mathbb{C})^3$.

2.5. **Amalgamating representations.** Let ξ denote the simple loop on M which separates boundary components A, B from C, D. The corresponding element of the fundamental group satisfies:

$$X = AB = (CD)^{-1}.$$

Denote by $M|\xi$ the compact surface obtained by *splitting* M along ξ: each of the two components M_1 and M_2 of $M|\xi$ are homeomorphic to a triply-punctured sphere. We express the character variety $\mathfrak{X}(M)$ in terms of the character variety $\mathfrak{X}(M|\xi)$.

The fundamental group of M_1 has presentation

$$\pi_1(M_1) = \langle A, B, \bar{X} \mid AB\bar{X} = I \rangle$$

and the fundamental group of M_2 has presentation

$$\pi_1(M_2) = \langle X, C, D \mid XCD = I \rangle.$$

The fundamental group of M is the amalgamated free product

$$\pi_1(M) = \pi_1(M_1) \coprod_{\mathbb{Z}} \pi_1(M_2)$$

amalgamated by the monomorphisms taking a generator of $\mathbb{Z}$ to $\bar{X}^{-1} \in \pi_1(M_1)$ and $X \in \pi_1(M_2)$ respectively. A representation

$$\rho : \pi_1(M) \longrightarrow \mathrm{SL}(2, \mathbb{C})$$

restricts to two representations

$$\rho_i : \pi_1(M_i) \longrightarrow \mathrm{SL}(2, \mathbb{C})$$

satisfying

$$\rho_1(\bar{X})\rho_2(X) = I. \tag{2.5.1}$$

Conversely two representations ρ_1, ρ_2 satisfying (2.5.1) define a unique representation ρ. (Compare Goldman [5].)

2.6. **Defining equations.** The boundary map $\partial_* : \mathfrak{X}(M) \longrightarrow \mathbb{C}^4$ is defined by four functions

$$a([\rho]) = \mathrm{tr}(\rho(A)), \quad b([\rho]) = \mathrm{tr}(\rho(B)),$$
$$c([\rho]) = \mathrm{tr}(\rho(C)), \quad d([\rho]) = \mathrm{tr}(\rho(D))$$

The elements $X = AB$, $Y = BC$ and $Z = CA$ of π are represented by simple loops on M. The corresponding traces x, y, z, together with the boundary traces a, b, c, d, define coordinates on $\mathfrak{X}(M)$ which are subject to the relation

$$\begin{aligned} x^2 + y^2 + z^2 + xyz = (ab + cd)x \\ + (ad + bc)y + (ac + bd)z \\ + (4 - a^2 - b^2 - c^2 - d^2 - abcd) \end{aligned} \tag{2.6.1}$$

(Compare Magnus [9].) Unlike the case in which π is free of rank two, $\mathfrak{X}(M)$ is not an affine space, but rather the hypersurface $V \subset \mathbb{C}^7$ consisting of points whose coordinates (a, b, c, d, x, y, z) satisfy (2.6.1). For each $(a, b, c, d) \in \mathbb{C}^4$, the set

$$\partial_*^{-1}(a, b, c, d) = V_{a,b,c,d}$$

consists of all $(x, y, z) \in \mathbb{C}^3$ satisfying (2.6.1). Rewrite the basic defining equation (2.6.1) in terms of x as follows:

$$\begin{aligned} \frac{2+x}{4}\left((y+z) - \frac{(a+b)(d+c)}{2+x}\right)^2 \\ + \frac{2-x}{4}\left((y-z) - \frac{(a-b)(d-c)}{2-x}\right)^2 \\ = \frac{(x^2 - abx + a^2 + b^2 - 4)(x^2 - cdx + c^2 + d^2 - 4)}{4 - x^2} \end{aligned} \tag{2.6.2}$$

We may write (2.6.2) as follows. For fixed a, b, c, d, x,

$$\begin{aligned} Q_x^{(a,b,c,d)}(y, z) = \\ \frac{2+x}{4}\left((y+z) - \frac{(a+b)(d+c)}{2+x}\right)^2 \\ + \frac{2-x}{4}\left((y-z) - \frac{(a-b)(d-c)}{2-x}\right)^2 \end{aligned}$$

is an inhomogeneous quadratic function of (y, z). Equation (2.6.2) is equivalent to:

$$Q_x^{(a,b,c,d)}(y, z) = \frac{\kappa_{a,b}(x)\kappa_{c,d}(x)}{4 - x^2}$$

which describes a conic $V_{a,b,c,d}(x)$ centered at (y_0, z_0) where

$$y_0 + z_0 = \frac{(a+b)(d+c)}{2+x}$$
$$y_0 - z_0 = \frac{(a-b)(d-c)}{2-x}$$

2.7. **Symmetries.** A group of 192 symmetries of the family $\mathfrak{X}(M)$ acts by simple linear transformations on the parameters (a, b, c, d). This action is generated by an action of the *central characters,* which multiply an even number of the generators by $-I$, and the *permutations* of the four generators A, B, C, D.

The symmetric group S_4 consisting of permutations of $\{A, B, C, D\}$ acts on the family $\mathfrak{X}(M)$ of relative character varieties $V_{a,b,c,d}$. The symmetries are apparent from the defining equation (2.6.1). For example the three products of disjoint transpositions act trivially on the (x, y, z) coordinates, while nontrivially permuting (a, b, c, d). However, the transpositions $A \leftrightarrow B$, $B \leftrightarrow C$, $C \leftrightarrow D$, act as follows:

$$\begin{bmatrix} a \\ b \\ c \\ d \\ x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} b \\ a \\ c \\ d \\ x \\ z \\ y \end{bmatrix}, \qquad \begin{bmatrix} a \\ b \\ c \\ d \\ x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} a \\ c \\ b \\ d \\ z \\ y \\ x \end{bmatrix}, \qquad \begin{bmatrix} a \\ b \\ c \\ d \\ x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} a \\ b \\ d \\ c \\ x \\ z \\ y \end{bmatrix}$$

Furthermore every character variety admits an action by *central characters,* as described in Lubotzky-Magid [8], §5 (where this operation is called *twisting* by a character.) If $Z \subset G$ is central, then $\mathrm{Hom}(\pi, Z)$ is a *group,* which acts on $\mathrm{Hom}(\pi, G)$ by pointwise multiplication. Since the center of $G = \mathrm{SL}(2, \mathbb{C})$ equals $\{\pm I\} \cong \mathbb{Z}/2$ and the fundamental group π of a quadruply-punctured sphere is free of rank 3, then

$$\mathrm{Hom}(\pi, \{\pm I\}) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2$$

acts on $V_{a,b,c,d}$ as follows. Such a central character maps the images of an even number of the generators A, B, C, D to their negatives: for example,

$$\begin{bmatrix} \alpha \\ \beta \\ \gamma \\ \delta \end{bmatrix} \mapsto \begin{bmatrix} -\alpha \\ -\beta \\ \gamma \\ \delta \end{bmatrix}$$

is the character corresponding to

$$(1,1,0) \in \mathbb{Z}/2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2$$

and whose action on the trace coordinates is given by:

$$\begin{bmatrix} a \\ b \\ c \\ d \\ x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} -a \\ -b \\ c \\ d \\ x \\ -y \\ -z \end{bmatrix}$$

S_4 normalizes this $\mathrm{Hom}(\pi, \{\pm I\})$ - action. Since the induced S_4-action on $\mathrm{Hom}(\pi, \{\pm I\})$ is transitive, the action of every central character is obtained by conjugating the above example by a permutation. These considerations of symmetry immediately imply the corresponding relative character varieties are isomorphic:

Lemma 2.7.1. *Let $(a,b,c,d) \in \mathbb{R}^4$. Let $\sigma : \mathbb{R}^4 \to \mathbb{R}^4$ be any permutation of coordinates, and let $\tau : \mathbb{R}^4 \to \mathbb{R}^4$ be any map which acts by reversing signs of two of the coordinates and leaving the other two coordinates unchanged. Then $V_{\sigma(a,b,c,d)}$ and $V_{\tau(a,b,c,d)}$ are both diffeomorphic to $V_{a,b,c,d}$.*

3. Singular points and reducible characters

3.1. A criterion for singularity.

Theorem 3.1.1. *$V_{a,b,c,d}$ is singular if and only if at least one of the following holds:*

- *One of a,b,c,d equals ± 2;*
- *Some reducible representation has boundary traces (a,b,c,d).*

The latter case occurs if and only if there exists $x \in \mathbb{C}$ such that $\kappa_{a,b}(x) = \kappa_{c,d}(x) = 0$.

3.2. Solutions of quadratic equations.

We will need the following technical lemmas on discriminant loci for the proof of Theorem 3.1.1 and for later results.

Lemma 3.2.1. *Let $n \geq 1$ be an integer, and let $\mathbf{k}$ be a field of characteristic not equal to 2. Pick $B(x), C(x) \in \mathbf{k}[x_1, \ldots, x_n]$. Let*

$$Y = \{(x_1, \ldots, x_n, z) \in \mathbf{k}^{n+1} \mid z^2 + B(x)z + C(x) = 0\}$$

and

$$Z = \{(x_1, \ldots, x_n) \in \mathbf{k}^n \mid B(x)^2 - 4C(x) = 0\}.$$

If Y is singular at $(x_1, \ldots, x_n, z)$, then Z is singular at $(x_1, \ldots, x_n)$. If Z is singular at $(x_1, \ldots, x_n)$, then there exists $z \in \mathbf{k}$ such that Y is singular at $(x_1, \ldots, x_n, z)$.

(If $n = 1$, then Z is singular at x if x is a multiple root of $B(x)^2 - 4C(x)$.)

Proof. Denote by $dB(x)$ the differential of $B(x)$ with respect to the x_i-variables *only*, and similarly for $dC(x)$. Let $(x_1, \ldots, x_n) \in \mathbf{k}^n$. Now Z is singular at $(x_1, \ldots, x_n)$ if and only if

$$\begin{cases} B(x)^2 - 4C(x) & = 0 \\ 2B(x)\, dB(x) - 4\, dC(x) & = 0 \end{cases} \tag{3.2.1}$$

and Y is singular at $(x_1, \ldots, x_n, z)$ if and only if

$$\begin{cases} z^2 + B(x)z + C(x) & = 0 \\ 2z + B(x) & = 0 \\ z\, dB(x) + dC(x) & = 0. \end{cases} \tag{3.2.2}$$

We must show that (3.2.1) is equivalent to the existence of $z \in \mathbf{k}$ satisfying (3.2.2). Note that

$$B(x)^2 - 4C(x) = 0$$

if and only if

$$z^2 + B(x)z + C(x)$$

has a double root $z \in \mathbf{k}$; necessarily this root must be

$$z = -\frac{1}{2}B(x).$$

Substituting $z = -\frac{1}{2}B(x)$ into the final equation of (3.2.2) and multiplying by -4 yields the second equation of (3.2.1). □

Lemma 3.2.2. *Let $\mathbf{k}$ be an algebraically closed field of characteristic not equal to 2. Suppose $A_1(x), B_1(x), C_1(x) \in \mathbf{k}[x]$ such that $A_1(x)$ has no multiple roots. Let*

$$Y = \{(x, y) \in \mathbf{k}^2 \mid A_1(x)y^2 + B_1(x)y + C_1(x) = 0\}$$

and

$$Z = \{x \in \mathbf{k} \mid B_1(x)^2 - 4A_1(x)C_1(x) = 0\}.$$

If Y is singular at (x, y), then Z is singular at x. If Z is singular at x, then there exists $y \in k$ such that Y is singular at (x, y).

Proof. The proof follows that of Lemma 3.2.1, modified in the case that $A_1(x) = 0$. Suppose that $A_1(x) = 0$. Then Z is singular at x if and only if

$$\begin{cases} B_1(x)^2 & = 0 \\ 2B_1(x)B_1'(x) - 4C_1(x)A_1'(x) & = 0 \end{cases} \tag{3.2.3}$$

and Y is singular at (x, y) if and only if there exists $y \in \mathbf{k}$ such that

$$\begin{cases} yB_1(x) + C_1(x) & = 0 \\ B_1(x) & = 0 \\ y^2A_1'(x) + yB_1'(x) + C_1'(x) & = 0 \end{cases} \tag{3.2.4}$$

Clearly (3.2.3) is equivalent to $B_1(x) = C_1(x) = 0$, which (3.2.4) implies. Since $A_1'(x) \neq 0$ (by hypothesis), and $\mathbf{k}$ is algebraically closed, some $z \in \mathbf{k}$ satisfies the last equation in (3.2.4), concluding the proof of Lemma 3.2.2. □

Lemma 3.2.3. *Let* $\mathbf{k} = \mathbb{R}$. *Suppose* $A_1(x), B_1(x), C_1(x) \in \mathbb{R}[x]$ *such that* $A_1(x)$ *has no multiple roots. Let*

$$Y = \{(x, y) \in \mathbb{R}^2 \mid A_1(x)y^2 + B_1(x)y + C_1(x) = 0\}$$
$$Z = \{x \in \mathbb{R} \mid B_1(x)^2 - 4A_1(x)C_1(x) = 0\}.$$

and let

$$\delta(x) = B_1'(x)^2 - 4A_1'(x)C_1'(x). \tag{3.2.5}$$

If Y *is singular at* (x, y), *then* Z *is singular at* x. *Suppose for some* $x \in \mathbb{R}$ *that*

$$A_1(x) = B_1(x) = C_1(x) = 0$$

and

$$\delta(x) \geq 0$$

If Z *is singular at* x, *then there exists* $y \in \mathbb{R}$ *such that* Y *is singular at* (x, y).

Proof. The only point at which the proof of Lemma 3.2.2 fails for $\mathbf{k} = \mathbb{R}$ is the final step, where the last equation in (3.2.4) may not have a solution $y \in \mathbb{R}$. Since the discriminant of this quadratic equation equals $\delta(x)$, our assumption $\delta(x) \geq 0$ guarantees the existence of a solution $y \in \mathbb{R}$. □

3.3. **Singularities of $V_{a,b,c,d}$.**

Proof of Theorem 3.1.1. Fix $(a,b,c,d) \in \mathbb{C}^4$, and rewrite (2.6.1) as:

$$z^2 + B(x,y)z + C(x,y) = 0$$

where

$$\begin{aligned} B(x,y) &= xy - (ac+bd) \\ C(x,y) &= x^2 + y^2 - (ab+cd)x - (ad+bc)y \\ &\qquad - (4 - a^2 - b^2 - c^2 - d^2 - abcd). \end{aligned} \tag{3.3.1}$$

By Lemma 3.2.1, $V_{a,b,c,d}$ is singular if and only if the curve $\mathcal{C}$ in the xy-plane defined by the discriminant

$$\begin{aligned} f(x,y) &= B(x,y)^2 - 4C(x,y) \\ &= x^2y^2 - 4x^2 - 4y^2 \\ &\qquad - 2(ac+bd)xy + 4(ab+cd)x + 4(ad+bc)y \\ &\qquad + a^2c^2 + b^2d^2 - 2abcd \\ &\qquad - 4a^2 - 4b^2 - 4c^2 - 4d^2 + 16 = 0 \end{aligned} \tag{3.3.2}$$

is singular. Now write $f(x,y)$ as a quadratic function in y:

$$f(x,y) = A_1(x)y^2 + B_1(x)y + C_1(x) \tag{3.3.3}$$

where

$$\begin{aligned} A_1(x) &= (x^2 - 4) \\ B_1(x) &= 4(ad+bc) - 2(ac+bd)x \\ C_1(x) &= -4x^2 + 4(ab+cd)x + \\ &\qquad a^2c^2 + b^2d^2 - 2abcd \\ &\qquad - 4a^2 - 4b^2 - 4c^2 - 4d^2 + 16 = 0. \end{aligned} \tag{3.3.4}$$

Since $A_1(x) = (x-2)(x+2)$ has simple zeroes, Lemma 3.2.2 implies that $V_{a,b,c,d}$ is singular if and only if the discriminant $g(x)$ of $f(x,y)$ as a quadratic function of y vanishes. This discriminant is easily computed to be:

$$\begin{aligned} g(x) &= B_1(x)^2 - 4A_1(x)C_1(x) \\ &= 16(x^2 - abx + a^2 + b^2 - 4)(x^2 - cdx + c^2 + d^2 - 4) \\ &= 16(\kappa_{a,b}(x))(\kappa_{c,d}(x)). \end{aligned} \tag{3.3.5}$$

Thus $V_{a,b,c,d}$ is singular if and only if there exists $x \in \mathbf{k}$ satisfying one of the following three conditions:

1. x is a double zero of $\kappa_{a,b}$;

2. x is a double zero of $\kappa_{c,d}$;
3. $\kappa_{a,b}(x) = \kappa_{c,d}(x) = 0$.

Now the first case occurs if and only if the discriminant of $\kappa_{a,b}$ is zero, that is, if and only if

$$(a^2 - 4)(b^2 - 4) = 0,$$

that is, if and only if $a = \pm 2$ or $b = \pm 2$. Similarly, the second case occurs if and only if $c = \pm 2$ or $d = \pm 2$.

Consider the remaining case

$$\kappa_{a,b}(x) = \kappa_{c,d}(x) = 0 \tag{3.3.6}$$

We must find a reducible representation having traces $(a, b, c, d; x)$.

Decompose M along ξ as above into two triply-punctured spheres M_1 and M_2. (Compare §2.5.) Then there exist representations

$$\rho_i : \pi_1(M_i) \longrightarrow \mathrm{SL}(2, \mathbb{C})$$

having characters (a, b, x) and (x, c, d) respectively. By (3.3.6), both ρ_1 and ρ_2 are reducible.

We distinguish two cases, depending on whether $x = \pm 2$ or not. If $x = \pm 2$, then choose representations ρ_1, ρ_2 so that $\rho_1(\bar{X}) \neq \pm I$ and $\rho_2(X) \neq \pm I$. Necessarily $\rho_1(\bar{X})$ and $\rho_2(X)$ have unique fixed points p_1 and p_2, respectively, in $\mathbb{CP}^1$. There exists $g \in G$ such that

$$g(p_1) = p_2, \qquad g\rho_1(\bar{X}^{-1})g^{-1} = \rho_2(X). \tag{3.3.7}$$

Define ρ by:

$$\begin{aligned} &\rho(A) = \rho_1(A) && \rho(B) = \rho_1(B) \\ &\rho(C) = g\rho_2(C)g^{-1} && \rho(D) = g\rho_1(D)g^{-1} \end{aligned} \tag{3.3.8}$$

Since the image of ρ fixes p_1, it is reducible and its character lies in $V_{a,b,c,d}$. (Alternatively, one can choose

$$\rho(\bar{X}) = \pm I$$

and $\rho(C), \rho(D)$ to be diagonal matrices.)

Now suppose $x \neq \pm 2$. Then $\rho_1(\bar{X}^{-1})$ and $\rho_2(X)$ are elements of G having trace x and are thus conjugate. Indeed, they are diagonalizable with with distinct eigenvalues λ, λ^{-1} where $x = \lambda + \lambda^{-1}$. Let $p_1 \in \mathbb{CP}^1$ be the fixed point of $\rho_1(\bar{X}^{-1})$ corresponding to the λ-eigenspace of $\rho_1(\bar{X}^{-1})$ and $p_2 \in \mathbb{CP}^1$ be the fixed point of $\rho_2(X^{-1}))$ corresponding to the λ-eigenspace of $\rho_2(X^{-1}))$. Then as above there exists g satisfying (3.3.7). The representation ρ defined by (3.3.8) fixes p_1 and its character lies in $V_{a,b,c,d}$. □

3.4. **A discriminant polynomial.** Given $(a,b,c,d) \in \mathbb{C}^4$, the existence of an $x \in \mathbb{C}$ satisfying condition (3.3.6) is equivalent to the vanishing of the *resultant* $\Delta(a,b,c,d)$ of the polynomials $\kappa_{a,b}$ and $\kappa_{c,d}$, which (remarkably) is a symmetric polynomial in a,b,c,d. This resultant is the sextic polynomial

$$\begin{aligned}
\Delta(a,b,c,d) &= \begin{vmatrix} 1 & -ab & a^2+b^2-4 & 0 \\ 0 & 1 & -ab & a^2+b^2-4 \\ 1 & -cd & c^2+d^2-4 & 0 \\ 0 & 1 & -cd & c^2+d^2-4 \end{vmatrix} \\
&= a^2b^2c^2 + a^2b^2d^2 + a^2c^2d^2 + b^2c^2d^2 \\
&\quad - a^3bcd - ab^3cd - abc^3d - abcd^3 \\
&\quad + a^4 + b^4 + c^4 + d^4 + 8abcd \\
&\quad - 2(a^2b^2 + a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2 + c^2d^2) \\
&= \big(2(a^2+b^2+c^2+d^2) - abcd - 16\big)^2 \\
&\quad - (4-a^2)(4-b^2)(4-c^2)(4-d^2).
\end{aligned} \tag{3.4.1}$$

Note that the vanishing of $\Delta(a,b,c,d)$ implies only that at least one character in $V_{a,b,c,d}$ is reducible. In general, a character

$$(a,b,c,d,x,y,z) \in V_{a,b,c,d}$$

is reducible if and only if

$$\begin{aligned}
\kappa_{a,b}(x) = \kappa_{c,d}(x) = \\
\kappa_{a,d}(y) = \kappa_{b,c}(y) = \\
\kappa_{a,c}(z) = \kappa_{b,d}(z) = 0
\end{aligned}$$

4. Topology of the set of $\mathbb{R}$-points

4.1. **Singular points.** For $(a,b,c,d) \in \mathbb{R}^4$, the variety $V_{a,b,c,d}$ is defined over $\mathbb{R}$; from now on we will only consider its set $V_{a,b,c,d}(\mathbb{R})$ of $\mathbb{R}$-points. By Prop. III.1.1 of Morgan-Shalen [11] (p.458), $V_{a,b,c,d}(\mathbb{R})$ consists of characters corresponding to representations in either SU(2) or $\mathrm{SL}(2,\mathbb{R})$. Since the leading term of the cubic polynomial defining $V_{a,b,c,d}$ is xyz, its closure in $\mathbb{RP}^3$ meets the plane at infinity in three lines in general position (the singular cubic defined by $xyz = 0$ in $\mathbb{RP}^2$). Thus for any values of $(a,b,c,d) \in \mathbb{R}^4$, the surface $V_{a,b,c,d}(\mathbb{R})$ has four ends. (Compare Figure 3.)



FIGURE 3. What $V_{a,b,c,d}(\mathbb{R})$ looks like near infinity

Theorem 4.1.1. *Let $(a,b,c,d) \in \mathbb{R}^4$. Then $V_{a,b,c,d}(\mathbb{R})$ contains a singular point if and only if one of the following holds:*

1. *One of a,b,c,d equals ± 2;*
2. *There is a reducible $\mathrm{SU}(2)$-representation having boundary traces (a,b,c,d), and $a,b,c,d \in [-2,2]$;*
3. *There is a reducible $\mathrm{SL}(2,\mathbb{R})$-representation having boundary traces (a,b,c,d), and $a,b,c,d \in \mathbb{R} \setminus (-2,2)$.*

The second condition is equivalent to

$$\Delta(a,b,c,d) = 0 \text{ and } a,b,c,d \in [-2,2];$$

the third condition is equivalent to

$$\Delta(a,b,c,d) = 0 \text{ and } a,b,c,d \in \mathbb{R} \setminus (-2,2).$$

We first verify the final statements. Suppose $x \in \mathbb{R}$ satisfies

$$\kappa_{a,b}(x) = \kappa_{c,d}(x) = 0$$

and $a,b,c,d \in [-2,2]$. By §3.4, $\Delta(a,b,c,d) = 0$. Conversely, given $\Delta(a,b,c,d) = 0$ with $a,b,c,d \in [-2,2]$, there exists a reducible $\mathrm{SL}(2,\mathbb{C})$-representation ρ with the given boundary traces. Furthermore,

$$x = \mathrm{tr}(\rho(X)) \in [-2,2]$$

by Lemma 2.3.1. By §2.2, the triples (a, b, x) and (c, d, x) correspond to SU(2)-representations. Amalgamating the representations as in the proof of Theorem 3.1.1 produces a SU(2)-representation ρ of $\pi_1(M)$. The equivalence of the third condition is proved similarly.

The proof of the rest of Theorem 4.1.1 is analogous to the proof of Theorem 3.1.1. The only difference is that we must use Lemma 3.2.3 instead of Lemma 3.2.2. Thus, it suffices to show:

Lemma 4.1.2. *Let $x, a, b, c, d \in \mathbb{R}$ and define $A_1(x), B_1(x), C_1(x)$ by (3.3.4). Then $\delta(x) \geq 0$ where $\delta(x)$ is defined in (3.2.5).*

Proof. By (3.3.4),

$$\begin{aligned} A_1'(x) &= 2x \\ B_1'(x) &= -2(ac + bd) \\ C_1'(x) &= -8x + 4(ab + cd) \end{aligned}$$

and

$$\begin{aligned} \delta(x) &= B_1'(x)^2 - 4A_1'(x)C_1'(x) \\ &= 4(ac + bd)^2 + 64x^2 - 32x(ab + cd) \\ &= 4\left(4x - (ac + bd)\right)^2 \geq 0. \end{aligned}$$

□

4.2. **Topological classification of the smooth real varieties.** Assume that $V_{a,b,c,d}(\mathbb{R})$ is nonsingular. We now classify the possible topologies for $V_{a,b,c,d}(\mathbb{R})$. The six topological types are depicted in Figures 11, 12, 13, 14, 15, 16.

Theorem 4.2.1. *If $(a, b, c, d) \in \mathbb{R}^4$ and $V_{a,b,c,d}(\mathbb{R})$ is nonsingular, then $V_{a,b,c,d}(\mathbb{R})$ is diffeomorphic to one of the following:*

1. *A quadruply-punctured sphere, in the case $a, b, c, d \in \mathbb{R} \setminus [-2, 2]$ with $abcd < 0$;*
2. *The disjoint union of a triply-punctured torus and a disk, in the case $a, b, c, d \in \mathbb{R} \setminus [-2, 2]$ with $abcd > 0$;*
3. *The disjoint union of a triply-punctured sphere and a disk, in the case that one of a, b, c, d is in $(-2, 2)$ and the rest are in $\mathbb{R} \setminus [-2, 2]$;*
4. *The disjoint union of an annulus and two disks, in the case that two of a, b, c, d are in $(-2, 2)$ and two are in $\mathbb{R} \setminus [-2, 2]$;*
5. *The disjoint union of four disks, in the case that three of a, b, c, d are in $(-2, 2)$ and the other is in $\mathbb{R} \setminus [-2, 2]$;*
6. *The disjoint union of four disks and a sphere, in the case that $a, b, c, d \in (-2, 2)$*

In the second case, the component of $V_{a,b,c,d}$ homeomorphic to a disc is the "Teichmüller space" of M, consisting of hyperbolic structures with geodesic boundary on M. Compare Goldman [6] and [5].

4.3. **Reduction to a plane quartic curve.** To prove Theorem 4.2.1, reduce the dimension by taking discriminants, as in the determination of the singularities. Again, difficulties arise because the discriminant $f(x,y)$ is not monic in y. The set $V_{a,b,c,d}(\mathbb{R})$ contains two points for every $(x,y) \in \mathbb{R}^2$ with $f(x,y) > 0$, one point for $f(x,y) = 0$, and none for $f(x,y) < 0$. Define

$$\mathcal{S} = \{(x,y) \in \mathbb{R}^2 : f(x,y) > 0\}$$

and

$$\mathcal{C} = \{(x,y) \in \mathbb{R}^2 : f(x,y) = 0\} = \partial\mathcal{S}.$$

The projection

$$(x,y) : V_{a,b,c,d}(\mathbb{R}) \longrightarrow \mathbb{R}^2$$

expresses $V_{a,b,c,d}(\mathbb{R})$ as a union of two copies of $\mathcal{S}$ identified along their boundary $\mathcal{C}$. This description as a "double branched cover" determines $V_{a,b,c,d}(\mathbb{R})$ up to homeomorphism (and hence diffeomorphism).

Fixing $x \in \mathbb{R}$ with $x \neq \pm 2$, for $|y|$ large,

$$f(x,y) \sim (x^2 - 4)y^2$$

and so $f(x,y)$ is positive if $|x| > 2$ and negative if $|x| < 2$. More precisely,

- for any x with $|x| < 2$, there exists $M > 0$ such that for $|y| > M$, $(x,y) \notin \mathcal{S}$;
- for any x with $|x| > 2$, there exists $M > 0$ such that for $|y| > M$, $(x,y) \in \mathcal{S}$.

To study $\mathcal{C}$, we reduce dimensions once more via discriminants. For fixed $x \neq \pm 2$, the tangent of $\mathcal{C}$ at (x,y) for some $y \in \mathbb{R}$ is vertical if and only if the discriminant

$$g(x) = 16\kappa_{a,b}(x)\kappa_{c,d}(x) = 0.$$

Lemma 4.3.1. *Let $x \in \mathbb{R} \setminus \{\pm 2\}$. Then the set of $y \in \mathbb{R}$ such that $(x,y) \in \mathcal{C}$ consists of:*

- *Exactly two points if $g(x) > 0$;*
- *Exactly one point if $g(x) = 0$.*
- *No points if $g(x) < 0$.*

(Recall the definition of $A_1(x), B_1(x), C_1(x)$ from (3.3.4).)

Lemma 4.3.2. *Suppose $x = \pm 2$ and $g(x) \neq 0$. Then $B_1(x) \neq 0$, and there exists exactly one $y \in \mathbb{R}$ such that $(x, y) \in \mathcal{C}$.*

Lemma 4.3.3. *Suppose that $x = \pm 2$ and $g(x) = 0$. Then $B_1(x) = 0$, and the set of $y \in \mathbb{R}$ such that $(x, y) \in \mathcal{C}$ is empty, if $C_1(x) \neq 0$ or $\mathbb{R}$, if $C_1(x) = 0$.*

Proofs. We write $f(x, y)$ as

$$f(x, y) = A_1(x)y^2 + B_1(x)y + C_1(x)$$

Lemma 4.3.1 follows because $f(x, y)$ is a quadratic polynomial in y with discriminant $g(x)$. Lemma 4.3.2 follows because $A_1(x) = 0$ and

$$g(x) = B_1(x)^2 - 4A_1(x)C_1(x).$$

Finally, Lemma 4.3.3 follows because its hypotheses imply $f(x, y) = C_1(x)$. □

Lemma 4.3.4. *Suppose that $a, b, c, d \in \mathbb{R} \setminus \{2, -2\}$.*

- *If $B_1(2) = C_1(2) = 0$, then $a = b$ and $c = d$.*
- *If $B_1(-2) = C_1(-2) = 0$, then $a = -b$ and $c = -d$.*

Furthermore, in either of these cases, $V_{a,b,c,d}(\mathbb{R})$ is singular if and only if a, b, c, and d are either all in $(-2, 2)$ or all in $\mathbb{R} \setminus [-2, 2]$.

Proof. $B_1(\pm 2) = 0$ factors as $(b \mp a)(c \mp d) = 0$, so without loss, $a = \pm b$. Substituting, $C_1(\pm 2) = 0$ becomes $(b^2 - 4)(c \mp d)^2 = 0$. Since $b \neq \pm 2$, we have $c = \pm d$. One calculates $\Delta(\pm b, b, \pm d, d) = 0$. Now apply Theorem 4.1.1. □

Lemma 4.3.5. *Suppose $V_{a,b,c,d}(\mathbb{R})$ contains no singular points but $g(x)$ has a real double root. Then this root is ± 2, and $a = \pm b$ and $c = \pm d$. One of the two possibilities must occur:*

- *$a \in (-2, 2)$ and $c \in \mathbb{R} \setminus [-2, 2]$;*
- *$c \in (-2, 2)$ and $a \in \mathbb{R} \setminus [-2, 2]$.*

Furthermore $\mathcal{C}$ contains the line $x = \pm 2$.

Proof. By Theorem 4.1.1, this situation can only arise if g has a double root and yet some of a, b, c, d are in $(-2, 2)$ and some are in $\mathbb{R} \setminus [-2, 2]$. Note that for $\kappa_{a,b}(x)$ to have a double root, one of a or b must be ± 2, but this does not happen since $V_{a,b,c,d}(\mathbb{R})$ is nonsingular. Thus, $\kappa_{a,b}(x)$ and $\kappa_{c,d}(x)$ have a common root. If this root lies in $(-2, 2)$, then Lemma 2.3.1 implies that all of a, b, c, d lie in $(-2, 2)$. Similarly, if this root lies in $\mathbb{R} \setminus [-2, 2]$, then all of a, b, c, d lie in $\mathbb{R} \setminus [-2, 2]$. In either case, Theorem 4.1.1 implies that $V_{a,b,c,d}(\mathbb{R})$ is singular. Therefore, the root is ± 2, implying $a = \pm b$ and $c = \pm d$. In addition, $B_1(\pm 2) = C_1(\pm 2) = 0$, and so Lemma 4.3.3 implies that $\mathcal{C}$ contains the line $x = \pm 2$. □

We continue the analysis of $\mathcal{C}$. The solution of the quadratic equation $f(x,y)=0$ for y when $x \neq \pm 2$ is $y = \phi_\pm(x)$ where

$$\phi_\pm(x) = \frac{-B_1(x) \pm \sqrt{g(x)}}{2(x^2-4)}$$

Note that $B_1(x)$ is linear in x and $g(x) \sim 16x^4$; so as $|x| \to \infty$,

$$\phi_\pm(x) \to \pm 2.$$

Similarly, fix $y \neq \pm 2$ and define $x = \psi_\pm(y)$ by solving $f(x,y) = 0$. Then, as $|y| \to \infty$,

$$\psi_\pm(y) \to \pm 2.$$

Thus, for any $\varepsilon > 0$, there is $M > 2 + \varepsilon$ for which

$$|x| > M \Longrightarrow \begin{cases} |\phi_+(x) - 2| & < \varepsilon \\ |\phi_-(x) + 2| & < \varepsilon \end{cases}$$

and for which

$$|y| > M \Longrightarrow \begin{cases} |\psi_+(y) - 2| & < \varepsilon \\ |\psi_-(y) + 2| & < \varepsilon \end{cases}$$

Fix $0 < \varepsilon < 1$, and let Q be the square $(-M, M)^2$. Then $\mathcal{C} \cap (\mathbb{R}^2 \setminus Q)$ is homeomorphic to a disjoint union of eight copies of $[M, \infty)$. For example, one of the eight homeomorphisms would map $t \in [M, \infty)$ to $(-t, \phi_+(-t))$. Note that $\mathcal{C}$ intersects ∂Q at the images of M under the eight homeomorphisms. $\mathcal{C}$ is smooth and hence a 1-manifold, so $\overline{Q} \cap \mathcal{C}$ is a compact 1-manifold with boundary $\partial\overline{Q} \cap \mathcal{C}$ consisting of eight points.

Lemma 4.3.6. *Assume $V_{a,b,c,d}(\mathbb{R})$ is non-singular. With Q as above, $\overline{Q} \cap \mathcal{C}$ has finitely many connected components. Each component is homeomorphic to either a closed interval $[-1,1]$, if at least one (and hence exactly two) points of the component lie in ∂Q, or S^1 otherwise. Furthermore, there must be exactly four components homeomorphic to an interval.*

$\mathcal{C}$ must look something like Figure 4. Outside Q, all points of $\mathcal{C}$ are close to the eight asymptotes. By choosing Q sufficiently large, $\mathcal{C}$ will be transverse to the four sides of ∂Q and $\mathcal{C} \cap \partial Q$ is a set of eight points,

$$P = \{p_1, p_2, p_3, p_4, p_5, p_6, p_7, p_8\},$$

numbered as in Figure 4. To determine the topologies of $\mathcal{C}$ and $\mathcal{S}$, it suffices to decide how these eight points of $\mathcal{C} \cap \partial Q$ are joined pairwise inside Q, and to find the number and positions of any bounded components.



FIGURE 4. How the curve $\mathcal{C}$ meets the square Q

In each of the following lemmas, the set of $\mathbb{R}$-points is assumed to be nonsingular.

Lemma 4.3.7. *If p_n is joined by $\mathcal{C}$ to p_m, then $n+m$ is odd.*

Proof. Suppose $n+m$ is even. Then on ∂Q, the points p_n and p_m separate the remaining points p_i into two sets P_1 and P_2 of odd cardinality. Since an arc α of $\mathcal{C} \cap Q$ joins p_n and p_m, this arc separates P_1 from P_2 in $\bar{Q}$. Furthermore the points of

$$P - \{p_m, p_n\} = P_1 \coprod P_2$$

are pairwise joined by arcs of $\mathcal{C} \cap Q$ which (since $\mathcal{C}$ is smooth) do not meet α. Thus the points of P_1 join to points of P_1, and similarly for P_2. This partitions P_1 into disjoint pairs, contradicting the odd cardinality of P_1. □

Lemma 4.3.8.

- *p_3 joins to p_4 if and only if $g(x)$ has a root in $(2, \infty)$;*
- *p_7 joins to p_8 if and only if $g(x)$ has a root in $(-\infty, -2)$.*

Proof. We'll show the first statement; the second is similar. Suppose first that p_3 joins to p_4. The component joining p_3 and p_4 must have a vertical tangent — corresponding to a root of $g(x)$ — at some point x_0. If $x_0 < 2$, then the component crosses the line $x = 2$ twice, contradicting Lemma 4.3.2. (If the component crossed the line at the same point each time, there would be a singularity. Alternatively, if $g(2) = 0$, then by Lemma 4.3.3, $\mathcal{C}$ must contain the whole line $x = 2$, and we would have singularities where the component crosses.) If $x_0 = 2$, then, by Lemma 4.3.3, either $\mathcal{C}$ doesn't intersect the line $x = 2$, or else $\mathcal{C}$ contains it. But a vertical tangent at $x = 2$ would force at least one point in $\mathcal{C}$ on that line; and if $\mathcal{C}$ contained the line, a component from the right touching the line at a point would result in a singularity. Thus $x_0 > 2$. Since $\mathcal{C}$ has a vertical tangent there, $g(x_0) = 0$.

Conversely, let $x_0 > 2$ be the root in question. Without loss, x_0 is the largest root of g. Thus $g(s) < 0$ for some $s \in (2, x)$, since $g(x) > 0$ for $x > x_0$, and x_0 is not a double root. If p_3 were to join to p_1, p_6, p_7, or p_8, then clearly $\mathcal{C}$ would cross the line $x = s$, contradicting Lemma 4.3.1. It also can't join to p_2 or p_5, since in that case, $\mathcal{C}$ would cross $x = s$ somewhere along the approach to the asymptote at $x = 2$. Thus p_3 must join to p_4. □

Lemma 4.3.9.

- *p_1 joins to p_6 if and only if g has two roots in $(-\infty, -2]$ and two roots in $[-2, 2]$;*
- *p_2 joins to p_5 if and only if g has two roots in $[2, \infty)$ and two roots in $[-2, 2]$.*

Proof. We'll show the first statement; the second is similar. To prove the forward direction, suppose p_1 and p_6 are joined by a component. If this component is the line $x = -2$, then Lemmas 4.3.3 and 4.3.4 imply that $a = -b$ and $c = -d$. Thus g has a double root at -2. By Lemma 4.3.5, we can assume without loss that $a, b \in (-2, 2)$ and $c, d \in \mathbb{R} \setminus [-2, 2]$. Therefore, by Lemma 2.3.1, g has a root in $(-\infty, -2]$, a double root at -2, and a root in $(-2, 2]$, and we are done.

Now assume $\mathcal{C}$ does not contain the line $x = -2$. Consider the whole component of $\mathcal{C}$ that joins p_1 to p_6, and not just the part lying inside Q; let α be a diffeomorphism mapping $\mathbb{R}$ onto this component. If -2 is a root of g, Lemma 4.3.3 implies that $\mathcal{C}$ does not intersect the line $x = -2$. If the x-value of $\alpha(t)$ increases from -2 as t increases from $-\infty$, then at some point it must decrease back to -2 as t increases to $+\infty$. Thus, there must be a vertical tangent — and hence a root of g — at some x-value greater than -2. This root must be less than 2, for the same reasons as in Lemma 4.3.8. Meanwhile, p_7 and p_8 must join

to each other, since they have nowhere else to go. Lemma 4.3.8 implies that g has a root in $(-\infty, -2)$. We now know that g has at least three real roots; it therefore has a fourth. By Lemma 2.3.1, this fourth root is somewhere in $(-\infty, 2]$, and we have the desired conclusion. The argument is similar if the x-value of α decreases and then increases.

On the other hand, if -2 is not a root of g, Lemma 4.3.1 implies that $\mathcal{C}$ must cross the line $x = -2$ at exactly one point. If any component other than the one in question were to cross this line, then it must cross it at least twice (once to get across, and once to get back) or at a singularlity or vertical tangent. Thus, the component in question must cross $x = -2$ exactly once. The x-value of α must increase from -2, decrease back down to -2 and below, and then increase back up to -2; as a result, we get at least two vertical tangents, one on each side of -2. By the same argument as above, both roots must be less than 2. Lemma 2.3.1 implies g must also have two other roots, one on each side of -2, and neither larger than 2.

For the converse, suppose that -2 is a simple root of g, and $x_1 \in (-\infty, -2)$ and $x_2, x_3 \in (-2, 2]$ are also roots. Lemma 4.3.8 implies that p_7 and p_8 join. Furthermore, g is negative on (x_2, x_3), and so there are no points of $\mathcal{C}$ above that interval. Thus p_1 and p_6 are cut off from the other points, and must join to each other. The argument is similar if -2 is a root, with two roots less than -2 and one greater, or if there are two roots less than -2 and two greater. Finally, if -2 is a multiple root of g, then Lemma 4.3.5 implies $a = -b$ and $c = -d$. As a result, $B_1(-2) = C_1(-2) = 0$. Lemma 4.3.3 implies that $\mathcal{C}$ contains the line $x = -2$, and so p_1 and p_6 join. □

(Note that in the above lemma, the case of a double root at ± 2 is considered as a deformation of the case of one root just above and one just below.)

Lemma 4.3.10. *If $\mathcal{C}$ has a compact component, then g has four distinct real roots $x_1 < x_2 < x_3 < x_4$, and the projection of the compact component onto the x-axis is $[x_2, x_3]$.*

Proof. The projection of the component onto the x-axis must be $[x_i, x_j]$, where x_i and x_j are distinct roots of g. Furthermore, g must be strictly positive on (x_i, x_j), and negative for some $x < x_i$ and for some $x > x_j$. Thus, g must have four real roots, and x_i and x_j must be the middle two. □

Corollary 4.3.11. *There is at most one compact component of $\mathcal{C}$.*

Lemma 4.3.12. *If g has four real roots $x_1 < x_2 < x_3 < x_4$ and $2, -2 \notin [x_2, x_3]$, then $\mathcal{C}$ has a compact component projecting onto $[x_2, x_3]$.*

Proof. Since the leading coefficient of f is nonzero throughout $[x_2, x_3]$, and the discriminant g is zero at x_2 and x_3 and positive in between, the part of $\mathcal{C}$ that lies above the $[x_2, x_3]$ must consist of two line segments glued at the endpoints. As g is negative just below x_2 and just above x_3, it follows $\mathcal{C}$ has a compact component. □

4.4. **The six cases.** We now analyze individual cases to find the resulting topologies.

In the first case, we are given $a, b, c, d \in \mathbb{R} \setminus [-2, 2]$ and $abcd < 0$. By applying Lemma 2.7.1, we can assume that $a, b, c > 2$ and $d < -2$. Thus, by Lemma 2.3.1, g has two roots in $[2, +\infty)$ and two in $(-\infty, 2]$. Note that by Lemma 4.3.5, the roots of g are all distinct; in particular, there must be at least one root greater than 2. Now by Lemma 4.3.8, p_3 and p_4 join, and p_7 and p_8 join. p_1 and p_5 do not join, by Lemma 4.3.7. If p_1 and p_6 were to join, then by Lemma 4.3.9, all four roots would be in $(-\infty, 2]$, contradicting the existence of a root greater than 2. Thus, p_1 and p_2 join; finally, p_5 and p_6 must join. Furthermore, there cannot be a compact component, since by Lemma 4.3.10 such a component would have to project onto $[x_2, x_3]$, with $x_2 \leq -2$ and $x_3 \geq 2$; but such a component would cross the lines $x = \pm 2$ twice, contradicting Lemma 4.3.2 and Lemma 4.3.3. Thus, $\mathcal{C} \cap \overline{Q}$ consists of just the four above-mentioned components.

Since f is negative for $x = 0$ and y large, we see that $\mathcal{S}$ is homeomorphic to a closed disk minus four points of the boundary. (See Figure 5.) Doubling along the boundary, we see that $V_{a,b,c,d}(\mathbb{R})$ is homeomorphic, and hence diffeomorphic, to a quadruply-punctured sphere. (See Figure 11.)

In the second case, all $a, b, c, d \in \mathbb{R} \setminus [-2, 2]$ and $abcd > 0$. Applying Lemma 2.7.1, we may assume that $a, b, c, d > 2$. Lemma 2.3.1 implies that $g(x)$ has four roots in $[2, +\infty)$. Lemma 4.3.8 implies that p_3 and p_4 must join, and p_7 and p_8 cannot join. Lemma 4.3.9 implies that p_1 and p_6 cannot join, nor can p_2 and p_5. Thus p_1 joins either p_2 or p_8, by Lemma 4.3.7. If p_1 joins p_2, then p_7 must join p_6 and so p_8 joins p_5; or, if p_1 joins p_8, then p_5 joins p_6 and p_2 joins p_7. Meanwhile, Lemma 4.3.12 implies that $\mathcal{C}$ has a compact component somewhere between the component joining p_3 to p_4 and the line $x = 2$. Thus $\mathcal{C}$ consists of a circle, together with one of the two possible sets of four non-compact components; both setups result in the same topology for $\mathcal{S}$.

Since f is negative for $x = 0$ and y large, $\mathcal{S}$ is homeomorphic to a closed annulus minus three points of the outer boundary, along with a closed disk minus one point of the boundary. (See Figure 6.) Doubling



FIGURE 5. Case 1

again, $V_{a,b,c,d}(\mathbb{R})$ is homeomorphic, and hence diffeomorphic, to a triply-punctured torus and a disk. (See Figure 12.)

In the third case, three of $a, b, c, d \in \mathbb{R} \setminus [-2, 2]$ and the fourth in $(-2, 2)$. By applying Lemma 2.7.1, we can assume that $d \in (-2, 2)$ and $a, b, c > 2$. By Lemma 2.3.1, g has two roots in $[2, +\infty)$ and no other real roots. Thus, by Lemma 4.3.10, there cannot be a compact component. Furthermore, Lemma 4.3.8 and Lemma 4.3.9 imply that p_3 and p_4 must join, but p_7 and p_8 cannot join, p_1 and p_6 cannot join, and p_2 and p_5 cannot join. Thus, we end up with the same arrangement of non-compact components as in Case 2, but no compact component.

From the sign of f, we see that $\mathcal{S}$ is homeomorphic to a closed disk minus three points of the boundary, along with a closed disk minus one point of the boundary. (See Figure 7.) Doubling again, we see that $V_{a,b,c,d}(\mathbb{R})$ is homeomorphic, and hence diffeomorphic, to a triply-punctured sphere and a disk. (See Figure 13.)



FIGURE 6. Case 2

In the fourth case, two of $a, b, c, d \in \mathbb{R} \setminus [-2, 2]$ and two in $(-2, 2)$. Applying Lemma 2.7.1, we can assume that $a, b \in (-2, 2)$ and $c, d > 2$. There are two possibilities.

Suppose first that $g(x)$ has a double root. Lemma 4.3.5 implies this double root must be 2; there is also one root in $[-2, 2)$ and one in $(2, \infty)$. By Lemma 4.3.10, there is no compact component and Lemma 4.3.5 implies that p_2 and p_5 join by a vertical line. Lemma 4.3.8 implies that p_3 and p_4 join, but p_7 and p_8 do not. Thus p_1 and p_8 must join, as must p_6 and p_7.

The second possibility is that all of the roots of g are distinct; thus, two distinct roots lie in $[-2, 2]$ and two distinct roots lie in $[2, +\infty)$, and 2 is not a double root. Once again, Lemma 4.3.10 precludes a compact component, and Lemma 4.3.8 implies that p_3 and p_4 join. In addition, Lemma 4.3.9 implies that p_2 and p_5 join, while p_1 and p_6 do not. Thus, p_1 and p_8 join, as do p_6 and p_7.

Both possibilities give the same result: $\mathcal{S}$ is homeomorphic to a closed disk minus two points of the boundary, along with two closed



FIGURE 7. Case 3

disks missing one boundary point each. (See Figure 8.) Doubling again, $V_{a,b,c,d}(\mathbb{R})$ is homeomorphic, and hence diffeomorphic, to a disjoint union of an annulus and two disks. (See Figure 14.)

In the fifth case, one of a, b, c, d lie in $\mathbb{R} \setminus [-2, 2]$ and the other three in $(-2, 2)$. Applying Lemma 2.7.1, we can assume that $a, b, c \in (-2, 2)$ and $d > 2$. Lemma 2.3.1 implies g has two roots in $[-2, 2]$ and no other real roots. Lemma 4.3.10 precludes a compact component. Furthermore, Lemma 4.3.8 and Lemma 4.3.9 imply that p_3 can't join to p_4, p_7 can't join to p_8, p_1 can't join to p_6, and p_2 can't join to p_5. Furthermore, g is negative for some $x_0 \in (-2, 2)$; thus, none of the points on the left can join to any of the points on the right, by Lemma 4.3.1. Therefore p_1 joins to p_8, p_2 joins to p_3, p_4 joins to p_5, and p_6 joins to p_7.

From the sign of f, we see that $\mathcal{S}$ is homeomorphic to a union of four disjoint closed disks missing one boundary point each. (See Figure 9.) Doubling, we see that $V_{a,b,c,d}(\mathbb{R})$ is homeomorphic, and hence diffeomorphic, to a union of four disjoint disks. (See Figure 15.)



FIGURE 8. Case 4

In the sixth case, all of a, b, c, d lie in $(-2, 2)$. Lemma 2.3.1 implies that g has four roots in $[-2, 2]$. Lemma 4.3.12 implies that $\mathcal{C}$ has a compact component between the lines $x = -2$ and $x = 2$. The rest of the analysis is exactly as in Case 5; the result is that p_1 joins to p_8, p_2 joins to p_3, p_4 joins to p_5, and p_6 joins to p_7.

From the sign of f, we see that $\mathcal{S}$ is homeomorphic to a closed disk, along with four closed disks missing one boundary point each. (See Figure 10.) Doubling, we see that $V_{a,b,c,d}(\mathbb{R})$ is homeomorphic, and hence diffeomorphic, to four disks and a sphere. (See Figure 16.) This concludes our case-by-case analysis and the proof of Theorem 4.2.1.

5. A compact component of $\mathrm{SL}(2, \mathbb{R})$-representations

When $(a, b, c, d) \in \mathbb{R}$ and $V_{a,b,c,d}(\mathbb{R})$ is smooth, then each connected component of $V_{a,b,c,d}(\mathbb{R})$ consists of characters of representations in either $\mathrm{SL}(2, \mathbb{R})$ or $\mathrm{SU}(2)$. Components of spaces of representations in $\mathrm{SU}(2)$ are compact. Conversely, when M is a once-punctured torus,



FIGURE 9. Case 5

the only compact components of $\mathfrak{X}(M)(\mathbb{R})(t)$ are characters of SU(2)-representations. (See Goldman [6] for a description of the topology of the relative character varieties for a punctured torus.) However, for the quadruply punctured sphere, there are compact components of $V_{a,b,c,d}(\mathbb{R})$ for certain values of (a,b,c,d) which do not correspond to SU(2)-representations.

If $V_{a,b,c,d}(\mathbb{R})$ contains a compact component, then Theorem 4.2.1 implies that

$$-2 < a, b, c, d \ < 2,$$

and conversely, every such $(a,b,c,d) \in (-2,2)^4$ for which $V_{a,b,c,d}(\mathbb{R})$ is smooth contains a compact component. (Compare the analysis of Case 6.) Here is a criterion for when this compact component consists of $\mathrm{SL}(2,\mathbb{R})$-characters or SU(2)-characters, in terms of the function $\Delta(a,b,c,d)$ defined in §3.4.

Proposition 5.1. *Suppose $a,b,c,d \in (-2,2)$ and $V_{a,b,c,d}$ is smooth. Then the compact component of $V_{a,b,c,d}(\mathbb{R})$ consists of characters of*



FIGURE 10. Case 6

$\mathrm{SL}(2,\mathbb{R})$-*representations if and only if*

$$\Delta(a,b,c,d) > 0 \tag{5.0.1}$$

$$16 - abcd - 2(a^2+b^2+c^2+d^2) > 0. \tag{5.0.2}$$

Otherwise, each element in the component is the character of an $\mathrm{SU}(2)$*-representation.*

Proof. Decompose the quadruply-punctured sphere M along ξ, as in §2.5, as a union of two triply-punctured spheres. A given representation $\pi_1(M) \longrightarrow \mathrm{SL}(2,\mathbb{C})$ has image in $\mathrm{SU}(2)$ only if its restriction to the fundamental group of each component of $M|\xi$ has image in $\mathrm{SU}(2)$. Furthermore there is an $\mathrm{SU}(2)$ character on M with boundary traces (a,b,c,d) if and only if there is an x such that (a,b,x) and (c,d,x) are $\mathrm{SU}(2)$ characters on triply-punctured spheres. To construct this character, amalgamate the two representations along the common boundary X as in §2.5; since two elements of $\mathrm{SU}(2)$ have the same trace if and only if they are conjugate, we can ensure that the two representations agree on this boundary.

Recall from §2.2 that (a, b, x) is an SU(2)-character for the triply-punctured sphere if and only if $a, b, x \in [-2, 2]$ and $\kappa_{a,b}(x) \leq 0$. For $a, b \in \mathbb{R}$, define the interval $I_{a,b} \subset \mathbb{R}$ by:

$$\begin{aligned} I_{a,b} &= \{x \in \mathbb{R} \mid \kappa_{a,b}(x) \leq 0\} \\ &= [i^-_{a,b}, i^+_{a,b}] \end{aligned}$$

where

$$\begin{aligned} i^-_{a,b} &= \frac{ab - \sqrt{(a^2-4)(b^2-4)}}{2} \\ i^+_{a,b} &= \frac{ab + \sqrt{(a^2-4)(b^2-4)}}{2} \end{aligned}$$

Since $\kappa_{a,b}(\pm 2) \geq 0$,

$$I_{a,b} \subset [-2, 2]$$

For given (a, b, c, d) with $-2 < a, b, c, d < 2$, the values of $x \in \mathbb{R}$ such that (a, b, c, d, x) extends to a character of an SU(2)-representation comprise the intersection

$$I_{a,b} \cap I_{c,d} \subset [-2, 2].$$

Furthermore the possible $(y, z) \in \mathbb{R}^2$ comprise a conic, which by (2.6.2), is an ellipse. Thus characters of SU(2)-representations with boundary traces (a, b, c, d) form a pencil of ellipses over the interval $I_{a,b} \cap I_{c,d}$.

However, if $I_{a,b} \cap I_{c,d} = \emptyset$, there will be a compact component of $V_{a,b,c,d}(\mathbb{R})$ not corresponding to SU(2)-representations. By applying a permutation if necessary, assume that

$$i^+_{a,b} < i^-_{c,d}$$

Then a component of $V_{a,b,c,d}(\mathbb{R})$ will map under

$$x : V_{a,b,c,d}(\mathbb{R}) \longrightarrow \mathbb{R}$$

to the interval $[i^+_{a,b}, i^-_{c,d}]$. By (2.6.2), the restriction of x to $x^{-1}([-2, 2])$ is proper. If

$$x_0 \in I_{a,b} \cup I_{c,d}$$

then

$$\frac{\kappa_{a,b}(x_0)\kappa_{c,d}(x_0)}{4 - x_0^2} < 0$$

and by (2.6.2), the preimage $x^{-1}(x_0) = \emptyset$. Thus

$$x^{-1}([i^+_{a,b}, i^-_{c,d}])$$

is a compact component of $V_{a,b,c,d}(\mathbb{R})$ not containing characters of SU(2)-representations. Since real characters are characters of either

SU(2)-representations or $\mathrm{SL}(2,\mathbb{R})$-representations, this compact component must consist of $\mathrm{SL}(2,\mathbb{R})$-representations.

Thus, the compact component of $V_{a,b,c,d}(\mathbb{R})$ consists of SU(2)-representations if and only if

$$I_{a,b} \cap I_{c,d} \neq \emptyset$$

and otherwise consists of $\mathrm{SL}(2,\mathbb{R})$-representations. The latter case occurs when

$$i^+_{a,b} < i^-_{c,d}$$

or

$$i^+_{c,d} < i^-_{a,b}.$$

These conditions are equivalent to conditions (5.0.1) and (5.0.2) by the following argument:

The condition $i^+_{a,b} < i^-_{c,d}$ is equivalent to:

$$\sqrt{(a^2-4)(b^2-4)} + \sqrt{(c^2-4)(d^2-4)} < cd - ab$$

and $i^+_{c,d} < i^-_{a,b}$ is equivalent to:

$$\sqrt{(a^2-4)(b^2-4)} + \sqrt{(c^2-4)(d^2-4)} < ab - cd$$

The following lemma (whose straightforward proof is omitted) then applies:

Lemma 5.0.1. *Let p, q, r be nonnegative real numbers. Then*

$$\sqrt{p} + \sqrt{q} < r$$

if and only if

$$r^4 - 2r^2(p+q) + (p-q)^2 > 0$$
$$r^2 - p - q > 0.$$

Taking

$$p = (a^2-4)(b^2-4)$$
$$q = (c^2-4)(d^2-4)$$
$$r = |ab - cd|$$

and calculating that

$$r^4 - 2r^2(p+q) + (p-q)^2 = 16\Delta(a,b,c,d)$$
$$r^2 - p - q = 2(16 - abcd - 2(a^2+b^2+c^2+d^2)),$$

the compact component corresponds to $\mathrm{SL}(2,\mathbb{R})$-representations if and only if (5.0.1) and (5.0.2). □

Here is an explicit example. Let $a, b, c = 3/2$ and $d = -3/2$. For example,

$$\alpha = \begin{bmatrix} 1 & 1 \\ -1/2 & 1/2 \end{bmatrix}, \quad \beta = \begin{bmatrix} 1 & 1/2 \\ -1 & 1/2 \end{bmatrix}, \quad \gamma = \begin{bmatrix} 1 & 1 \\ -1/2 & 1/2 \end{bmatrix}$$

generates an $\mathrm{SL}(2, \mathbb{R})$-representation with the desired character. Furthermore, its (x, y, z) traces are $(0, 0, 1/4)$. Now the equation for the relative character variety at these boundary traces is

$$x^2 + y^2 + z^2 + xyz - 1/16 = 0$$

which does not intersect the planes $x = \pm 2$, $y = \pm 2$, or $z = \pm 2$ at real points.

References

1. Benedetto,R., Senior Thesis, Harvard University (1993)
2. Benedetto,R. and Goldman, W., *Investigating moduli spaces with Geomview,* in "Imagine That!," A quarterly publication of the Geometry Center, (1994)
3. Fricke, R., and Klein, F., *Vorlesungen der Automorphen Funktionen,* Teubner, Leipzig, Vol. I (1897), VolII (1912)
4. Geomview, Software from the Geometry Center, University of Minnesota
5. Goldman, W., *Geometric structures and varieties of representations,* in "The Geometry of Group Representations," Contemp. Math. **74**, Amer. Math. Soc. (1988), 169–198
6. Goldman, W., *Topological components of spaces of representations,* Inv. Math. Vol. 93 No. 3, (1988), pp. 557–607.
7. Goldman, W., *Ergodic theory on moduli spaces,* Ann. Math. (to appear)
8. Lubotzky, A. and Magid, A., *Varieties of representations of finitely generated groups,*, Memoirs Amer. Math. Soc. **336** (vol. 58) (1985)
9. Magnus, W., *Rings of Fricke characters and automorphism groups of free groups,* Math. Zeit. **170** (1980), 91–103
10. Mathematica, Software from Wolfram Research, Inc.
11. Morgan, J.W. and Shalen, P.B., *Valuations, trees and degenerations of hyperbolic structures I,* Ann. Math. **120** (1984), 401–476
12. Procesi, C., *Invariant theory of n by n matrices,* Adv. Math. **19** (1976), 306–381

List of Figures

1 A triply-punctured sphere and its three boundary components 7

2 A quadruply-punctured sphere 9

3 What $V_{a,b,c,d}(\mathbb{R})$ looks like near infinity 18

4 How the curve $\mathcal{C}$ meets the square Q 23

5 Case 1 27

6 Case 2 28

7 Case 3 29
8 Case 4 30
9 Case 5 31
10 Case 6 32
11 Class 1: $abcd < 0$ and none of $|a|, |b|, |c|, |d| < 2$ 37
12 Class 2: $abcd > 0$ and none of $|a|, |b|, |c|, |d| < 2$ 37
13 Class 3: One of $|a|, |b|, |c|, |d| < 2$ 38
14 Class 4: Two of $|a|, |b|, |c|, |d| < 2$ 38
15 Class 5: Three of $|a|, |b|, |c|, |d| < 2$ 39
16 Class 6: All of $|a|, |b|, |c|, |d| < 2$ 39

Mathematics Department, Brown University, Providence, RI 02906 USA *(Benedetto)*, Mathematics Department, University of Maryland, College Park, MD 20742 USA *(Goldman)*

E-mail address: benny@math.brown.edu *(Benedetto)*, wmg@math.umd.edu *(Goldman)*



FIGURE 11. Class 1: $abcd < 0$ and none of $|a|, |b|, |c|, |d| < 2$



FIGURE 12. Class 2: $abcd > 0$ and none of $|a|, |b|, |c|, |d| < 2$



FIGURE 13. Class 3: One of $|a|, |b|, |c|, |d| < 2$



FIGURE 14. Class 4: Two of $|a|, |b|, |c|, |d| < 2$



FIGURE 15. Class 5: Three of $|a|, |b|, |c|, |d| < 2$



FIGURE 16. Class 6: All of $|a|, |b|, |c|, |d| < 2$