MATH858D

BASIC CONCEPTS OF PROBABILITY

MARIA CAMERON

Contents

1. Definitions 1
2. Expected values and moments 3
3. Independence, joint distributions, covariance 4
4. Chebyshev inequality 8
5. Types of convergence of random variables 9
6. Laws of Large Numbers and the Central Limit Theorem 12
7. Conditional probability and conditional expectation 13
References 19

1. Definitions

- A **sample space** Ω is the set of all possible outcomes.
- An **event** A is a subset of Ω.
- A **σ-algebra** $\mathcal{B}$ is a subset of the set of all subsets of Ω satisfying the following axioms
  (1) $\emptyset \in \mathcal{B}$ and $\Omega \in \mathcal{B}$;
  (2) If $B \in \mathcal{B}$ then $B^c \in \mathcal{B}$ (B^c is the complement of B in Ω, i.e., $B^c \equiv \Omega \backslash B$).
  (3) If $\mathcal{A} = \{A_1, \dots, A_n, \dots\}$ is a finite or countable collection in $\mathcal{B}$ then

$$\bigcup_i A_i \in \mathcal{B}.$$

Corollary: If $\mathcal{A} = \{A_1, \dots, A_n, \dots\}$ is a finite or countable collection in $\mathcal{B}$ then

$$\bigcap_i A_i \in \mathcal{B}.$$

Indeed,

$$\bigcap_i A_i = \left(\bigcup_i A_i^c \right)^c.$$

Example 1 Suppose you are tossing a die. For a single throw, the sample space is $\Omega = \{1, 2, 3, 4, 5, 6\}$. If you are interested in particular number on the top, the natural choice of the σ-algebra is the set of all

subsets of Ω. Then $|\mathcal{B}| = 2^6 = 64$. If you are interested only in whether the outcome is odd or even, then a reasonable choice of σ-algebra is

$$\mathcal{B} = \{\emptyset, \{1,3,5\}, \{2,4,6\}, \{1,2,3,4,5,6\}\}.$$

If you are interested only whether there is an outcome or not, you can choose the coarsest σ-algebra

$$\mathcal{B} = \{\emptyset, \{1,2,3,4,5,6\}\}.$$

- A **probability measure** P is a function $P : \mathcal{B} \to [0,1]$ such that
  (1) $P(\Omega) = 1$;
  (2) $0 \le P(A) \le 1$ for all $A \in \mathcal{B}$.
  (3) **Countable additivity:** If $\mathcal{A} = \{A_1, \dots, A_n, \dots\}$ is a finite or countable collection in $\mathcal{B}$ such that $A_i \cap A_j = \emptyset$ for any i, j, then

$$P\left(\bigcup_i A_i\right) = \sum_i P(A_i).$$

  Corollary: $P(\emptyset) = 0$. Indeed,

$$1 = P(\Omega) = P(\Omega \cup \emptyset) = P(\Omega) + P(\emptyset) = 1 + P(\emptyset).$$

  Hence, $P(\emptyset) = 0$.
- A **probability space** is the triple $(\Omega, \mathcal{B}, P)$.
- A **random variable** η is a $\mathcal{B}$-measurable function $\eta : \Omega \to \mathbb{R}$.
  A function is called $\mathcal{B}$-measurable if the preimage of any measurable subset of $\mathbb{R}$ is in $\mathcal{B}$. It is proven in analysis that it is suffices to check that

$$\{\omega \in \Omega \mid \eta(\omega) \le x\} \in \mathcal{B} \text{ for any } x \in \mathbb{R}.$$

- A **probability distribution function** of a random variable η is defined by

$$F_\eta(x) = P(\{\omega \in \Omega \mid \eta(\omega) \le x\}) = P(\eta \le x).$$

Theorem 1. *If F is a probability distribution function then*
(1) *F is nondecreasing, i.e. $x < y$ implies $F(x) \le F(y)$.*
(2) $\lim_{x\to-\infty} F(x) = 0, \quad \lim_{x\to\infty} F(x) = 1.$
(3) *$F(x)$ is continuous from the right for every $x \in \mathbb{R}$, i.e.,*

$$\lim_{y\to x+0} F(y) = F(x).$$

Example 2 Suppose you are tossing a die. Consider the probability space

$$(\Omega = \{1,2,3,4,5,6\}, \mathcal{B} = 2^\Omega, P(\omega) = \tfrac{1}{6}), \tag{1}$$

where 2^Ω is the set of all subsets of Ω, and $\omega \in \Omega = \{1,2,3,4,5,6\}$. Consider the random variable $\eta(\omega) = \omega$. The probability distribution

function is given by

$$F_\eta(x) = \begin{cases} 0, & x < 1, \\ j/6, & j \le x < j+1,\ j = 1,2,3,4,5 \\ 1, & x \ge 6. \end{cases}$$

- Suppose $F'_\eta(x)$ exists. Then $f_\eta(x) \equiv F'_\eta(x)$ is called the **probability density function (pdf)** of the random variable η, and

$$P(x < \eta \le x + dx) = F_\eta(x + dx) - F_\eta(x) = f_\eta(x)dx + o(dx).$$

Example 3 Gaussian density

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-m)^2}{2\sigma^2}},$$

where m and σ are constants. m is the mean, while σ is the standard deviation.

Example 4 Density of an exponential random variable

$$f(x) = \begin{cases} e^{-x}, & x \ge 0, \\ 0, & x < 0. \end{cases}.$$

2. Expected values and moments

Definition 1. *Let $(\Omega, \mathcal{B}, P)$ be a probability space, and η be a random variable. Then the expected value, or mean, of the random variable η is defined as*

$$E[\eta] = \int_\Omega \eta(\omega) dP. \tag{2}$$

If Ω is a discrete set,

$$E[\eta] = \sum_i \eta(\omega_i) P(\omega_i).$$

Example 5 Suppose you are tossing a die. Consider the probability space (1) and the random variable $\eta(\omega) = \omega$, $\omega = 1,2,3,4,5,6$. The expected value of η is

$$E[\eta] = \sum_{j=1}^{6} j \tfrac{1}{6} = 3.5$$

Suppose that the random variable η is fixed. Then we will omit the subscript in the notation of its probability distribution function: $F_\eta(x) \equiv F(x)$.

The integral in Eq. (2) can be rewritten using $F(x)$:

$$E[\eta] = \int_{\mathbb{R}} x P(x < \eta \le x + dx) = \int_{-\infty}^{\infty} x dF(x).$$

If a derivative $f(x)$ of the probability distribution function F exists, then

$$E[\eta] = \int_{-\infty}^{\infty} x f(x) dx.$$

If g is a continuous function defined on the range of the random variable η (on $\eta(\Omega)$), then the expected value of this function is

$$E[g(\eta)] = \int_{-\infty}^{\infty} g(x) dF(x).$$

Moments: Let us take $g(x) = x^n$.

$$E[\eta^n] = \int_{-\infty}^{\infty} x^n dF(x).$$

Central moments: Let us take $g(x) = (x - E[\eta])^n$.

$$E[(\eta - E[\eta])^n] = \int_{-\infty}^{\infty} (x - E[\eta])^n dF(x).$$

Variance = 2nd central moment:

$$\mathrm{Var}(\eta) = E[(\eta - E[\eta])^2) = \int_{-\infty}^{\infty} (x - E[\eta])^2 dF(x).$$

Example 6 Suppose you are tossing a die. Consider the probability space (1) and the random variable $\eta(\omega) = \omega$, $\omega = 1, 2, 3, 4, 5, 6$. The variance of η is

$$\mathrm{Var}(\eta) = \frac{1}{6} \sum_{j=1}^{6} (j - 3.5)^2 = \frac{35}{12} = 2.91(6).$$

The standard deviation:

$$\sigma(\eta) = \sqrt{\mathrm{Var}(\eta)}.$$

3. Independence, joint distributions, covariance

- Two events $A, B \in \mathcal{B}$ are **independent** if

$$P(A \cap B) = P(A)P(B).$$

- Two random variables η_1 and η_2 are independent if the events

$$\{\omega \in \Omega \mid \eta_1(\omega) \le x\} \text{ and } \{\omega \in \Omega \mid \eta_2(\omega) \le y\} \tag{3}$$

are independent for all $x, y \in \mathbb{R}$.

Example 7 Suppose you are tossing a die twice. Consider the probability space

$$(\Omega = \{1, 2, 3, 4, 5, 6\}^2, \mathcal{B} = 2^{\Omega^2}, P(\{\omega_1, \omega_2\}) = 1/36), \quad 1 \le \omega_1, \omega_2 \le 6. \tag{4}$$

Let η_1 and η_2 be random variables equal to the outcomes of the first and the second throws respectively. These random variables are independent.

TABLE 1. Two throws of a die. Values of the random variables $\xi(\omega_1, \omega_2) = \omega_1 + \omega_2$ (left) and $\beta(\omega_1, \omega_2) = \omega_1 - \omega_2$ (right).

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

	1	2	3	4	5	6
1	0	1	2	3	4	5
2	-1	0	1	2	3	4
3	-2	-1	0	1	2	3
4	-3	-2	-1	0	1	2
5	-4	-3	-2	-1	0	1
6	-5	-4	-3	-2	-1	0

Now consider the random variables $\eta(\omega_1, \omega_2) = \omega_1$ and $\xi(\omega_1, \omega_2) = \omega_1 + \omega_2$ (see Table 1, left). We can show that η and ξ are dependent by taking e.g., $x = 1$ and $y = 2$ in Eq. (3):

$$P(\eta \leq 1 \ \& \ \xi \leq 2) = \tfrac{1}{36} \neq P(\eta \leq 1)P(\xi \leq 2) = \tfrac{1}{6} \cdot \tfrac{1}{36} = \tfrac{1}{216}.$$

Finally, we consider the random variables $\xi(\omega_1, \omega_2) = \omega_1 + \omega_2$ and $\beta(\omega_1, \omega_2) = \omega_1 - \omega_2$ (see Table 1, right). We can show that they are dependent by taking e.g., $x = 2$ and $y = -1$ in Eq. (3):

$$P(\xi \leq 2 \ \& \ \beta \leq -1) = 0 \neq P(\xi \leq 2)P(\beta \leq -1) = \tfrac{1}{36} \cdot \tfrac{15}{36} = \tfrac{5}{432}.$$

- The joint distribution function of two random variables η_1 and η_2 is given by

$$F_{\eta_1\eta_2}(x, y) = P(\{\omega \in \Omega \mid \eta_1(\omega) \leq x, \ \eta_2(\omega) \leq y\}) = P(\eta_1(\omega) \leq x, \ \eta_2(\omega) \leq y).$$

- If the second mixed derivative of $F_{\eta_1\eta_2}$ exists, it is called the **joint probability density** of η_1 and η_2 and denoted by

$$f_{\eta_1\eta_2}(x, y) := \frac{\partial F_{\eta_1\eta_2}(x, y)}{\partial x \partial y}.$$

In this case,

$$F_{\eta_1,\eta_2}(x, y) = \int_{-\infty}^{x} \int_{-\infty}^{y} f_{\eta_1\eta_2}(x, y) dx dy.$$

Exercise Show that two random variables are independent if and only if

$$F_{\eta_1\eta_2}(x, y) = F_{\eta_1}(x) F_{\eta_2}(y).$$

Furthermore, if the joint pdf $f_{\eta_1\eta_2}(x, y)$ exists, then

$$f_{\eta_1\eta_2}(x, y) = f_{\eta_1}(x) f_{\eta_2}(y).$$

- Given the joint pdf $f_{\eta_1\eta_2}$, one can obtain $f_{\eta_1}(x)$ by

$$f_{\eta_1}(x) = \int_{-\infty}^{\infty} f_{\eta_1\eta_2}(x, y) dy.$$

In this equation, f_{η_1} is called a **marginal** of $f_{\eta_1\eta_2}$, and the variable η_2 is said to have been integrated out.

- **Properties of expected value and variance** It follows from the definition, that the expected value is a linear functional:

$$E[a\eta_1 + b\eta_2] = aE[\eta_1] + bE[\eta_2]. \tag{5}$$

-

$$\mathrm{Var}(a\eta) = a^2\mathrm{Var}(\eta). \tag{6}$$

- If η_1 and η_2 are independent, then

$$\mathrm{Var}(\eta_1 + \eta_2) = \mathrm{Var}(\eta_1) + \mathrm{Var}(\eta_2). \tag{7}$$

Example 8 Suppose you are tossing a die twice. Consider the probability space and random variables introduced in Example 7. Then

$$E[\xi] = E[\eta_1 + \eta_2] = E[\eta_1] + E[\eta_2] = 7.$$

$$E[\beta] = E[\eta_1 - \eta_2] = E[\eta_1] + E[-\eta_2] = 0.$$

$$\mathrm{Var}[\xi] = \mathrm{Var}[\eta_1 + \eta_2] = \mathrm{Var}[\eta_1] + \mathrm{Var}[\eta_2] = \tfrac{35}{6} = 5.8(3).$$

$$\mathrm{Var}[\beta] = \mathrm{Var}[\eta_1 - \eta_2] = \mathrm{Var}[\eta_1] + \mathrm{Var}[-\eta_2] = \mathrm{Var}[\eta_1] + \mathrm{Var}[\eta_2] = \tfrac{35}{6} = 5.8(3).$$

Example 9 Consider the Bernoulli random variable

$$\eta = \begin{cases} 1, & P(1) = p, \\ 0, & P(0) = 1 - p. \end{cases} \tag{8}$$

Its expected value and variance are

$$E[\eta] = 1 \cdot p + 0 \cdot (1 - p) = p,$$

$$\mathrm{Var}(\eta) = (1 - p)^2 \cdot p + (0 - p)^2 \cdot (1 - p) = p(1 - p).$$

Now consider the sum of n independent copies of η:

$$\xi := \sum_{i=1}^{n} \eta_i.$$

Using Eq. (5) we calculate $E[\xi]$:

$$E[\xi] = \sum_{\imath=1}^{n} E[\eta_i] = np.$$

Since η_i, $1 \le i \le n$, are independent, we can calculate $\mathrm{Var}(\xi)$ using Eq. (7):

$$\mathrm{Var}(\xi) = \sum_{i=1}^{n} \mathrm{Var}(\eta_i) = np(1 - p).$$

Finally, consider the average of n independent copies of η:

$$\zeta := \frac{1}{n}\sum_{i=1}^{n}\eta_i \equiv \frac{\xi}{n}.$$

Using Eqs. (5) and (6), we find

$$E[\zeta] = p,$$

$$\mathrm{Var}(\zeta) = \mathrm{Var}\left(\frac{\xi}{n}\right) = \frac{1}{n^2}\mathrm{Var}(\xi) = \frac{p(1-p)}{n}.$$

- The **covariance** of two random variables η_1 and η_2 is defined by

$$\mathrm{Cov}(\eta_1,\eta_2) = E[(\eta_1 - E[\eta_1])(\eta_2 - E[\eta_2])].$$

Remark If η_1 and η_2 are independent, then $\mathrm{Cov}(\eta_1,\eta_2) = 0$. If $\mathrm{Cov}(\eta_1,\eta_2) = 0$ then η_1 and η_2 are uncorrelated. Note that uncorrelated random variables are not necessarily independent.

Example 10 Suppose you are tossing a die twice. Consider the probability space and random variables introduced in Example 7. As we have established in Example 7, ξ and β are dependent. However, they are uncorrelated. Indeed,

$$\mathrm{Cov}(\xi,\beta) = \sum_{1\le\omega_1\le 6,\ 1\le\omega_2\le 6}(\omega_1+\omega_2-7)(\omega_1-\omega_2)P(\{\omega_1,\omega_2\})$$

$$= \frac{1}{36}\left(\sum_{\omega_1<\omega_2}(\omega_1+\omega_2-7)(\omega_1-\omega_2) + \sum_{\omega_1>\omega_2}(\omega_1+\omega_2-7)(\omega_1-\omega_2)\right) = 0.$$

Example 11 A vector-valued random variable $\eta = [\eta_1,\dots,\eta_n]$ is jointly Gaussian if

$$P(x_1 < \eta_1 \le x_1 + dx_1,\dots,x_n < \eta_n \le x_n + dx_n) = \frac{1}{Z}e^{-\frac{1}{2}(x-m)^\top A^{-1}(x-m)}dx + o(dx),$$

where $x = [x_1,\dots,x_n]^\top$, $m = [m_1,\dots,m_n]^\top$, $dx = dx_1\dots dx_n$, and A is a symmetric positive definite matrix. The normalization constant Z is given by

$$Z = (2\pi)^{n/2}|A|^{1/2}, \text{ where } |A| = \det A.$$

In the case of jointly Gaussian random variables, the covariance matrix C whose entries are

$$C_{ij} = E[(\eta_i - E[\eta_i])(\eta_j - E[\eta_j])]$$

is equal to A. Two jointly Gaussian random variables are independent if and only if they are uncorrelated.

4. Chebyshev inequality

Theorem 2. *Let η be a random variable. Suppose $g(x)$ is a nonnegative, nondecreasing function (i.e., $g(x) \geq 0$, $g(a) \leq g(b)$ whenever $a < b$). Then for any $a \in \mathbb{R}$*

$$P(\eta \geq a) \leq \frac{E[g(\eta)]}{g(a)}. \tag{9}$$

Proof.

$$\begin{aligned} E[g(\eta)] &= \int_{-\infty}^{\infty} g(x)dF(x) \\ &\geq \int_{a}^{\infty} g(x)dF(x) \geq g(a)\int_{a}^{\infty} dF(x) = g(a)P(\eta \geq a). \end{aligned}$$

□

Given a random variable η we define a random variable

$$\xi := |\eta - E[\eta]|.$$

Define

$$g(x) = \begin{cases} x^2, & x \geq 0, \\ 0, & x < 0. \end{cases}.$$

Plugging this into Eq. (9) we obtain

$$P(|\eta - E[\eta]| \geq a) \leq \frac{\mathrm{Var}(\eta)}{a^2}.$$

Example 12 Suppose you are tossing a die twice. Consider the probability space and random variables introduced in Example 7. We will compare the exact probabilities with their Chebyshev estimates.

$P(|\xi - 7| \geq 1) = P(\xi \neq 7) = 1 - \frac{6}{36} = \frac{5}{6} = 0.8(3)$, $\quad \dfrac{\mathrm{Var}(\xi)}{1} = \frac{35}{6} = 5.8(3)$;

$P(|\xi - 7| \geq 2) = P(\xi \leq 5 \text{ or } \xi \geq 9) = \frac{20}{36} = \frac{5}{9} = 0.(5)$, $\quad \dfrac{\mathrm{Var}(\xi)}{4} = \frac{35}{24} = 1.458(3)$;

$P(|\xi - 7| \geq 3) = P(\xi \leq 4 \text{ or } \xi \geq 10) = \frac{12}{36} = \frac{1}{3} = 0.(3)$, $\quad \dfrac{\mathrm{Var}(\xi)}{9} = \frac{35}{54} = 0.6(481)$;

$P(|\xi - 7| \geq 4) = P(\xi \in \{2, 3, 11, 12\}) = \frac{6}{36} = \frac{1}{6} = 0.1(6)$, $\quad \dfrac{\mathrm{Var}(\xi)}{16} = \frac{35}{96} = 0.36458(3)$;

$P(|\xi - 7| \geq 5) = P(\xi \in \{2, 12\}) = \frac{2}{36} = \frac{1}{18} = 0.0(5)$, $\quad \dfrac{\mathrm{Var}(\xi)}{25} = \frac{35}{150} = 0.2(3)$;

Choosing $a = k\sigma$ we get

$$P(|\eta - E[\eta]| \geq k\sigma) \leq \frac{1}{k^2}.$$

This means that for *any* random variable η defined on *any* probability space we have that the probability that η deviates from its expected value by at least k standard deviations does not exceed $1/k^2$.

Note that the power of Chebyshev's inequality is its weakness at the same time. It is universal as it is valid for *any* random variable defined on *any* probability space. At the same time, due to its generality, it typically gives loose bounds. These bounds cannot be improved in principle, because they are exact for the random variable

$$\eta = \begin{cases} 1, & P = \frac{1}{2k^2}, \\ 0, & P = 1 - \frac{1}{k^2}, \\ -1, & P = \frac{1}{2k^2}. \end{cases}.$$

It is easy to check that $E[\eta] = 0$, $\mathrm{Var}(\eta) = \frac{1}{k^2}$, and

$$P(|\eta| \geq 1) = \tfrac{1}{k^2}.$$

Chebyshev's inequality gives the precise upper bound:

$$P(|\eta| \geq 1) \leq \frac{\mathrm{Var}(\eta)}{1^2} = \frac{1}{k^2}.$$

5. Types of convergence of random variables

Suppose we have a sequence of random variables $\{\eta_1, \eta_2, \ldots\}$. In probability theory, there exist several different notions of convergence of a sequence of random variables $\{\eta_1, \eta_2, \ldots\}$ to some limit random variable η.

- $\{\eta_1, \eta_2, \ldots\}$ **converges in distribution** or **converges weakly**, or **converges in law** to η if

$$\lim_{n\to\infty} F_n(x) = F(x) \text{ for every } x \text{ where } F(x) \text{ is continuous,} \tag{10}$$

  where F_n and F are the probability distribution functions of η_n and η respectively.

  Remark Convergence of pdfs $f_n(x)$ implies convergence of $F_n(x)$. The converse is not true in general. For example, consider $F_n(x) = x - \frac{1}{2\pi n}\sin(2\pi n x)$, $x \in (0,1)$. The corresponding pdf is $f_n(x) = 1 - \cos(2\pi n x)$, $x \in (0,1)$. $\{F_n(x)\}$ converges to $F(x) = x$, i.e., to the uniform distribution, while $\{f_n(x)\}$ does not converge at all.

  Remark In the discrete case, the convergence of probability distributions $f(k) := P(\eta = k)$ implies the convergence of the probability distribution functions.

  Example 13 Consider the sum of n independent copies of the Bernoulli random variable as in Example 9:

$$\xi = \sum_{i=1}^{n} \eta_i, \text{ where } \eta_i = \begin{cases} 1, & P(1) = p, \\ 0, & P(0) = 1 - p. \end{cases} \tag{11}$$

Its probability distribution is the binomial distribution given by

$$f(k;n,p) \equiv P(\xi = k) = \left(\begin{array}{c} n \\ k \end{array} \right) p^k (1-p)^{n-k}, \tag{12}$$

where $\left(\begin{array}{c} n \\ k \end{array} \right)$ is the number of k-combinations of the set of n elements:

$$\left(\begin{array}{c} n \\ k \end{array} \right) = \frac{n!}{k!(n-k)!}.$$

Now we let $n \to \infty$ and $p \to 0$ in such a manner that the product np (i.e., the expected value of ξ) remains constant. We introduce the parameter

$$\lambda := np.$$

Consider the sequence of random variables ξ_n where ξ_n is the sum of n independent copies of Bernoulli random variable with $p = \lambda/n$, i.e,

$$\xi_n = \sum_{i=1}^{n} \eta_i^{(n)}, \text{ where } \eta_i^{(n)} = \begin{cases} 1, & P(1) = \lambda/n, \\ 0, & P(0) = 1 - \lambda/n. \end{cases} \tag{13}$$

Plugging in $p = \lambda/n$ in the results of Example 9 we find the expected value and the variance:

$$E[\xi_n] = n\frac{\lambda}{n} = \lambda.$$

$$\mathrm{Var}(\xi_n) = n\frac{\lambda}{n}\left(1 - \frac{\lambda}{n}\right) = \lambda\left(1 - \frac{\lambda}{n}\right).$$

We will show that the sequence ξ_n converges to the Poisson random variable with parameter λ in distribution. Consider the limit

$$\lim_{n\to\infty} f(k;n,\lambda/n) = \lim_{n\to\infty} \frac{n(n-1)\dots(n-k+1)}{k!}\frac{\lambda^k}{n^k}\left(1 - \frac{\lambda}{n}\right)^{n-k} =$$

$$\frac{\lambda^k}{k!}\lim_{n\to\infty}\frac{n(n-1)\dots(n-k+1)}{n^k}\lim_{n\to\infty}\left(1 - \frac{\lambda}{n}\right)^{n}\lim_{n\to\infty}\left(1 - \frac{\lambda}{n}\right)^{-k}$$

The first limit in the equation above is 1 as $n(n-1)\dots(n-k+1) = n^k + O(n^{k-1})$. The second limit can be calculated using the well-known fact that

$$\lim_{n\to\infty}\left(1 + \frac{1}{n}\right)^n = e.$$

Hence

$$\lim_{n\to\infty}\left(1 - \frac{\lambda}{n}\right)^n = e^{-\lambda}.$$

The third limit is 1. Therefore,

$$\lim_{n\to\infty}\frac{n!}{k!(n-k)!}\left(\frac{\lambda}{n}\right)^k\left(1 - \frac{\lambda}{n}\right)^{n-k} = \frac{\lambda^k}{k!}e^{-\lambda},$$

which is the Poisson distribution with parameter λ.

- $\{\eta_1, \eta_2, \ldots\}$ **converges in probability** to η if for any $\epsilon > 0$

$$\lim_{n\to\infty} P(|\eta_n - \eta| \geq \epsilon) = 0 \tag{14}$$

Remark Convergence in probability implies convergence in distribution.

Proof. We will prove this fact for the case of scalar random variables. We have $\lim_{n\to\infty} P(|\eta_n - \eta| \geq \epsilon) = 0$, we need to prove $\lim_{n\to\infty} P(\eta_n \leq x) = P(\eta \leq x)$ for every x where F_η is continuous. First we show an auxiliary fact that for any two random variables ξ and ζ, $x \in \mathbb{R}$ and $\epsilon > 0$

$$P(\xi \leq a) \leq P(\zeta \leq a + \epsilon) + P(|\xi - \zeta| > \epsilon). \tag{15}$$

Indeed,

$$\begin{aligned} P(\xi \leq a) &= P(\xi \leq a \;\&\; \zeta \leq a + \epsilon) + P(\xi \leq a \;\&\; \zeta > a + \epsilon) \\ &\leq P(\zeta \leq a + \epsilon) + P(\xi - \zeta \leq a - \zeta \;\&\; a - \zeta < -\epsilon) \\ &\leq P(\zeta \leq a + \epsilon) + P(\zeta - \xi < -\epsilon) \\ &\leq P(\zeta \leq a + \epsilon) + P(\zeta - \xi < -\epsilon) + P(\zeta - \xi > \epsilon) \\ &= P(\zeta \leq a + \epsilon) + P(|\zeta - \xi| > \epsilon). \end{aligned}$$

Applying Eq. (15) to $\xi = \eta_n$ and $\zeta = \eta$ with $a = x$ and $a = x - \epsilon$, we get

$$\begin{aligned} P(\eta_n \leq x) &\leq P(\eta \leq x + \epsilon) + P(|\eta_n - \eta| > \epsilon) \\ P(\eta \leq x - \epsilon) &\leq P(\eta_n \leq x) + P(|\eta_n - \eta| > \epsilon). \end{aligned}$$

$$P(\eta \leq x - \epsilon) - P(|\eta_n - \eta| > \epsilon) \leq P(\eta_n \leq x) \leq P(\eta \leq x + \epsilon) + P(|\eta_n - \eta| > \epsilon).$$

Taking the limit $n \to \infty$ and taking into account that $\lim_{i\to\infty} P(|\eta_n - \eta| \geq \epsilon) = 0$, we get

$$F_\eta(x - \epsilon) \leq \lim_{n\to\infty} F_{\eta_n}(x) \leq F_\eta(x + \epsilon).$$

If x is a point of continuity of F_η,

$$\lim_{\epsilon\to 0} F_\eta(x - \epsilon) = \lim_{\epsilon\to 0} F_\eta(x + \epsilon) = F_\eta(x).$$

Therefore, taking the limit $\epsilon \to 0$ we obtain the weak convergence:

$$\lim_{n\to\infty} F_{\eta_n}(x) = F_\eta(x)$$

for any x where $F_\eta(x)$ is continuous. □

Remark The converse is, generally, not true. However, convergence in distribution to a *constant* random variable implies convergence in probability.

- $\{\eta_1, \eta_2, \ldots\}$ **converges almost surely** or **almost everywhere** or **with probability 1** or **strongly** to η if

$$P\left(\lim_{n\to\infty} \eta_n = \eta\right) = 1. \tag{16}$$

Remark Convergence almost surely implies convergence in probability (by Fatou's lemma) and in distribution.

- To summarize,

$$\boxed{\eta_i \to \eta \text{ almost surely}} \Rightarrow \boxed{\eta_i \to \eta \text{ in probability}} \Rightarrow \boxed{\eta_i \to \eta \text{ in distribution}} \tag{17}$$

6. Laws of Large Numbers and the Central Limit Theorem

- Let $\{\eta_1, \eta_2, \ldots\}$ be a sequence of random variables with finite expected values $\{m_1 = E[\eta_1], m_2 = E[\eta_2], \ldots\}$. Define

$$\xi_n = \frac{1}{n}\sum_{i=1}^{n} \eta_i, \quad \bar{\xi}_n = \frac{1}{n}\sum_{i=1}^{n} m_i.$$

Definition 2. (1) *The sequence of random variables η_n satisfies the Law of Large Numbers if $\xi_n - \bar{\xi}_n$ converges to zero in probability, i.e., for any $\epsilon > 0$*

$$\lim_{n\to\infty} P(|\xi_n - \bar{\xi}_n| > \epsilon) = 0.$$

(2) *The sequence of random variables η_n satisfies the Strong Law of Large Numbers if $\xi_n - \bar{\xi}_n$ converges to zero almost surely, i.e., for almost all $\omega \in \Omega$*

$$\lim_{n\to\infty} \xi_n - \bar{\xi}_n = 0.$$

- If the random variables η_n are independent and if $\mathrm{Var}(\eta_i) \le V < \infty$, then the Law of Large Numbers holds by the Chebyshev Inequality (9):

$$\begin{aligned} P(|\xi_n - \bar{\xi}_n| > \epsilon) &= P\left(\left|\sum_{i=1}^{n} \eta_i - \sum_{i=1}^{n} m_i\right| > n\epsilon\right) \\ &\le \frac{\mathrm{Var}(\eta_1 + \ldots + \eta_n)}{\epsilon^2 n^2} \le \frac{V}{\epsilon^2 n} \to 0 \text{ as } n \to \infty. \end{aligned}$$

- **Theorem 3.** *(Khinchin) A sequence of independent identically distributed random variables $\{\eta_i\}$ with $\mathbb{E}[\eta_i] = m$ and $\mathbb{E}[|\eta_i|] < \infty$ satisfies the Law of Large Numbers.*

- **Theorem 4.** *(Kolmogorov) A sequence of independent identically distributed random variables with finite expected value and variance satisfies the Strong Law of Large Numbers.*

Theorem 5. (The central limit theorem) *Let $\{\eta_1, \eta_2, \ldots\}$ be a sequence of independent identically distributed (i.i.d.) random variables with $m = E[\eta_i]$ and $0 < \sigma^2 = \mathrm{Var}(\eta_i) < \infty$, then*

$$\frac{(\sum_{i=1}^n \eta_i) - nm}{\sigma\sqrt{n}} \longrightarrow N(0,1) \text{ in distribution,} \tag{18}$$

i.e., converges weakly to the standard normal distribution $N(0,1)$ (i.e., the Gaussian distribution with mean 0 and variance 1) as $n \to \infty$.

A proof via Fourier transform can be found in [1]. Another proof making use of characteristic functions can be found in [2].

Remark Eq. (18) can be recasted as

$$\frac{1}{n}\sum_{i=1}^n \eta_i \longrightarrow N\left(m, \frac{\sigma^2}{n}\right) \text{ in distribution,} \tag{19}$$

i.e., the average of the first n i.i.d. random variables η_i converges in distribution to the Gaussian random variable with mean $m = E[\eta_i]$ and variance σ^2/n.

7. Conditional probability and conditional expectation

- The conditional probability of an event B given that the event A has happened is given by

$$P(B|A) = \frac{P(A \cap B)}{P(A)}.$$

Note that if A and B are independent, then $P(A \cap B) = P(A)P(B)$ and hence

$$P(B|A) = \frac{P(A)P(B)}{P(A)} = P(B).$$

Example 14 Suppose you are tossing a die twice. Consider the probability space (4). Let A be the event that the outcome of the first throw is even, and B be the event that the sum of the outcomes is ≥ 10. Then (see Table 1)

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{4/36}{1/2} = \frac{2}{9}.$$

Note that $P(B) = 1/6 < P(B|A)$. Hence the events A and B are dependent.

If the event A is fixed, then $P(B|A)$ defines a probability measure on $(\Omega, \mathcal{B})$.

- If η is a random variable on Ω, then conditional expectation of η given the event A is

$$E[\eta|A] = \int_\Omega \eta(\omega) P(d\omega|A) = \int_\Omega \eta(\omega) \frac{P(d\omega \cap A)}{P(A)} = \frac{\int_A \eta(\omega) P(dw)}{P(a)}.$$

Example 15 . Suppose you are tossing a die twice. Consider the probability space (4). Let A be the event that the outcome of the first throw is even, and η be the random variable whose value is the sum of outcomes, i.e., $\eta(\{\omega_1,\omega_2\}) = \omega_1+\omega_2$. Then

$$E[\eta|A] = \sum_{\omega_1=1}^{6}\sum_{\omega_2=1}^{6}(\omega_1+\omega_2)P(\{\omega_1,\omega_2\} \mid \omega_1\in\{2,4,6\}).$$

Let us calculate $P(\{\omega_1,\omega_2\} \mid \omega_1\in\{2,4,6\})$.

$$P(\{\omega_1,\omega_2\} \mid \omega_1\in\{2,4,6\}) = \frac{P(\{\omega_1,\omega_2\}\cap(\omega_1\in\{2,4,6\}))}{P(\omega_1\in\{2,4,6\}}$$

$$= \begin{cases} 0, & \omega_1\in\{1,3,5\},\\ \frac{P(\{\omega_1,\omega_2\})}{P(\omega_1\in\{2,4,6\})} = \frac{1/36}{1/2} = \frac{1}{18}, & \omega_1\in\{2,4,6\}. \end{cases}$$

Now we continue our calculation:

$$E[\omega_1+\omega_2 \mid \omega_1\in\{2,4,6\}] = \sum_{\omega_1\in\{2,4,6\}}\sum_{\omega_2=1}^{6}(\omega_1+\omega_2)\frac{1}{18} = \frac{135}{18} = 7.5.$$

Note that $E[\eta] = 7 \neq E[\eta|A] = 7.5$.

- Now we show how one can construct new random variables using conditional probability. For simplicity, we start with partitioning the set of outcomes Ω into a finite or countable number of disjoint measurable subsets:

$$\Omega = \bigcup_i A_i, \quad \text{where} \quad A_i\in\mathcal{B}, \quad A_i\cap A_j = \emptyset.$$

Definition 3. *Let η be a random variable on the probability space $(\Omega,\mathcal{B},P)$. Let $\mathcal{A} = \{A_i\}$ be a partition of Ω as above. Define a new random variable $E[\eta|\mathcal{A}]$ as follows:*

$$E[\eta|\mathcal{A}] = \sum_i E[\eta|A_i]\chi(A_i), \tag{20}$$

where $\chi(A_i)$ is the indicator function of A_i:

$$\chi(A_i;\omega) = \begin{cases} 1, & \omega\in A_i,\\ 0, & \omega\notin A_i. \end{cases}$$

Remark Note that $E[\eta|\mathcal{A}]$ is a random variable as it is a function of the outcome ω. Indeed,

$$E[\eta|\mathcal{A}](\omega) = E[\eta|A_i] \text{ where } A_i\ni\omega.$$

Example 16 Suppose you are tossing a die twice. Let us partition the set of outcomes as follows:

$$\Omega = \bigcup_{i=1}^{6} \{(\omega_1, \omega_2) \mid \omega_1 = i\}.$$

The corresponding partition $\mathcal{A}$ is

$$\mathcal{A} = \{\{(\omega_1, \omega_2) \mid \omega_1 = i\}\}_{i=1}^{6}.$$

Take the random variable $\xi = \omega_1 + \omega_2$ (see Table 1, left), the sum of numbers on the top. Construct a new random variable

$$\begin{aligned} E[\xi|\mathcal{A}] &= \sum_{i=1}^{6} E[\xi|\omega_1 = i]\chi(\omega_1 = i) = \sum_{i=1}^{6} (i + 3.5)\chi(\omega_1 = i) \\ &= 4.5\chi(\omega_1 = 1) + 5.5\chi(\omega_1 = 2) + 6.5\chi(\omega_1 = 3) \\ &+ 7.5\chi(\omega_1 = 4) + 8.5\chi(\omega_1 = 5) + 9.5\chi(\omega_1 = 6). \end{aligned}$$

- Now we define the conditional expectation of one random variable η given the other random variable θ. First we assume that θ assumes a finite or countable number of values $\{\theta_1, \theta_2, \ldots\}$. Define the partition $\mathcal{A}$ where

$$A_i = \{\omega \in \Omega \mid \theta = \theta_i\}.$$

Definition 4. *We define a new random variable $E[\eta|\theta]$ as a the following function of the random variable θ:*

$$E[\eta|\theta] := E[\eta|\mathcal{A}], \quad \text{i.e.,} \quad E[\eta|\theta] = E[\eta|A_i] \text{ if } \theta = \theta_i.$$

Example 17 Suppose you are tossing a die twice. Let (ω_1, ω_2) be the numbers on the top. Define random variables $\xi = \omega_1 + \omega_2$ and $\theta = \omega_1$. Then it follows from our calculation from the previous example that

$$E[\xi|\theta] = 3.5 + \theta.$$

- Now we give generalizations of $E[\eta|\mathcal{A}]$ and $E[\eta|\theta]$ defined for a partition of Ω into discrete subsets.

Definition 5. *Let $(\Omega, \mathcal{B}, P)$ be a probability space and η be a random variable. Let $\mathcal{A}$ be another σ-algebra defined on Ω that is coarser than $\mathcal{B}$, i.e., if $A \in \mathcal{A}$ then $A \in \mathcal{B}$ (i.e., $\mathcal{A} \subset \mathcal{B}$). Then the conditional expectation of η with respect to the σ-algebra $\mathcal{A}$ is the random variable denoted by $E[\eta|\mathcal{A}]$ satisfying*

$$\int_A E[\eta|\mathcal{A}]P(d\omega) = \int_A \eta(\omega)P(d\omega) \text{ for any } A \in \mathcal{A}.$$

Suppose θ is another random variable on $(\Omega, \mathcal{B}, P)$. The σ-algebra generated by θ is the σ-algebra $\sigma(\theta)$ generated by the sets

$$\{\omega \in \Omega \mid \theta(\omega) \le x\},$$

I.e., $\sigma(\theta)$ is the smallest σ-algebra containing all of these sets. Obviously, since θ is $\mathcal{B}$-measurable, $\sigma(\theta) \subset \mathcal{B}$.

Example 18 Consider the probability space with the set of outcomes $\mathbb{R}^2$, Borel σ-algebra $\mathcal{B}$ (i.e., the one generated by all open sets) and the probability measure

$$P(B) = \int_B \frac{1}{Z} e^{-(x^2+y^2)} dxdy, \quad Z = \int_{\mathbb{R}^2} e^{-(x^2+y^2)} dxdy = \pi, \quad B \in \mathcal{B}.$$

Consider the random variables $\eta(x,y) = x$ and $\theta(x,y) = \sqrt{x^2+y^2}$. The σ-algebra $\sigma(\theta)$ is generated by all balls centered at the origin:

$$\{(x,y) \in \mathbb{R}^2 \mid \sqrt{x^2+y^2} \le z\}.$$

Definition 6. *The conditional expectation $E[\eta|\theta]$ of a random variable η given a random variable θ is the conditional expectation of η with respect to the σ-algebra $\sigma(\theta)$ generated by the random variable θ, i.e.,*

$$E[\eta|\theta] = E[\eta|\sigma(\theta)].$$

- Consider the case where the joint pdf of random variables η and θ $f_{\eta,\theta}(x,y)$ exists. Then we define the conditional probability distribution

$$f_{\eta|\theta}(x|y) := \frac{f_{\eta,\theta}(x,y)}{f_\theta(y)}. \tag{21}$$

Then

$$P(a < \eta \le b \mid \theta = y) = \int_a^b f_{\eta|\theta}(x|y) dx,$$

where the left-hand side of the equation above is understood as

$$P(a < \eta \le b \mid \theta = y) = \lim_{\epsilon \to 0+0} P(a < \eta \le b \mid |\theta - y| < \epsilon).$$

Example 19 Consider the probability space as in Example 18. Define the random variables $\eta(x,y) = x$ and $\theta(x,y) = \sqrt{x^2+y^2}$. We want to calculate

$$P(a < \eta \le b \mid \theta = z) = P(a < x \le b \mid \sqrt{x^2+y^2} = z)$$

The set $\sqrt{x^2+y^2} = z$ is a circle centered at the origin of radius z. Since the probability density on every circle is uniform, this probability is the ratio of the total arc length of segments of the circle with $a < x \le b$ to the arc length of the circle (see Fig. 1). Therefore,

$$P(a < \eta \le b \mid \theta = z) = \frac{1}{\pi} \left(\arccos\left(\frac{\max\{a,-z\}}{z}\right) - \arccos\left(\frac{\min\{b,z\}}{z}\right) \right);$$



FIGURE 1. Illustration to Example 19. The subsets on the circle $\theta = \sqrt{x^2+y^2} = z$ where $a < \eta = x \le b$ are shown in red.

The conditional expectation of η given θ is

$$E[\eta|\theta] = \int_{-\infty}^{\infty} x f_{\eta|\theta}(x|y)dx.$$

The conditional variance is defined by

$$\mathrm{Var}(\eta|\theta) := E[|\eta - E[\eta|\theta]|^2 \mid \theta].$$

Example 20 Suppose the joint pdf of random variables η and θ is given by

$$f_{\eta,\theta}(x,y) = \frac{1}{Z} e^{-\beta(x^2+y^2+x^2y^2)} \quad \text{where} \quad Z := \int_{\mathbb{R}^2} e^{-\beta(x^2+y^2+x^2y^2)} dxdy$$

is the partition function. Note that this pdf is the Gibbs measure for the overdamped Langevin dynamics in the potential energy landscape $V(x,y) = x^2 + y^2 + x^2y^2$. Level sets of this potential are shown in Fig. 2. Let us find $f_{\eta|\theta}(x|y)$, $E[\eta|\theta]$, and $\mathrm{Var}(\eta|\theta)$. First we find the marginal density

$$f_\theta(y) = \frac{1}{Z}\int_{-\infty}^{\infty} e^{-\beta(x^2+y^2+x^2y^2)} dx = \frac{1}{Z}\sqrt{\frac{\pi}{\beta(1+y^2)}} e^{-\beta y^2}.$$

Next, we find

$$f_{\eta|\theta}(x|y) = \frac{\frac{1}{Z} e^{-\beta(x^2+y^2+x^2y^2)}}{\frac{1}{Z}\sqrt{\frac{\pi}{\beta(1+y^2)}} e^{-\beta y^2}} = \sqrt{\frac{\beta(1+y^2)}{\pi}} e^{-\beta x^2(y^2+1)}.$$



FIGURE 2. Level sets of the potential $V(x,y) = x^2 + y^2 + x^2y^2$.

Then the conditional expectation of η given θ is

$$E[\eta|\theta] = \int_{-\infty}^{\infty} x\sqrt{\frac{\beta(1+y^2)}{\pi}}e^{-\beta x^2(y^2+1)}dx = 0.$$

Finally, we find $\mathrm{Var}(\eta|\theta)$:

$$\begin{aligned}\mathrm{Var}(\eta|\theta) =& \int_{-\infty}^{\infty} x^2\sqrt{\frac{\beta(1+y^2)}{\pi}}e^{-\beta x^2(y^2+1)}dx\\ =&\sqrt{\frac{\beta(1+y^2)}{\pi}}\frac{\sqrt{\pi}}{2\beta^{3/2}(1+y^2)^{3/2}} = \frac{1}{2\beta(1+y^2)}.\end{aligned}$$

- **Conditional expectation as the best approximation.** Imagine that you are considering two random variables η and θ, and you wish to approximate η with a function of θ. We will show that the best approximation of η by a function of θ in the least squares sense is $E[\eta|\theta]$.

Theorem 6. *Let $g(\theta)$ be any measurable function of θ. Then*

$$E[(\eta - E[\eta|\theta])^2] \le E[(\eta - g(\theta))^2]. \tag{22}$$

Proof. We will prove this fact for the case where the set of values of θ is at most countable: $\theta(\omega) \in \{\theta_1, \theta_2, \ldots\}$. Any function $g(\theta)$ can be written as

$$g(\theta) = E[\eta|\theta] + (g(\theta) - E[\eta|\theta]).$$

We plug this into the right-hand side of Eq. (22) and partition the set of outcomes Ω into nonintersecting subsets

$$Z_i = \{\omega \in \Omega \mid \theta(\omega) = \theta_i\}.$$

We have:

$$\begin{aligned}
E[(\eta - g(\theta))^2] &= \int_\Omega (\eta(\omega) - E[\eta|\theta] - (g(\theta) - E[\eta|\theta]))^2 P(d\omega) \\
&= \sum_i \int_{Z_i} (\eta - E[\eta|\theta] - (g(\theta) - E[\eta|\theta]))^2 P(d\omega) \\
&= \sum_i \int_{Z_i} (\eta - E[\eta|\theta])^2 P(d\omega) \\
&-2 \sum_i (g(\theta_i) - E[\eta|Z_i]) \int_{Z_i} (\eta - E[\eta|Z_i]) P(d\omega) \\
&+ \sum_i (g(\theta_i) - E[\eta|Z_i])^2 \int_{Z_i} P(d\omega).
\end{aligned}$$

Taking into account that

$$\int_{Z_i} (\eta - E[\eta|Z_i]) P(d\omega) = E[\eta|Z_i] - E[\eta|Z_i] = 0,$$

we continue:

$$\begin{aligned}
E[(\eta - g(\theta))^2] &= E[(\eta - E[\eta|\theta])^2] + \sum_i (g(\theta_i) - E[\eta|Z_i])^2 P(Z_i) \\
&\geq E[(\eta - E[\eta|\theta])^2].
\end{aligned}$$

□

References

[1] A. Chorin and O. Hald, Stochastic Tools in Mathematics and Science, 3rd edition, Springer 2013
[2] L. Koralov and Ya. Sinai, Theory of probability and stochastic processes, 2nd edition, Springer, 2007
[3] Wikipedia