

BEYOND MEAN FIELD: ON THE ROLE OF PAIR EXCITATIONS IN THE EVOLUTION OF CONDENSATES.

M. GRILLAKIS AND M. MACHEDON

This paper is dedicated to Professor Choquet-Bruhat, whose work on nonlinear wave equations was an inspiration to us earlier in our careers

1. INTRODUCTION

ABSTRACT. This paper is in part a summary of our earlier work [17, 18, 19], and in part an announcement introducing a refinement of the equations for the pair excitation function used in our previous work with D. Margetis. The new equations are Euler-Lagrange equations, and the solutions conserve energy and the number of particles.

2. INTRODUCTION

The problem, which has received a lot of attention in recent years, is concerned with the evolution of the N -body linear Schrödinger equation

$$\begin{aligned} \frac{1}{i} \frac{\partial}{\partial t} \psi_N(t, \cdot) &= H_N \psi_N(t, \cdot) \text{ with} \\ \psi_N(0, x_1, \dots, x_N) &= \phi_0(x_1) \phi_0(x_2) \cdots \phi_0(x_N) \\ \|\psi_N(t, \cdot)\|_{L^2(\mathbb{R}^{3N})} &= 1 \end{aligned}$$

The Hamiltonian is an operator of the form

$$H_N = \sum_{j=1}^N \Delta_{x_j} - \frac{1}{N} \sum_{i < j} v_N(x_i - x_j)$$

where $v_N(x) := N^{3\beta} v(N^\beta x)$ with $0 \leq \beta \leq 1$ models the strength of two body interactions. Notice that if $\beta > 0$ then $v_N(x) \rightarrow \delta(x)$ as $N \rightarrow \infty$.

The authors thank Sergiu Klainerman for the interest shown for this work, and John Millson for many discussions related to the symplectic group and its representations. The authors would like to thank the Department of Applied Mathematics at the University of Crete and ACMAC for their hospitality during the preparation of the present work.

For simplicity we assume that $v \in C_0$ and $v \geq 0$. The goal is to show, in a sense to be made precise,

$$\psi_N(t, x_1, \dots, x_N) \simeq e^{iN\chi(t)} \phi(t, x_1) \phi(t, x_2) \cdots \phi(t, x_N) \quad (1)$$

where ϕ satisfies a suitable non-linear Schrödinger equation. In particular, this approximation is not true in $L^2(\mathbb{R}^{3N})$.

The motivation for this problem is that in the presence of a trap the ground state of H_N looks like

$$\Psi_N(x_1, x_2, \dots, x_N) \simeq \phi_0(x_1) \phi_0(x_2) \cdots \phi_0(x_N)$$

This is suggested by the result of Lieb and Seiringer who showed in [26] that

$$\gamma_1^N(x, x') \rightarrow \phi_0(x) \overline{\phi_0(x')}$$

where

$$\gamma_1^N(x, x') = \int \Psi_N(x, x_2, \dots, x_N) \overline{\Psi_N(x', x_2, \dots, x_N)} dx_2 \cdots dx_N$$

Here $\|\phi_0\|_{L^2} = 1$ and ϕ_0 minimizes the Gross-Pitaevskii functional. See [25] for extensive background.

The reason for the recent attention to this problem is two-fold. On the one hand experimental advances during the last twenty years made the creation and manipulation of condensates in the laboratory possible, on the other hand recent mathematical developments made possible the rigorous treatment of the equations when the number of particles, namely N , is large.

While this is a "classical PDE problem" (as opposed to a Fock space problem), the PDE approach to this problem has only been studied systematically during the last 10-15 years, in the series of papers of Erdős and Yau [8], and Erdős, Schlein and Yau [9] to [11]. See also [7]. These papers prove

$$\gamma_1^N(t, x, x') \rightarrow \phi(t, x) \overline{\phi(t, x')} \quad (2)$$

in trace norm as $N \rightarrow \infty$, and similarly for the higher order marginal density matrices γ_k^N , where k is fixed. The problem becomes more difficult and interesting as the parameter β in the definition of v_N approaches 1. The strategy of these papers is based on the older work of Spohn [30]. Recent simplifications and generalizations, based on harmonic analysis techniques and a "boardgame argument" inspired by the Feynman diagram approach of Erdős, Schlein and Yau, were given in [21], [22], [6], [3], [4], [5]. See also [14], [27] for a different approach.

The symmetric Fock space approach to the problem is much older. It originated in physics, with the papers by Lee, Huang and Yang [23] in the static case, and Wu [31] in the time-dependent case. See also [2]. It continued with the mathematically rigorous work of Hepp [20], and Ginibre and Velo [15].

Motivated by the goal of obtaining a convergence rate to solutions of NLS in (2), Rodnianski and Schlein resumed the rigorous Fock space approach in [28]. This paper, as well as the older work of Wu, served as an inspiration for our work. Our goal is to obtain a refinement to (1) which provides an $L^2(\mathbb{R}^{3N})$ and Fock space estimate. This leads to the introduction of the pair excitation function k .

We also mention the recent preprint [1] where a similar approach (but with an explicit choice of pair excitation function k) is used to prove convergence of the density matrices in the critical case $\beta = 1$.

3. FOCK SPACE

In this section we briefly review symmetric Fock space, following the notation of [19]. See [28], for more details. The elements of \mathbb{F} are vectors of the form

$$|\psi\rangle = (\psi_0, \psi_1(x_1), \psi_2(x_1, x_2), \dots)$$

where $\psi_0 \in \mathbb{C}$ and ψ_k are symmetric L^2 functions. The norm of such a vector is,

$$\| |\psi\rangle \|_{\mathcal{F}}^2 = \langle \psi | \psi \rangle = |\psi_0|^2 + \sum_{n=1}^{\infty} \|\psi_n\|_{L^2}^2.$$

The creation and annihilation distribution valued operators denoted by a_x^* and a_x respectively which act on vectors of the form $(0, \dots, \psi_{n-1}, 0, \dots)$ and $(0, \dots, \psi_{n+1}, 0, \dots)$ by

$$a_x^*(\psi_{n-1}) := \frac{1}{\sqrt{n}} \sum_{j=1}^n \delta(x - x_j) \psi_{n-1}(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_n)$$

$$a_x(\psi_{n+1}) := \sqrt{n+1} \psi_{n+1}([x], x_1, \dots, x_n)$$

with $[x]$ indicating that the variable x is frozen. The vacuum state is defined as follows:

$$|0\rangle := (1, 0, 0 \dots)$$

and $a_x|0\rangle = 0$. One can easily check that $[a_x, a_y^*] = \delta(x - y)$ and since the creation and annihilation operators are distribution valued we can

form operators that act on \mathbb{F} by introducing a field, say $\phi(x)$, and form

$$a(\bar{\phi}) := \int dx \{ \bar{\phi}(x) a_x \} \quad \text{and} \quad a^*(\phi) := \int dx \{ \phi(x) a_x^* \}$$

where by convention we associate a with $\bar{\phi}$ and a^* with ϕ . These operators are well defined, unbounded, on \mathbb{F} provided that ϕ is square integrable. The creation and annihilation operators provide a way to introduce coherent states in \mathbb{F} in the following manner, first define the skew-Hermitian operator

$$\mathcal{A}(\phi) := \int dx \{ \bar{\phi}(x) a_x - \phi(x) a_x^* \} \quad (3)$$

and then introduce N -particle coherent states as

$$|\psi(\phi)\rangle := e^{-\sqrt{N}\mathcal{A}(\phi)}|0\rangle. \quad (4)$$

This is the Weyl operator used by Rodnianski and Schlein in [28]. It is easy to check that

$$e^{-\sqrt{N}\mathcal{A}(\phi)}|0\rangle = \left(\dots c_n \prod_{j=1}^n \phi(x_j) \dots \right) \quad \text{with} \quad c_n = (e^{-N} N^n / n!)^{1/2}.$$

In particular, by Stirling's formula, the main term that we are interested in has the coefficient

$$c_N \approx (2\pi N)^{-1/4} \quad (5)$$

Thus a coherent state introduces a tensor product in each sector of \mathbb{F} .

For the construction analogous to (3) involving quadratics, start with the Lie algebra of real or complex symplectic "matrices" of the form

$$L := \begin{pmatrix} d(x, y) & l(x, y) \\ k(x, y) & -d^T(x, y) \end{pmatrix}$$

where d , k and l are kernels in L^2 , and k and l are symmetric in (x, y) . We denote this Lie algebra $sp(\mathbb{C})$ or $sp(\mathbb{R})$ depending on whether the kernels d , k and l are complex or real. The natural setting for us (which will insure that the Fock space operator $e^{\mathcal{I}(L)}$ defined below, is unitary, see also the appendix of [18]) is the subalgebra $sp_c(\mathbb{R}) = \mathcal{W}sp(\mathbb{R})\mathcal{W}^{-1}$ where

$$\mathcal{W} = \frac{1}{\sqrt{2}} \begin{pmatrix} I & iI \\ I & -iI \end{pmatrix}$$

The elements of $sp_c(\mathbb{R})$ look like

$$L := \begin{pmatrix} id(x, y) & \bar{k}(x, y) \\ k(x, y) & -id^T(x, y) \end{pmatrix} \quad (6)$$

with L^2 kernels d complex and self-adjoint, k complex and symmetric.

Remark 3.1. The corresponding group elements $E \in Sp_c(\mathbb{R})$ (in particular $E = e^L$, $L \in sp_c(\mathbb{R})$) satisfy the following three properties:

- E commutes with the real structure σ defined by $\sigma(\phi, \psi) = (\overline{\psi}, \overline{\phi})$, in other words E is of the form

$$E = \begin{pmatrix} P(x, y) & Q(x, y) \\ \overline{Q}(x, y) & \overline{P}(x, y) \end{pmatrix}$$

- E belongs to the infinite dimensional analogue of $U(n, n)$, in other words

$$E^* \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} E = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}$$

- E is in the symplectic group, meaning

$$E^T \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} E = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$$

In fact, any two of the above imply the third. The conceptual reason for this is that the symplectic inner product $\left((\phi_1, \psi_1), (\phi_2, \psi_2) \right) = \int \phi_1 \psi_2 - \int \psi_1 \phi_2$ and the " $U(n, n)$ " inner product $\left\langle (\phi_1, \psi_1), (\phi_2, \psi_2) \right\rangle = \int \phi_1 \overline{\phi_2} - \int \psi_1 \overline{\psi_2}$ are related by $\left\langle (\phi_1, \psi_1), (\phi_2, \psi_2) \right\rangle = \left((\phi_1, \psi_1), \sigma(\phi_2, \psi_2) \right)$.

See Folland's book [13] for more along these lines in the finite dimensional case. These matrices are called Bogoliubov rotations in [1].

Our approach is based on the map from $L \in sp(\mathbb{C})$ to quadratic polynomials in (a, a^*) in the following manner,

$$\begin{aligned} \mathcal{I}(L) &= \frac{1}{2} \int dx dy \left\{ (a_x, a_x^*) \begin{pmatrix} d(x, y) & l(x, y) \\ k(x, y) & -d(y, x) \end{pmatrix} \begin{pmatrix} -a_y^* \\ a_y \end{pmatrix} \right\} \\ &= -\frac{1}{2} \int dx dy \{ d(x, y) a_x a_y^* + d(y, x) a_x^* a_y + k(x, y) a_x^* a_y^* - l(x, y) a_x a_y \}. \end{aligned} \quad (7)$$

This is the infinite dimensional Segal-Shale-Weil infinitesimal representation. The group representation was studied in [29]. The crucial property of this map is the Lie algebra isomorphism

$$[\mathcal{I}(L_1), \mathcal{I}(L_2)] = \mathcal{I}([L_1, L_2]) \quad (8)$$

Notice that if $L \in sp_c(\mathbb{R})$, then L has the form (6) and $\mathcal{I}(L)$ is skew-Hermitian, thus $e^{\mathcal{I}(L)}$ is a unitary operator on Fock space. For the

applications that follow we will only use the self-adjoint elements of $sp_c(\mathbb{R})$

$$K = \begin{pmatrix} 0 & \bar{k}(t, x, y) \\ k(t, x, y) & 0 \end{pmatrix} \quad (9)$$

and the corresponding

$$\mathcal{B}(k) := \mathcal{I}(K) = \frac{1}{2} \int dx dy \{ \bar{k}(t, x, y) a_x a_y - k(t, x, y) a_x^* a_y^* \} . \quad (10)$$

$$K = \begin{pmatrix} 0 & \bar{k}(t, x, y) \\ k(t, x, y) & 0 \end{pmatrix} \quad (11)$$

We easily compute

$$e^K = \begin{pmatrix} \text{ch}(k) & \overline{\text{sh}(k)} \\ \text{sh}(k) & \overline{\text{ch}(k)} \end{pmatrix}$$

where

$$\text{sh}(k) := k + \frac{1}{3!} k \circ \bar{k} \circ k + \dots , \quad (12a)$$

$$\text{ch}(k) := \delta(x - y) + \frac{1}{2!} \bar{k} \circ k + \dots , \quad (12b)$$

This particular construction and the corresponding unitary operator $e^{\mathcal{B}}$ were introduced in [17].

The Fock Hamiltonian is

$$\mathcal{H} := \mathcal{H}_1 - N^{-1} \mathcal{V} \quad \text{where,} \quad (13a)$$

$$\mathcal{H}_1 := \int dx dy \{ \Delta_x \delta(x - y) a_x^* a_y \} \quad \text{and} \quad (13b)$$

$$\mathcal{V} := \frac{1}{2} \int dx dy \{ v_N(x - y) a_x^* a_y^* a_x a_x \} . \quad (13c)$$

It is a diagonal operator on Fock space, and it acts as a regular PDE Hamiltonian in n variable

$$H_{n, PDE} = \sum_{j=1}^n \Delta_{x_j} - \frac{1}{2N} \sum_{x_j \neq x_k} N^{3\beta} v(N^\beta(x_j - x_k))$$

on the n th component of \mathbb{F} .

4. OUTLINE OF OLDER RESULTS

Our goal is to study the evolution of coherent initial conditions of the form

$$|\psi_{exact}\rangle = e^{it\mathcal{H}} e^{-\sqrt{N}\mathcal{A}(\phi_0)} |0\rangle \quad (14)$$

The papers [17, 18, 19] propose an approximation of the form

$$|\psi_{appr}\rangle := e^{-\sqrt{N}\mathcal{A}(\phi(t))} e^{-\mathcal{B}(k(t))} |0\rangle, \quad (15)$$

and derive Schrödinger type equations for $\phi(t, x)$, $k(t, x, y)$ so that $|\psi_{exact}(t)\rangle \approx e^{iN\chi(t)} |\psi_{appr}(t)\rangle$, with $\chi(t)$ a real phase factor, and find precise estimates in Fock space, see Theorem (4.1) below. Our strategy is to consider

$$|\psi_{red}\rangle = e^{\mathcal{B}(t)} e^{\sqrt{N}\mathcal{A}(t)} e^{it\mathcal{H}} e^{-\sqrt{N}\mathcal{A}(0)} |0\rangle$$

and then find a "reduced Hamiltonian" H_{red} so that

$$\frac{1}{i} \partial_t |\psi_{red}\rangle = \mathcal{H}_{red} |\psi_{red}\rangle. \quad (16)$$

The reduced Hamiltonian is

$$\begin{aligned} \mathcal{H}_{red} := & \frac{1}{i} (\partial_t e^{\mathcal{B}}) e^{-\mathcal{B}} \\ & + e^{\mathcal{B}} \left(\frac{1}{i} (\partial_t e^{\sqrt{N}\mathcal{A}}) e^{-\sqrt{N}\mathcal{A}} + e^{\sqrt{N}\mathcal{A}} \mathcal{H} e^{-\sqrt{N}\mathcal{A}} \right) e^{-\mathcal{B}}. \end{aligned}$$

It can be written abstractly as a composition (in space only) of operators

$$\mathcal{H}_{red} = \frac{1}{i} \frac{\partial}{\partial t} + e^{\mathcal{B}} e^{\sqrt{N}\mathcal{A}} \left(-\frac{1}{i} \frac{\partial}{\partial t} + \mathcal{H} \right) \circ e^{-\sqrt{N}\mathcal{A}} e^{-\mathcal{B}}$$

Explicitly it is

$$\begin{aligned} \mathcal{H}_{red} = & N\mathcal{P}_0 + N^{1/2} e^{\mathcal{B}} \mathcal{P}_1 e^{-\mathcal{B}} \\ & + \mathcal{H}_G + \mathcal{I}(R) - N^{-1/2} e^{\mathcal{B}} \mathcal{P}_3 e^{-\mathcal{B}} - N^{-1} e^{\mathcal{B}} \mathcal{P}_4 e^{-\mathcal{B}} \end{aligned} \quad (17)$$

where the various terms are defined below. \mathcal{P}_n indicate polynomials of degree n in a, a^* to be given explicitly:

$$\begin{aligned} \mathcal{P}_0 := & \int dx \left\{ \frac{1}{2i} (\phi \bar{\phi}_t - \bar{\phi} \phi_t) - |\nabla \phi|^2 \right\} \\ & - \frac{1}{2} \int dx dy \{ v_N(x-y) |\phi(x)|^2 |\phi(y)|^2 \}. \end{aligned} \quad (18)$$

$$\begin{aligned} \mathcal{P}_1 := & \int dx \{ h(t, x) a_x^* + \bar{h}(t, x) a_x \} \\ = & a^*(h(t, \cdot) + a(\bar{h}(t, \cdot))) \end{aligned} \quad (19)$$

where $h := -(1/i)\partial_t\phi + \Delta\phi - (v_N * |\phi|^2)\phi$.

$$\mathcal{H}_G := \frac{1}{2} \int dx dy \{ -g_N(t, x, y) a_y^* a_x - g_N(t, y, x) a_x^* a_y \} \quad (20a)$$

where $\quad (20b)$

$$g_N(t, x, y) := -\Delta_x \delta(x - y) + (v_N * |\phi|^2)(t, x) \delta(x - y) + v_N(x - y) \bar{\phi}(t, x) \phi(t, y) \quad (20c)$$

and

$$\begin{aligned} R &= \frac{1}{i} \left(\frac{\partial}{\partial t} e^K \right) e^{-K} + [G, e^K] e^{-K} + e^K M e^{-K} = \\ &= \begin{pmatrix} -\overline{\mathbf{W}(\text{ch}(k))} & -\overline{\mathbf{S}(\text{sh}(k))} \\ \mathbf{S}(\text{sh}(k)) & \mathbf{W}(\text{ch}(k)) \end{pmatrix} \circ \begin{pmatrix} \text{ch}(k) & -\overline{\text{sh}(k)} \\ -\text{sh}(k) & \overline{\text{ch}(k)} \end{pmatrix} \\ &+ e^K M e^{-K} \end{aligned} \quad (21)$$

where \mathbf{S} describes a Schrödinger type evolution, while \mathbf{W} is a Wigner type operator by

$$\mathbf{S}(s) := \frac{1}{i} s_t + g^T \circ s + s \circ g \quad \text{and} \quad \mathbf{W}(p) := \frac{1}{i} p_t + [g^T, p]$$

$$\text{while } M := \begin{pmatrix} 0 & \bar{m} \\ -m & 0 \end{pmatrix} \quad \text{where}$$

$$m(x, y) := -v_N(x - y) \phi(x) \phi(y), \quad v_N(x) = N^{3\beta} v(N^\beta x)$$

$$\text{and } G := \begin{pmatrix} g & 0 \\ 0 & -g^T \end{pmatrix}.$$

Finally,

$$\mathcal{P}_3 := [\mathcal{A}, \mathcal{V}] = \int dx dy \{ v_N(x - y) (\phi(y) a_x^* a_y^* a_x + \bar{\phi}(y) a_x^* a_x a_y) \} \quad (22a)$$

$$\mathcal{P}_4 := \mathcal{V} = (1/2) \int dx dy \{ v_N(x - y) a_x^* a_y^* a_x a_y \} . \quad (22b)$$

The main result of [19], building on the previous papers of the authors and D. Margetis [17, 18], can be summarized as follows.

Theorem 4.1. *Let ϕ and k satisfy*

$$\frac{1}{i}\partial_t\phi - \Delta\phi + (v_N * |\phi|^2)\phi = 0 \quad (23a)$$

and either one of the following equivalent equations: (23b)

$$1) \left(\mathbf{S}(\text{sh}(k)) - \overline{\text{ch}(k)} \circ m \right) \circ \text{ch}(k) = \left(\mathbf{W}(\overline{\text{ch}(k)}) + \text{sh}(k) \circ \overline{m} \right) \circ \text{sh}(k) \quad (23c)$$

or else the equivalent non-linear equation (23d)

$$2) \mathbf{S}(\text{th}(k)) = m + \text{th}(k) \circ \overline{m} \circ \text{th}(k) \quad (23e)$$

where $\text{th}(k) := \overline{\text{ch}(k)}^{-1} \circ \text{sh}(k)$ (23f)

or else the equivalent system of linear equations (23g)

$$3a) \mathbf{S}(\text{sh}(2k)) = m_N \circ \text{ch}(2k) + \overline{\text{ch}(2k)} \circ m_N \quad (23h)$$

$$3b) \mathbf{W}(\overline{\text{ch}(2k)}) = m_N \circ \overline{\text{sh}(2k)} - \text{sh}(2k) \circ \overline{m}_N . \quad (23i)$$

with prescribed initial conditions $\phi(0, \cdot) = \phi_0$, $k(0, \cdot, \cdot) = 0$. If ϕ , k satisfy the above equations, then there exists a real phase function χ such that

$$\left\| \left| \psi_{\text{exact}}(t) \right\rangle - e^{iN\chi(t)} \left| \psi_{\text{appr}}(t) \right\rangle \right\|_{\mathcal{F}} \leq \frac{C(1+t)\log^4(1+t)}{N^{(1-3\beta)/2}} . \quad (24)$$

provided $0 < \beta < \frac{1}{3}$.

The purpose of the present paper is to introduce and study a coupled refinement of the system (23a), (23h), (23i) which, we believe, is the correct system describing the case $\beta = 1$. These equations occur as Euler-Lagrange equations, and are written down explicitly in Theorem (8.1).

5. MAIN NEW RESULTS

Since \mathcal{H}_{red} is a fourth order polynomial in a and a^* ,

$$\mathcal{H}_{\text{red}}|0\rangle = (X_0, X_1, X_2, X_3, X_4, 0, \dots). \quad (25)$$

Definition 5.1. Define the Lagrangian

$$\mathcal{L} = - \int X_0(t) dt \quad (26)$$

The new, coupled equations for ϕ and k that we introduce in this paper are $X_1 = 0$ and $X_2 = 0$.

We first prove that \mathcal{L} is indeed the Lagrangian for these equations. We start by showing "abstractly" that

$$\frac{\delta X_0}{\delta \bar{\phi}} = \sqrt{N} \int (X_1(t, x) \text{ch}(k)(t, x, \cdot) - \bar{X}_1(t, x) \text{sh}(k)(t, x, y)) dx \quad (27)$$

$$\frac{\delta X_0}{\delta \bar{\zeta}} = \frac{1}{\sqrt{2}} \overline{\text{ch}(k)} \circ X_2 \circ \text{ch}(k) \quad (28)$$

where $\zeta = \text{th}(k) = \overline{\text{ch}(k)}^{-1} \circ \text{sh}(k)$. We then compute explicitly the zeroth order term $X_0(t)$ in $\mathcal{H}_{red}|0\rangle$ (which provides the Lagrangian density for our coupled equations):

$$\begin{aligned} -X_0(t) &= N \int dx_1 \left\{ -\Im(\phi_1 \overline{\partial_t \phi_1}) + |\nabla \phi_1|^2 \right\} \\ &+ \frac{N}{2} \int dx_1 dx_2 v_{1-2}^N |\phi_1 \phi_2 + \frac{1}{N} (\text{sh} \circ \text{ch})_{1,2}|^2 \\ &+ \frac{1}{2} \int dx_1 dx_2 dx_3 v_{1-2}^N |\phi_1 \text{sh}_{2,3} + \phi_2 \text{sh}_{1,3}|^2 \\ &+ \frac{1}{2} \left(\int dx_1 dx_2 \left\{ -\Im(\text{sh}_{1,2} \overline{\partial_t \text{sh}_{1,2}}) + |\nabla_{1,2} \text{sh}_{1,2}|^2 \right\} \right. \\ &\left. + \frac{1}{2N} \int dt dx_1 dx_2 v_{1-2}^N \left\{ |(\text{sh} \circ \bar{\text{sh}})_{1,2}|^2 + (\text{sh} \circ \bar{\text{sh}})_{1,1} (\bar{\text{sh}} \circ \text{sh})_{2,2} \right\} \right). \end{aligned}$$

where $\text{sh}_{1,2}$ is an abbreviation for $\text{sh}(k)(t, x_1, x_2)$, $v_{1-2}^N = v_N(x_1 - x_2)$, etc, and the products are pointwise products, while compositions are denoted by \circ . Then we proceed to compute explicitly the coupled equations $X_1 = 0$ and $X_2 = 0$, derive conserved quantities, and formulate a conjecture. The resulting equations are similar to those of Theorem (4.1), except that $m = -v_N(x_1 - x_2)\phi(t, x_1)\phi(t, x_2)$ is replaced by

$$\Theta = -v_N(x_1 - x_2) \left(\phi(t, x_1)\phi(t, x_2) + \frac{1}{2N} \text{sh}(2k)(t, x_1, x_2) \right),$$

and similar $O(\frac{1}{N})$ coupling corrections apply to the Hartree operator as well as \mathbf{S} and \mathbf{W} .

Remark 5.2. The static terms of $X_0(t)$ (not involving time derivatives) also appear in the recent preprint [1], but do not serve as a Lagrangian there.

6. THE LAGRANGIAN AND THE EQUATIONS, ABSTRACT
FORMULATION

Proposition 6.1. *Let k and ϕ be fixed.*

$$\begin{aligned} & \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} X_0(\phi + \epsilon h, k) \\ &= 2\sqrt{N}\Re \int X_1(t, x) \left(\text{ch}(k)(t, x, y)\bar{h}(t, y) - \overline{\text{sh}(k)}(t, x, y)h(t, y) \right) dx dy \end{aligned} \quad (29)$$

In particular, if this vanishes for all h , then $X_1(t, x) = 0$.

Proof. \mathcal{H}_{red} can be written as

$$\mathcal{H}_{red} = \frac{1}{i} \frac{\partial}{\partial t} + e^{\mathcal{B}} e^{\sqrt{N}\mathcal{A}} \left(-\frac{1}{i} \frac{\partial}{\partial t} + \mathcal{H} \right) \circ e^{-\sqrt{N}\mathcal{A}} e^{-\mathcal{B}}$$

in the sense of compositions (in space only) of operators. During this proof, denote $\mathcal{H}_t = -\frac{1}{i} \frac{\partial}{\partial t} + \mathcal{H}$.

Let h be an L^2 function and let $\mathcal{A}_\epsilon = \sqrt{N}(a(\bar{\phi} + \epsilon\bar{h}) - a^*(\phi + \epsilon h))$. Thus we have

$$X_0(\phi + \epsilon h, k) = \left\langle e^{\mathcal{B}} e^{\mathcal{A}_\epsilon} H_t e^{-\mathcal{A}_\epsilon} e^{-\mathcal{B}} |0\rangle, |0\rangle \right\rangle$$

We compute

$$\begin{aligned} & \left(\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} e^{\mathcal{A}_\epsilon} \right) e^{-\mathcal{A}_0} = \dot{\mathcal{A}}_0 + \frac{1}{2} [\mathcal{A}_0, \dot{\mathcal{A}}_0] \\ &= \sqrt{N} (a(\bar{h}) - a^*(h)) + \frac{N}{2} [a(\bar{\phi}) - a^*(\phi), a(\bar{h}) - a^*(h)] \\ &= \sqrt{N} (a(\bar{h}) - a^*(h)) + iN\Im \int \phi \bar{h} \end{aligned}$$

and

$$e^{\mathcal{A}_0} \left(\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} e^{-\mathcal{A}_\epsilon} \right) = - \left(\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} e^{\mathcal{A}_\epsilon} \right) e^{-\mathcal{A}_0}$$

thus

$$\begin{aligned}
& \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} \left\langle e^{\mathcal{B}} e^{\mathcal{A}_\epsilon} \mathcal{H}_t e^{-\mathcal{A}_\epsilon} e^{-\mathcal{B}} |0\rangle, |0\rangle \right\rangle \\
&= \left\langle e^{\mathcal{B}} \left[\sqrt{N} a(\bar{h}) - \sqrt{N} a^*(h), e^{\mathcal{A}_0} \mathcal{H}_t e^{-\mathcal{A}_0} \right] e^{-\mathcal{B}} |0\rangle, |0\rangle \right\rangle \\
&= \left\langle \left[e^{\mathcal{B}} \left(\sqrt{N} a(\bar{h}) - \sqrt{N} a^*(h) \right) e^{-\mathcal{B}}, e^{\mathcal{B}} e^{\mathcal{A}_0} \mathcal{H}_t e^{-\mathcal{A}_0} e^{-\mathcal{B}} \right] |0\rangle, |0\rangle \right\rangle \\
&= \left\langle \left[a(\bar{l}) - a^*(l), e^{\mathcal{B}} e^{\mathcal{A}_0} \mathcal{H}_t e^{-\mathcal{A}_0} e^{-\mathcal{B}} \right] |0\rangle, |0\rangle \right\rangle \\
&= 2\Re \left\langle \mathcal{H}_{red} |0\rangle, a^*(l) |0\rangle \right\rangle := I
\end{aligned}$$

where we denoted

$$e^{\mathcal{B}} \left(\sqrt{N} a(\bar{h}) - \sqrt{N} a^*(h) \right) e^{-\mathcal{B}} = a(\bar{l}) - a^*(l)$$

Explicitly,

$$\begin{aligned}
& e^{\mathcal{B}} \left(a(\bar{h}) - a^*(h) \right) e^{-\mathcal{B}} \\
&= a(\text{ch}(k) \circ \bar{h}) + a^*(\text{sh}(k) \circ \bar{h}) \\
&\quad - a(\overline{\text{sh}(k)} \circ h) - a^*(\overline{\text{ch}(k)} \circ h)
\end{aligned}$$

so

$$l = \sqrt{N} \left(\overline{\text{ch}(k)} \circ h - \text{sh}(k) \circ \bar{h} \right)$$

Thus,

$$\begin{aligned}
I &= 2\sqrt{N} \Re \int X_1(t, x) \left(\text{ch}(k)(t, x, y) \bar{h}(y) - \overline{\text{sh}(k)}(t, x, y) h(y) \right) dx dy \\
&= 2\sqrt{N} \Re \int \left(\overline{\text{ch}(k)} \circ X_1 - \text{sh}(k) \circ \overline{X_1} \right) (y) \bar{h}(y) dy
\end{aligned}$$

□

In order to state the corresponding result for X_2 , we have to introduce a new set of coordinates for our basic matrices

$$e^K = \begin{pmatrix} \text{ch}(k) & \overline{\text{sh}(k)} \\ \text{sh}(k) & \overline{\text{ch}(k)} \end{pmatrix}$$

where

$$K = \begin{pmatrix} 0 & \bar{k}(t, x, y) \\ k(t, x, y) & 0 \end{pmatrix} \quad (30)$$

The most obvious coordinate system is, of course, provided by k . We recall the following proposition, proved in [18].

Proposition 6.2. *The exponential map is one-to-one and onto from matrices of the form (30) ($k \in L^2$, symmetric) to positive definite matrices E satisfying the three properties of Remark (3.1) for which $\|I - E\|_{L^2}$ is finite.*

For our purposes, a better coordinate system is provided by $\zeta = \text{th}(k) = \overline{\text{ch}(k)}^{-1} \circ \text{sh}(k)$.

Proposition 6.3. *There is a bijection between $k \in L^2$, symmetric, and $\zeta \in L^2$, symmetric, $\|\zeta\|_{op} < 1$ (op stands for the operator norm) such that*

$$\begin{aligned} e^K &:= \begin{pmatrix} \text{ch}(k) & \overline{\text{sh}(k)} \\ \text{sh}(k) & \text{ch}(k) \end{pmatrix} \\ &= E_\zeta := \begin{pmatrix} I & \bar{\zeta} \\ 0 & I \end{pmatrix} \begin{pmatrix} (I - \bar{\zeta} \circ \zeta)^{1/2} & 0 \\ 0 & (I - \zeta \circ \bar{\zeta})^{-1/2} \end{pmatrix} \begin{pmatrix} I & 0 \\ \zeta & I \end{pmatrix} \\ &= \begin{pmatrix} (I - \bar{\zeta} \circ \zeta)^{-1/2} & \bar{\zeta} \circ (I - \zeta \circ \bar{\zeta})^{-1/2} \\ \zeta \circ (I - \bar{\zeta} \circ \zeta)^{-1/2} & (I - \zeta \circ \bar{\zeta})^{-1/2} \end{pmatrix} \end{aligned} \quad (31)$$

where the square root is taken in the operator sense.

Proof. Given k , define $\zeta = \overline{\text{ch}(k)}^{-1} \circ \text{sh}(k)$. The decomposition (31) is an algebraic identity, and it is clear that ζ is symmetric and L^2 . Since $I - \text{ch}(k)^{-2} = \bar{\zeta} \circ \zeta$, we see that $\|\zeta\|_{op} < 1$. In fact, $\|\zeta v\|_{L^2}^2 = \|v\|_{L^2}^2 - \|\text{ch}(k)^{-1}v\|_{L^2}^2$. Conversely, given ζ a symmetric Hilbert-Schmidt kernel with $\|\zeta\|_{op} < 1$ define E_ζ by (31). It is easy to check that E_ζ is positive definite, satisfies the symmetries of remark (3.1) and $\|I - E_\zeta\|_{HS} < \infty$. (HS stands for the Hilbert-Schmidt norm), thus we can apply Proposition (6.2) and find the corresponding K . \square

We also record the following consequence:

Proposition 6.4. *Let $\zeta_\epsilon = \zeta + \epsilon h$ ($h \in L^2$, symmetric, $\|\zeta_\epsilon\|_{op} < 1$), and K_ϵ corresponding to ζ_ϵ according to the previous proposition. Then*

$$\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} e^{K_\epsilon} e^{-K} = \begin{pmatrix} ia & \bar{b} \\ b & -ia^T \end{pmatrix}$$

with

$$b = \overline{\text{ch}(k)} \circ h \circ \text{ch}(k)$$

Proof. We compute

$$\begin{aligned} & \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} e^{K_\epsilon} e^{-K} \\ &= \begin{pmatrix} \text{ch}(k)' \circ \text{ch}(k) - \overline{\text{sh}(k)'} \circ \text{sh}(k) & -\text{ch}(k)' \circ \overline{\text{sh}(k)} + \overline{\text{sh}(k)'} \circ \overline{\text{ch}(k)} \\ \text{sh}(k)' \circ \text{ch}(k) - \overline{\text{ch}(k)'} \circ \text{sh}(k) & -\text{sh}(k)' \circ \overline{\text{sh}(k)} + \overline{\text{ch}(k)'} \circ \overline{\text{ch}(k)} \end{pmatrix} \end{aligned}$$

An easy calculation shows that $b = -\overline{\text{ch}(k)'} \circ \text{sh}(k) + \text{sh}(k)' \circ \text{ch}(k) = \overline{\text{ch}(k)} \circ \zeta' \circ \text{ch}(k)$. \square

We are ready to prove

$$\frac{\delta X_0}{\delta \zeta} = \frac{1}{\sqrt{2}} \overline{\text{ch}(k)} \circ X_2 \circ \text{ch}(k)$$

Proposition 6.5. *Let k_ϵ correspond to $\zeta + \epsilon h$ as in the previous proposition. Then*

$$\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} X_0(\phi, k_\epsilon) = \sqrt{2} \Re \int \overline{\text{ch}(k)} \circ X_2 \circ \text{ch}(k)(t, z, w) \bar{h}(t, z, w) dz dw$$

In particular, if the above vanishes for all h , then $X_2 = 0$.

Proof. Let $B_\epsilon = B(k_\epsilon)$.

$$X_0(\phi, k_\epsilon) = \left\langle e^{\mathcal{B}_\epsilon} e^{\sqrt{N}\mathcal{A}} H_t e^{-\sqrt{N}\mathcal{A}} e^{-\mathcal{B}_\epsilon} |0\rangle, |0\rangle \right\rangle$$

and

$$\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} X_0(\phi, k_\epsilon) = -2\Re \left\langle H_{red} |0\rangle, \psi |0\rangle \right\rangle \quad (32)$$

where

$$\psi = \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} e^{\mathcal{B}_\epsilon} e^{-B} = \mathcal{I} \left(\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} e^{K_\epsilon} e^{-K} \right)$$

Using the isomorphism (7) and proposition (6.4) we see that

$$\psi |0\rangle = (i\theta, 0, -\frac{1}{\sqrt{2}} \overline{\text{ch}(k)} \circ h \circ \text{ch}(k)(t, x_1, x_2), 0, \dots)$$

where θ is a real number coming from the trace of the self-adjoint a . Since X_0 is real, $i\theta$ does not contribute to (32), and the result follows. \square

7. EXPLICIT FORM OF THE LAGRANGIAN

The goal of this section is the following proposition.

Proposition 7.1. *The zeroth order term in $\mathcal{H}_{red}|0\rangle$ (which provides the Lagrangian density for our coupled equations) is $X_0(t)$ where*

$$\begin{aligned}
-X_0(t) = & N \int dx_1 \left\{ -\Im(\phi_1 \overline{\partial_t \phi_1}) + |\nabla \phi_1|^2 \right\} \\
& + \frac{N}{2} \int dx_1 dx_2 v_{1-2}^N |\phi_1 \phi_2 + \frac{1}{N} (\text{sh} \circ \text{ch})_{1,2}|^2 \\
& + \frac{1}{2} \int dx_1 dx_2 dx_3 v_{1-2}^N |\phi_1 \text{sh}_{2,3} + \phi_2 \text{sh}_{1,3}|^2 \\
& + \frac{1}{2} \left(\int dx_1 dx_2 \left\{ -\Im(\text{sh}_{1,2} \overline{\partial_t \text{sh}_{1,2}}) + |\nabla_{1,2} \text{sh}_{1,2}|^2 \right\} \right. \\
& \left. + \frac{1}{2N} \int dt dx_1 dx_2 v_{1-2}^N \left\{ |(\text{sh} \circ \overline{\text{sh}})_{1,2}|^2 + (\text{sh} \circ \overline{\text{sh}})_{1,1} (\overline{\text{sh}} \circ \text{sh})_{2,2} \right\} \right).
\end{aligned}$$

where $\text{sh}_{1,2}$ is an abbreviation for $\text{sh}(k)(t, x_1, x_2)$, etc, and the products are pointwise products, while compositions are denoted by \circ .

The proof follows from several lemmas, which can be proved by explicit calculations. We proceed to compute X_0 in (25). The only terms in (17) which contribute to X_0 are $N\mathcal{P}_0$ which is already explicit, the zeroth order terms in $\mathcal{I}(R)|0\rangle$, as well as the zeroth order terms in $N^{-1}e^{\mathcal{B}}\mathcal{P}_4e^{-\mathcal{B}}|0\rangle$.

Lemma 7.2. *The term $N\mathcal{P}_0$ is given by*

$$\begin{aligned}
N\mathcal{P}_0 = & N \int dx \left\{ \frac{1}{2i} (\phi \bar{\phi}_t - \bar{\phi} \phi_t) - |\nabla \phi|^2 \right\} \\
& - \frac{N}{2} \int dx_1 dx_2 \left\{ v_{1-2}^N |\phi_1 \phi_2|^2 \right\}.
\end{aligned}$$

We used abbreviations $v_{1-2}^N = v_N(x_1 - x_2)$, $\phi_1 = \phi(x_1)$, etc., and for the following two lemmas we will denote $u_{1,2} = \text{sh}(k)(t, x_1, x_2)$ and $c_{1,2} = \text{ch}(k)(t, x_1, x_2)$.

Lemma 7.3. *The zeroth order term in $\mathcal{I}(R)|0\rangle$ is*

$$\begin{aligned} & -\frac{1}{2} \left(\int dx_1 dx_2 \left\{ \frac{1}{2i} (\bar{u}_{1,2} \partial_t u_{1,2} - \partial_t \bar{u}_{1,2} u_{1,2}) + |\nabla_{1,2} u_{1,2}|^2 \right\} \right. \\ & + \int dx_1 dx_2 dx_3 \{ v_{1-2}^N |\phi_1 u_{2,3}|^2 + |\phi_2 u_{1,3}|^2 \} \\ & + 2\Re \int dx_1 dx_2 dx_3 \{ v_{1-2}^N \phi_2 u_{1,3} \overline{\phi_1 u_{2,3}} \} \\ & \left. + 2\Re \int dx_1 dx_2 \{ v_{1-2}^N (u \circ c)_{1,2} \bar{\phi}_1 \bar{\phi}_2 \} \right) \end{aligned}$$

Lemma 7.4. *The zeroth order term in $-\frac{1}{N} e^{\mathcal{B}} \mathcal{V} e^{-\mathcal{B}} |0\rangle$ is*

$$\begin{aligned} & -\frac{1}{2} \int dx_1 dx_2 v_{1-2}^N \left\{ (u \circ c)_{1,2} \overline{(u \circ c)_{1,2}} \right. \\ & \left. + |(u \circ \bar{u})_{1,2}|^2 + (u \circ \bar{u})_{1,1} (\bar{u} \circ u)_{2,2} \right\} . \end{aligned}$$

8. EXPLICIT FORM OF THE EQUATIONS

In this section we derive the following theorem, thus introducing our new equations. First, some notation. Consider the kernels

$$\begin{aligned} \omega_c(t, x, y) &= \bar{\phi}(t, x) \phi(t, y) \\ \omega_p(t, x, y) &= \overline{\text{sh}(k)} \circ \text{sh}(k)(t, x, y) \end{aligned}$$

and their trace densities

$$\begin{aligned} \rho_c &= |\phi|^2(t, x) \\ \rho_p(t, x) &= \text{sh}(k) \circ \overline{\text{sh}(k)}(t, x, x) \end{aligned}$$

Here c stands for condensate, and p for pair. In this notation, the old operator kernel g_N defined in (20c) is

$$\begin{aligned} g_N(t, x, y) &:= -\Delta_x \delta(x - y) + (v_N * \rho_c)(t, x) \delta(x - y) \\ &+ v_N(x - y) \omega_c(t, x, y) \end{aligned}$$

Define the new operator kernel

$$\begin{aligned} \tilde{g}_N(t, x, y) &:= -\Delta_x \delta(x - y) \\ &+ (v_N * \rho_c)(t, x) \delta(x - y) + v_N(x - y) \omega_c(t, x, y) \end{aligned} \quad (33)$$

$$+ \frac{1}{N} ((v_N * \rho_p)(t, x) \delta(x - y) + v_N(x - y) \omega_p(t, x, y)) \quad (34)$$

Also denote $\alpha_c = (33)$, $\frac{1}{N}\alpha_p = (34)$ and $\alpha = \alpha_c + \frac{1}{N}\alpha_p$. Define

$$\tilde{\mathbf{S}}(s) := \frac{1}{i}s_t + \tilde{g}_N^T \circ s + s \circ \tilde{g}_N \quad \text{and} \quad \tilde{\mathbf{W}}(p) := \frac{1}{i}p_t + [\tilde{g}_N^T, p]$$

Finally, define $\Theta(t, x_1, x_2) = -v_N(t, x_1, x_2) (\phi(t, x_1)\phi(t, x_2) + \frac{1}{2N}\text{sh}(2k)(t, x_1, x_2))$.

Theorem 8.1. *The equation $X_1 = 0$ is equivalent to*

$$\frac{1}{i}\partial_t\phi(t, x_1) - \Delta\phi - \int \Theta(t, x_1, x_2)\bar{\phi}(t, x_2)dx_2 + \int \frac{1}{N}\alpha_p^T(t, x_1, x_2)\phi(t, x_2)dx_2 = 0$$

The equation $X_2 = 0$ is equivalent to either of :

1) *the equation*

$$\tilde{\mathbf{S}}(\text{th}(k)) = \Theta + \text{th}(k) \circ \bar{\Theta} \circ \text{th}(k)$$

2) *the pair of equations (in fact, 2a) implies 2b))*

$$2a) \tilde{\mathbf{S}}(\text{sh}(2k)) = \Theta \circ \text{ch}(2k) + \overline{\text{ch}(2k)} \circ \Theta \quad (35)$$

$$2b) \tilde{\mathbf{W}}(\overline{\text{ch}(2k)}) = \Theta \circ \overline{\text{sh}(2k)} - \text{sh}(2k) \circ \bar{\Theta}$$

Remark 8.2. One can go back and fourth between ζ and $\text{ch}(2k)$, $\text{sh}(2k)$ using

$$\begin{aligned} \overline{\text{sh}(k)} \circ \text{sh}(k) &= (1 - \bar{\zeta} \circ \zeta)^{-1} - 1 = \frac{1}{2}(\text{ch}(2k) - 1) \\ \zeta &= \text{sh}(2k)(1 + \text{ch}(2k))^{-1} \end{aligned}$$

Proof. A direct calculation for X_1 shows that

$$X_1 = -\sqrt{N} \left(\overline{\text{ch}(k)} \circ \widetilde{\text{Har}}_k(\phi) + \text{sh}(k) \circ \overline{\widetilde{\text{Har}}_k(\phi)} \right)$$

where

$$\begin{aligned} &\widetilde{\text{Har}}_k(\phi)(t, x_1) \\ &= \frac{1}{i}\partial_t\phi - \Delta\phi - \int \Theta(t, x_1, x_2)\bar{\phi}(t, x_2)dx_2 \\ &+ \frac{1}{N} \int v_N(x_1 - x_2)(\text{sh} \circ \bar{\text{sh}})(x_1, x_2)\phi(x_2)dx_2 \\ &+ \frac{1}{N} \phi(x_1) \int v_N(x_1 - x_2)(\text{sh} \circ \bar{\text{sh}})(x_2, x_2)dx_2 \end{aligned}$$

In conjunction with Proposition (6.1) this shows that

$$\frac{\delta\mathcal{L}}{\delta\bar{\phi}} = N\widetilde{\text{Har}}_k(\phi)$$

which can also be easily verified directly from Proposition (7.1).

A direct calculation also shows that, if X_2 denotes the second component of $\mathcal{H}_{red}|0\rangle$, then

$$\begin{aligned}
& -\sqrt{2}X_2(t, y_1, y_2) = \tag{36} \\
& \left(\left(\mathbf{S}(\text{sh}(k)) - \overline{\text{ch}(k)} \circ m \right) \circ \text{ch}(k) - \left(\mathbf{W}(\overline{\text{ch}(k)}) + \text{sh}(k) \circ \overline{m} \right) \circ \text{sh}(k) \right) \\
& + (1/N) \int dx_1 dx_2 \quad \left\{ \right. \\
& \left(\overline{\text{ch}}(y_1, x_2) \text{sh}(x_2, y_2) (\overline{\text{sh}} \circ \text{sh})(x_1, x_1) v_N(x_1 - x_2) + \right. \\
& \overline{\text{ch}}(y_1, x_2) \text{sh}(x_1, y_2) (\overline{\text{sh}} \circ \text{sh})(x_1, x_2) v_N(x_1 - x_2) + \\
& \overline{\text{ch}}(y_1, x_1) \text{sh}(x_2, y_2) (\text{sh} \circ \overline{\text{sh}})(x_1, x_2) v_N(x_1 - x_2) + \\
& \left. \overline{\text{ch}}(y_1, x_1) \text{sh}(x_1, y_2) (\text{sh} \circ \overline{\text{sh}})(x_2, x_2) v_N(x_1 - x_2) \right)_{\text{symm}} + \\
& \text{sh}(y_1, x_1) \text{sh}(x_2, y_2) (\overline{\text{sh}} \circ \overline{\text{ch}})(x_1, x_2) v_N(x_1 - x_2) + \\
& \left. \overline{\text{ch}}(y_1, x_1) \text{ch}(x_2, y_2) (\overline{\text{ch}} \circ \text{sh})(x_1, x_2) v_N(x_1 - x_2) \right\}.
\end{aligned}$$

where *symm* stands for "symmetrized". The time dependence in the last six lines has been omitted. Recalling $\zeta = \overline{\text{ch}(k)}^{-1} \circ \text{sh}(k) = \text{sh}(k) \circ \overline{\text{ch}(k)}^{-1}$, compose on the left with $\overline{\text{ch}(k)}^{-1}$ and on the right with $\text{ch}(k)^{-1}$ to get

$$\overline{\text{ch}(k)}^{-1} \circ X_2 \circ \text{ch}(k)^{-1} = \mathbf{S}(\zeta) - \Theta - \zeta \circ \overline{\Theta} \circ \zeta + \frac{1}{N} \mathbf{N} \tag{37}$$

where N is given by

$$\begin{aligned}
\mathbf{N}(t, y_1, y_2) = & \zeta(t, y_1, y_2) \left(\int dx \left((\overline{\text{sh}} \circ \text{sh} + \text{sh} \circ \overline{\text{sh}})(t, x, x) v_N(x - y_1) \right) \right)_{\text{symm}} + \\
& \left(\int dx \zeta(t, x, y_2) (\overline{\text{sh}} \circ \text{sh} + \text{sh} \circ \overline{\text{sh}})(t, x, y_1) v_N(x - y_1) \right)_{\text{symm}}
\end{aligned}$$

where *symm* stands for symmetrizing in y_1, y_2 . In other words,

$$\mathbf{N} = \zeta \circ \alpha_p + \alpha_p \circ \zeta$$

Thus, in ζ coordinates, the equation $X_2 = 0$ becomes

$$\tilde{\mathbf{S}}(\zeta) - \Theta - \zeta \circ \overline{\Theta} \circ \zeta = 0 \tag{38}$$

Now we can get an equation for $\tilde{\mathbf{W}}(\overline{\text{ch}(2k)})$ and $\tilde{\mathbf{S}}(\text{sh}(2k))$. We will use the general formulas

$$\begin{aligned}\tilde{\mathbf{W}}(f^{-1}) &= -f^{-1} \circ \tilde{\mathbf{W}}(f) \circ f^{-1} \\ \tilde{\mathbf{W}}(f \circ \bar{g}) &= \tilde{\mathbf{S}}(f) \circ \bar{g} - f \circ \overline{\tilde{\mathbf{S}}(g)} \\ \tilde{\mathbf{S}}(f \circ g) &= \tilde{\mathbf{S}}(f) \circ g - f \circ \overline{\tilde{\mathbf{W}}(\bar{g})}\end{aligned}$$

Thus

$$\begin{aligned}\tilde{\mathbf{W}}((1 - \zeta \circ \bar{\zeta})^{-1}) &= (1 - \zeta \circ \bar{\zeta})^{-1} \circ \left(\tilde{\mathbf{S}}(\zeta) \circ \bar{\zeta} - \zeta \circ \overline{\tilde{\mathbf{S}}(\zeta)} \right) \circ (1 - \zeta \circ \bar{\zeta})^{-1} \\ &= (1 - \zeta \circ \bar{\zeta})^{-1} \circ \left((\Theta + \zeta \circ \bar{\Theta} \circ \zeta) \bar{\zeta} + \zeta \circ \overline{(\Theta + \zeta \circ \bar{\Theta} \circ \zeta)} \right) (1 - \zeta \circ \bar{\zeta})^{-1}\end{aligned}$$

Similarly we get a formula for $\tilde{\mathbf{S}}(\text{sh}(2k))$, using

$$\begin{aligned}\tilde{\mathbf{S}}(\zeta \circ (1 - \bar{\zeta} \circ \zeta)^{-1}) &= (1 - \zeta \circ \bar{\zeta})^{-1} \circ \left(\tilde{\mathbf{S}}(\zeta) - \zeta \circ \overline{\tilde{\mathbf{S}}(\zeta)} \circ \zeta \right) \circ (1 - \bar{\zeta} \circ \zeta)^{-1}\end{aligned}$$

$$\begin{aligned}\tilde{\mathbf{S}}(\zeta \circ (1 - \bar{\zeta} \circ \zeta)^{-1}) &= (1 - \zeta \circ \bar{\zeta})^{-1} \circ \left(\Theta + \zeta \circ \bar{\Theta} \circ \zeta + \zeta \circ \overline{(\Theta + \zeta \circ \bar{\Theta} \circ \zeta)} \circ \zeta \right) \circ (1 - \bar{\zeta} \circ \zeta)^{-1} \\ &= \left((1 - \zeta \circ \bar{\zeta})^{-1} - \frac{1}{2} \right) \circ \Theta + \Theta \circ \left((1 - \bar{\zeta} \circ \zeta)^{-1} - \frac{1}{2} \right)\end{aligned}$$

□

9. CONSERVED QUANTITIES

We start by motivating the introduction of some conserved quantities. Recall the Lagrangian

$$\begin{aligned}
\mathcal{L}(\phi, \text{sh}(k)) &= N \int dt dx_1 \left\{ -\Im (\phi_1 \overline{\partial_t \phi_1}) + |\nabla \phi_1|^2 \right\} \\
&+ \frac{N}{2} \int dt dx_1 dx_2 v_{1-2}^N |\phi_1 \phi_2 + \frac{1}{N} (\text{sh} \circ \text{ch})_{1,2}|^2 \\
&+ \frac{1}{2} \int dt dx_1 dx_2 dx_3 v_{1-2}^N |\phi_1 \text{sh}_{2,3} + \phi_2 \text{sh}_{1,3}|^2 \\
&+ \frac{1}{2} \left(\int dt dx_1 dx_2 \left\{ -\Im (\text{sh}_{1,2} \overline{\partial_t \text{sh}_{1,2}}) + |\nabla_{1,2} \text{sh}_{1,2}|^2 \right\} \right. \\
&\left. + \frac{1}{2N} \int dt dx_1 dx_2 v_{1-2}^N \left\{ |(\text{sh} \circ \overline{\text{sh}})_{1,2}|^2 + (\text{sh} \circ \overline{\text{sh}})_{1,1} (\overline{\text{sh}} \circ \text{sh})_{2,2} \right\} \right).
\end{aligned}$$

where $\text{sh}_{1,2}$ is an abbreviation for $\text{sh}(k)(t, x_1, x_2)$, etc, and the products are pointwise products, while compositions are denoted by \circ . Introduce the energy \mathcal{E}

$$\begin{aligned}
\mathcal{E}(\phi, \text{sh}(k))(t) &= N \int dx_1 \left\{ |\nabla \phi_1|^2 \right\} \\
&+ \frac{N}{2} \int dx_1 dx_2 v_{1-2}^N |\phi_1 \phi_2 + \frac{1}{N} (\text{sh} \circ \text{ch})_{1,2}|^2 \\
&+ \frac{1}{2} \int dx_1 dx_2 dx_3 v_{1-2}^N |\phi_1 \text{sh}_{2,3} + \phi_2 \text{sh}_{1,3}|^2 \\
&+ \frac{1}{2} \left(\int dx_1 dx_2 \left\{ |\nabla_{1,2} \text{sh}_{1,2}|^2 \right\} \right. \\
&\left. + \frac{1}{2N} \int dx_1 dx_2 v_{1-2}^N \left\{ |(\text{sh} \circ \overline{\text{sh}})_{1,2}|^2 + (\text{sh} \circ \overline{\text{sh}})_{1,1} (\overline{\text{sh}} \circ \text{sh})_{2,2} \right\} \right).
\end{aligned}$$

Our equations for ϕ and $\text{sh}(k)$ are equivalent to

$$N \frac{1}{i} \frac{\partial \phi}{\partial t} = - \frac{\delta \mathcal{E}}{\delta \overline{\phi}} \quad (39)$$

$$\frac{1}{i} \frac{\partial \text{sh}(k)}{\partial t} = - \frac{\delta \mathcal{E}}{\delta \overline{\text{sh}(k)}} \quad (40)$$

The relation

$$\begin{aligned} 0 &= \frac{d}{d\theta} \Big|_{\theta=0} \mathcal{E}(e^{i\theta}\phi, e^{2i\theta}\text{sh}(k)) \\ &= 2\Re \left(\int \frac{\delta\mathcal{E}}{\delta\bar{\phi}}(-i\bar{\phi})dx_1 + \int \frac{\delta\mathcal{E}}{\delta\text{sh}(k)}(-i\overline{\text{sh}(k)})dx_1dx_2 \right) \end{aligned}$$

together with (39), (40), leads to the conservation

$$\frac{d}{dt} \left(\int |\phi(t, x_1)|^2 dx_1 + \frac{1}{N} \int |\text{sh}(k)(t, x_1, x_2)|^2 dx_1 dx_2 \right) = 0$$

thus we define the density

$$\begin{aligned} \rho(t, x_1) &= |\phi(t, x_1)|^2 + \frac{1}{N} \int |\text{sh}(k)(t, x_1, x_2)|^2 dx_2 \\ &= \rho_c(t, x_1) + \frac{1}{N} \rho_p(t, x_1) \end{aligned}$$

Similarly, let $\phi_\epsilon(t, x) = \phi(t, x + \epsilon e_j)$, $\text{sh}(k)_\epsilon(t, x, y) = \text{sh}(k)(t, x + \epsilon e_j, y + \epsilon e_j)$ ($e_j =$ unit vector, $1 \leq j \leq 3$). The relation

$$\begin{aligned} 0 &= \frac{d}{d\epsilon} \Big|_{\epsilon=0} \mathcal{E}(\phi_\epsilon, \text{sh}(k)_\epsilon) \\ &= 2\Re \left(\int \frac{\delta\mathcal{E}}{\delta\bar{\phi}} \partial_j \bar{\phi} dx_1 + \int \frac{\delta\mathcal{E}}{\delta\text{sh}(k)} (\partial_j \overline{\text{sh}(k)}) dx_1 dx_2 \right) \end{aligned}$$

together with (39), (40) leads to the conservation

$$\frac{d}{dt} \left(N \int \Im(\phi \partial_j \bar{\phi}) dx_1 + \int \Im(\text{sh}(k) \partial_j \overline{\text{sh}(k)}) dx_1 dx_2 \right) = 0$$

thus we define the momentum density

$$\begin{aligned} p_j(t, x_1) &= \Im(\phi \partial_j \bar{\phi}) + \frac{1}{N} \int \Im(\text{sh}(k) \partial_j \overline{\text{sh}(k)}) dx_2 \\ &:= p_{c,j}(t, x_1) + \frac{1}{N} p_{p,j}(t, x_1) \end{aligned}$$

Finally, using (39), (40) we see that

$$\frac{\partial}{\partial t} \mathcal{E}(t) = 0$$

so we define the energy density

$$\begin{aligned}
e(t, x_1) &= N |\nabla \phi_1|^2 \\
&+ \frac{N}{2} \int dx_2 v_{1-2}^N |\phi_1 \phi_2 + \frac{1}{N} (\text{sh} \circ \text{ch})_{1,2}|^2 \\
&+ \frac{1}{2} \int dx_2 dx_3 v_{1-2}^N |\phi_1 \text{sh}_{2,3} + \phi_2 \text{sh}_{1,3}|^2 \\
&+ \frac{1}{2} \left(\int dx_2 \left\{ |\nabla_{1,2} \text{sh}_{1,2}|^2 \right\} \right. \\
&\left. + \frac{1}{2N} \int dx_2 v_{1-2}^N \left\{ |(\text{sh} \circ \overline{\text{sh}})_{1,2}|^2 + (\text{sh} \circ \overline{\text{sh}})_{1,1} (\overline{\text{sh}} \circ \text{sh})_{2,2} \right\} \right).
\end{aligned}$$

10. A CONJECTURE

We conjecture that, if ϕ , k satisfy the equations of Theorem (8.1) and $|\psi_{exact}\rangle$, $|\psi_{appr}\rangle$, are defined by (14), (15), then, in the critical case $\beta = 1$,

$$\| |\psi_{exact}\rangle - |\psi_{appr}\rangle \|_{\mathcal{F}} \rightarrow 0$$

as $N \rightarrow \infty$, at an explicit rate.

REFERENCES

- [1] Benedikter, N., de Oliveira, G., Schlein B. : Quantitative Derivation of the Gross-Pitaevskii Equation, preprint.
- [2] Bogoliubov, N. N.: Lectures on Quantum Statistics: Quasi-Averages, Gordon and Breach (1970)
- [3] Chen, X. : On the Rigorous Derivation of the 3D Cubic Nonlinear Schrödinger Equation with A Quadratic Trap, Archive for Rational Mechanics and Analysis, **210** (2013), 365-408
- [4] Chen, X, Holmer, J. : On the Klainerman-Machedon Conjecture of the Quantum BBGKY Hierarchy with Self-interaction, preprint.
- [5] Chen, X, Holmer, J. : Focusing Quantum Many-body Dynamics: The Rigorous Derivation of the 1D Focusing Cubic Nonlinear Schrödinger Equation, preprint.
- [6] Chen, T. and Pavlović, N. : Derivation of the cubic NLS and Gross-Pitaevskii hierarchy for manybody dynamics in $d = 2, 3$ based on spacetime estimates, preprint.
- [7] Elgart, A., Erdős, L., Schlein, B., Yau, H. T.: Gross-Pitaevskii equation as the mean field limit of weakly coupled bosons. Arch. Rat. Mech. Anal. **179**, 265–283 (2006)
- [8] Erdős, L., Yau, H. T.: Derivation of the non-linear Schrödinger equation from a many-body Coulomb system. Adv. Theor. Math. Phys. **5**, 1169–1205 (2001)
- [9] Erdős, L., Schlein, B., Yau, H. T.: Derivation of the Gross-Pitaevskii hierarchy for the dynamics of Bose-Einstein condensate. Comm. Pure Appl. Math. **59**, 1659–1741 (2006)

- [10] Erdős, L., Schlein, B., Yau, H. T.: Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems. *Invent. Math.* **167**, 515–614 (2007)
- [11] Erdős, L., Schlein, B., Yau, H. T.: Rigorous derivation of the Gross-Pitaevskii equation. *Phys. Rev. Lett.* **98**, 040404 (2007)
- [12] Erdős, L., Schlein, B., Yau, H. T.: Derivation of the Gross-Pitaevskii equation for the dynamics of Bose-Einstein condensate. *Annals Math.* **172**, 291–370 (2010)
- [13] Folland, G. B.: *Harmonic analysis in phase space*. *Annals of Math. Studies*, Vol. **122**. Princeton, NJ: Princeton University Press, 1989
- [14] Frölich, J., Knowles, A., Schwarz, S., : On the mean-field limit of Bosons with Coulomb two-body interaction, *Comm. Math Phys.*, **288**, (209), 1023-1059.
- [15] Ginibre, J., Velo, G.: The classical field limit of scattering theory for non-relativistic many-boson systems, I and II. *Comm. Math. Phys.* **66**, 37–76 (1979) and **68**, 45–68 (1979)
- [16] Ginibre, J., Velo, G.: Scattering theory in the energy space for a class of nonlinear Schrödinger equations, *J. Math. Pures Appl.* **64** (1985), 363-401
- [17] Grillakis, M. Machedon, M., Margetis, D.: Second-order corrections to mean field evolution of weakly interacting Bosons. I. *Comm. Math. Phys.* **294**, 273–301 (2010)
- [18] Grillakis, M. Machedon, M., Margetis, D.: Second-order corrections to mean field evolution of weakly interacting Bosons. II. *Adv. in Math.* **228**, 1788-1815 (2011)
- [19] Grillakis, M. Machedon, M.: Pair excitations and the mean field approximation of interacting Bosons, I. To appear in *Communications in Mathematical Physics*.
- [20] Hepp, K.: The classical limit for quantum mechanical correlation functions. *Comm. Math. Phys.* **35**, 265–277 (1974)
- [21] Klainerman, S., Machedon, M. On the uniqueness of solutions to the Gross-Pitaevskii hierarchy. *Comm. Math. Phys.* **279**, 169-185 (2008)
- [22] Kirkpatrick, K., Schlein, B., Staffilani, G. :Derivation of the two dimensional nonlinear Schrödinger equation from manybody quantum dynamics, *Amer. J. Math.* **133** (2011), 91-130.
- [23] Lee, T. D., Huang, K., Yang, C. N.: Eigenvalues and eigenfunctions of a Bose system of hard spheres and its low-temperature properties. *Phys. Rev.* **106**, 1135–1145 (1957)
- [24] Lee, T. Yang, C. N.: Low-Temperature Behaviour of a Dilute System of Hard Spheres. I. Equilibrium Properties. *Phys. Rev.* **112**, 1419-1429 (1958)
- [25] Lieb, E. H., Seiringer, R., Solovej, J. P., Yngvanson, J.: *The mathematics of the Bose gas and its condensation*. Basel, Switzerland: Birkhäuser Verlag, 2005
- [26] Lieb, E. H., Seiringer: Proof of Bose-Einstein condensation for dilute trapped gasses, *Phys. Rev. Letters* **88** (2002), 170409.
- [27] Knowles, A., Pickl, P. : Mean-field dynamics: Singular potentials and rate of convergence, preprint.
- [28] Rodnianski, I., Schlein, B.: Quantum fluctuations and rate of convergence towards mean field dynamics. *Comm. Math. Phys.* **291**(2), 31–61 (2009)

- [29] Shale, D.: Linear symmetries of free Boson fields, *Trans. Amer. Math. Soc.* **103**(1), 149–167 (1962)
- [30] Spohn, H. : Kinetic equations from hamiltonian dynamics: Markovian limits. *Rev. Mod. Phys.* **52**, (1980), 569-615.
- [31] Wu, T. T.: Some nonequilibrium properties of a Bose system of hard spheres at extremely low temperatures. *J. Math. Phys.* **2**, 105–123 (1961)
- [32] Wu, T. T.: Bose-Einstein condensation in an external potential at zero temperature: General Theory. *Phys. Rev. A* **58**, 1465–1474 (1998)

UNIVERSITY OF MARYLAND, COLLEGE PARK
E-mail address: `mng@math.umd.edu`

UNIVERSITY OF MARYLAND, COLLEGE PARK
E-mail address: `mxm@math.umd.edu`