

Acknowledgments



This material is based upon work partially supported by the National Science Foundation under grant no. DMS-2510216 and by Simons Foundation. “Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.”

Collaborators:

Kasso Okoudjou (Tufts), Anirudha Poria (Bar-Ilan U.), Michael Rawson (UMD/PNNL/Microsoft), Yang Wang (HKUST), Rui Zhang (HKUST), Felix Kraemer (TUM).

Works:

- ① R. Balan, K. Okoudjou, A. Poria, *On a Feichtinger Problem*, Operators and Matrices vol. 12(3), 881-891 (2018) <http://dx.doi.org/10.7153/oam-2018-12-53>
- ② R. Balan, K.A. Okoudjou, M. Rawson, Y. Wang, R. Zhang, *Optimal l_1 Rank One Matrix Decomposition*, in "Harmonic Analysis and Applications", Rassias M., Ed. Springer (2021)
- ③ R. Balan, F. Jiang, *Factorization of positive-semidefinite operators with absolutely summable entries*, arXiv:2409.20372 [math.FA] [math.CA]

Motivation

A Problem by Feichtinger

At a 2004 Oberwolfach meeting, Hans Feichtinger asked the following question:

(Q) Given a positive semi-definite (psd) trace-class operator

$T : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$, $Tf(x) = \int K(x, y)f(y)dy$, with $K \in M^1(\mathbb{R}^d \times \mathbb{R}^d)$, is

there a factorization $T = \sum_k \langle \cdot, g_k \rangle g_k$ such that $\sum_k \|g_k\|_{M^1}^2 < \infty$?

C. Heil and D. Larson reformulated the problem equivalently as:

(Q') Given an infinite matrix $A = (A_{m,n})_{m,n \geq 1}$ so that: (1) $A = A^* \geq 0$, and (2)

$\|A\|_1 := \sum_{m,n \geq 1} |A_{m,n}| < \infty$ is there a factorization $A = \sum_{k \geq 1} f_k f_k^*$ so that

$\sum_{k \geq 1} \|f_k\|_1^2 < \infty$?

Motivation

A Problem by Feichtinger

At a 2004 Oberwolfach meeting, Hans Feichtinger asked the following question:

(Q) Given a positive semi-definite (psd) trace-class operator

$T : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$, $Tf(x) = \int K(x, y)f(y)dy$, with $K \in M^1(\mathbb{R}^d \times \mathbb{R}^d)$, is

there a factorization $T = \sum_k \langle \cdot, g_k \rangle g_k$ such that $\sum_k \|g_k\|_{M^1}^2 < \infty$?

C. Heil and D. Larson reformulated the problem equivalently as:

(Q') Given an infinite matrix $A = (A_{m,n})_{m,n \geq 1}$ so that: (1) $A = A^* \geq 0$, and (2)

$\|A\|_1 := \sum_{m,n \geq 1} |A_{m,n}| < \infty$ is there a factorization $A = \sum_{k \geq 1} f_k f_k^*$ so that

$\sum_{k \geq 1} \|f_k\|_1^2 < \infty$?

Based on [BandeiraMixonSteinerberger'24] and [B.Jiang'24], the answer is

NEGATIVE in general!

Today we focus on characterizations of operators that admit such factorizations.

Easy decomposition

(Q') Given an infinite matrix $A = (A_{m,n})_{m,n \geq 1}$ so that: (1) $A = A^* \geq 0$, and (2) $\|A\|_1 := \sum_{m,n \geq 1} |A_{m,n}| < \infty$ is there a factorization $A = \sum_{k \geq 1} f_k f_k^*$ so that $\sum_{k \geq 1} \|f_k\|_1^2 < \infty$?

In general, the answer to Q' is **NEGATIVE**. ...

However:

Given an infinite matrix $A = (A_{m,n})_{m,n \geq 1}$ so that: (1) $A = A^*$, and (2) $\|A\|_1 := \sum_{m,n \geq 1} |A_{m,n}| < \infty$, it is easy to see that there is a decomposition $A = \sum_{k \geq 1} \varepsilon_k g_k g_k^*$ with $\varepsilon_k \in \{-1, +1\}$ and $\sum_{k \geq 1} \|g_k\|_1^2 \leq 2 \sum_{m,n \geq 1} |A_{m,n}| < \infty$. Furthermore, $\sum_{m,n \geq 1} |A_{m,n}| \leq \sum_{k \geq 1} \|g_k\|_1^2$ is always true.

Thus, the question is whether one can choose g_k 's so that $\varepsilon_k = 1$ for every k .

Classes of Operators

Notations:

$$\mathbb{V} := \left\{ A = (A_{m,n})_{m,n \geq 1} : A = A^*, \|A\|_1 = \sum_{m,n=1}^{\infty} |A_{m,n}| < \infty \right\}$$

$$\Gamma := \{A \in \mathbb{V} : A \succcurlyeq 0 \text{ as a bilinear form on } \ell_2\}$$

$$\Gamma_+ := \left\{ A \in \Gamma : \exists (x_k)_k, A = \sum_k x_k x_k^*, \sum_k \|x_k\|_1^2 < \infty \right\}.$$

For $A \in \Gamma_+$, $\gamma_+(A) := \inf \left\{ \sum_{k \geq 1} \|x_k\|_1^2 : A = \sum_{k \geq 1} x_k x_k^* \text{ strongly} \right\}$.

Theorem (Corollary of the BMS Result)

$$\Gamma_+ \subsetneq \Gamma.$$

Our problem: Understand $\Gamma \setminus \Gamma_+$ and find conditions for an operator $A \in \Gamma_+$.

Topological properties of Γ_+ : $\|\cdot\|_1$ -topology

Theorem (Density)

Let $A \in \Gamma$ and $\varepsilon > 0$. There exist operators $B, C \in \Gamma_+$ so that:

- ① $A = B - C$;
- ② $\gamma_+(C) < \varepsilon$.

In particular, Γ_+ is dense in Γ in the $\|\cdot\|_1$ topology.

As remarked earlier, $A = \sum_k \varepsilon_k g_k g_k^*$ for some $g_k \in \ell_1$ and $\varepsilon_k \in \{-1, +1\}$, with $\sum_k \|g_k\|_1^2 < \infty$.

The result above says that $g_k = g_k^\varepsilon$'s can be chosen so that for arbitrary small ε ,

$$\sum_{k:\varepsilon_k=-1} \|g_k^\varepsilon\|_1^2 < \varepsilon.$$

However, for $A \in \Gamma \setminus \Gamma_+$,

$$\lim_{\varepsilon \rightarrow 0} \sum_k \|g_k^\varepsilon\|_1^2 = \infty.$$

Algebraic properties of Γ_+

Γ_+ is a convex cone. Furthermore, it enjoys the following two properties:

Theorem (Conjugacy invariance)

The class Γ_+ is invariant to conjugacy with $B(\ell_1, \ell_1)$ operators. Specifically, if $A \in \Gamma_+$ and $T : \ell_1 \rightarrow \ell_1$ bounded operator, then $TAT^ \in \Gamma_+$.*

Theorem (Algebraic Cone)

Let p be a univariate polynomial with nonnegative coefficients. Suppose $A \in \Gamma_+$. Then $p(A) \in \Gamma_+$.

Furthermore, let $f(z) = \sum_{k=1}^{\infty} c_k z^k$ with $c_k \geq 0$ for all k and $\sum_{k=1}^{\infty} c_k < \infty$. Let $g(z) = \frac{f(z)}{z} = \sum_{k=0}^{\infty} c_{k+1} z^k$. Then for every $A \in \Gamma_+$, we have $g(A) \in \Gamma_+$ with

$$\gamma_+(f(A)) \leq g(\|A\|_1) \cdot \gamma_+(A).$$

Sufficiency criterion for Γ_+

Theorem

Suppose $A \in \mathcal{B}(\ell_2, \ell_2)$ is positive and self-adjoint, and let $D := \sum_k \sqrt{A_{k,k}} \in [0, \infty]$.

- Ⓐ If $D < \infty$, then $A^{1/2} \in \Pi_2(\ell_2, \ell_1)$ (i.e., $A^{1/2}$ is 2-summing) with $\pi_2(A^{1/2}) \leq D$. Moreover, $A \in \Gamma_+$ with $\gamma_+(A) \leq D^2$.
- Ⓑ If $A^{1/2}$ is 2-summing, i.e. $A^{1/2} \in \Pi_2(\ell_2, \ell_1)$, then $A \in \Gamma_+$ and $D \leq \kappa_G \cdot \pi_2(A^{1/2})$, with κ_G being the Grothendieck constant.

Recall: an operator $T : \ell_2 \rightarrow \ell_1$ is called 2-summing if there is $C > 0$ such that for any finite collection $x_1, \dots, x_N \in \ell_2$, we have

$$\sum_{k=1}^N \|Tx_k\|_1^2 \leq C^2 \cdot \sup_{z \in \ell_2: \|z\|_2=1} \sum_{k=1}^N |\langle z, x_k \rangle|^2 = C^2 \left\| \sum_{k=1}^N x_k x_k^* \right\|_{2 \rightarrow 2}.$$

The infimum of all such C is denoted by $\pi_2(T)$, called the 2-summing norm of T . The set of such 2-summing operators is denoted by $\Pi_2(\ell_2, \ell_1)$.

Other factorizations

The previous factorization is the special case of the following result:

Theorem

If $A \in \Gamma$ admits a factorization $A = TT^*$ such that $T \in \Pi_2(\ell_2, \ell_1)$, then $A \in \Gamma_+$. Moreover, for any eigendecomposition $A = \sum_k \lambda_k e_k e_k^*$, we have

$$\gamma_+(A) \leq \sum_k \|A^{1/2} e_k\|_1^2 = \sum_k \lambda_k \|e_k\|_1^2 \leq \pi_2(T)^2 < \infty.$$

Remark: $A^{1/2} \in \Pi_2(\ell_2, \ell_1)$ if and only if $\sum_k \sqrt{A_{k,k}} < \infty$.

New Γ_+ -induced topology

Let τ be the topology on \mathbb{V} generated by open sets of the form $T + \mathcal{U}_\varepsilon$ for $\varepsilon > 0$, where $T \in \mathbb{V}$ and $\mathcal{U}_\varepsilon := \{A \in \Gamma_+ : \gamma_+(A) < \varepsilon\}$. Intuitively, τ measures “approximation from above”. Furthermore,

$$\mathcal{U}_\varepsilon \subsetneq \{A \in \Gamma_+ : \|A\|_{1,1} < \varepsilon\} \subsetneq \{A \in \mathbb{V} : A \geq 0, \|A\|_{1,1} < \varepsilon\} \subsetneq \{A \in \mathbb{V} : \|A\|_{1,1} < \varepsilon\}$$

Proposition

The following are true about τ .

- Ⓐ τ is Hausdorff, translation invariant, and first-countable.
- Ⓑ The family $\mathcal{F} = \{T + \mathcal{U}_\varepsilon : T \in \mathbb{V}, \varepsilon > 0\}$ forms a basis for τ , i.e., every open set in τ can be written as a union of elements in \mathcal{F} .
- Ⓒ If $A \in \mathcal{O} \in \tau$ then there exists $r > 0$ so that $A + \mathcal{U}_r \subset \mathcal{O}$. In particular, τ is not the discrete topology on \mathbb{V} .
- Ⓓ Let $A_0 \in \Gamma \setminus \Gamma_+$. Then the subspace $\{tA_0; t \in \mathbb{R}\}$ with the induced topology is discrete.
- Ⓔ τ is not second-countable.

Topological properties of Γ_+ : τ -topology

Theorem

Let \mathbb{V} , Γ , Γ_+ , and the topology τ generated by $T + \mathcal{U}_\varepsilon$ with $T \in \mathbb{V}$ and $\varepsilon > 0$. Then

- Ⓐ Γ_+ is dense in Γ with respect to τ ;
- Ⓑ Γ is the τ -closure of Γ_+ ;
- Ⓒ Γ_+ is reproducing, namely, $\mathbb{V} = \Gamma_+ - \Gamma_+$.

Remark: This result provides a more precise approximation and density than the $\|\cdot\|_1$ induced topology.

Thank you for listening! ... QUESTIONS?

Warm-Up Exercise

Let $A \in \mathbb{R}^{n \times n}$ be a matrix and consider the following optimization problem:

$$\gamma(A) := \inf_{A = \sum_{k \geq 1} x_k y_k^T} \sum_k \|x_k\|_1 \|y_k\|_1$$

Warm-Up Exercise

Let $A \in \mathbb{R}^{n \times n}$ be a matrix and consider the following optimization problem:

$$\gamma(A) := \inf_{A = \sum_{k \geq 1} x_k y_k^T} \sum_k \|x_k\|_1 \|y_k\|_1$$

Note:

$$A = [A_{i,j}]_{i,j \in [n]} = \begin{bmatrix} A_{11} \\ A_{21} \\ \vdots \\ A_{n1} \end{bmatrix} \cdot [1, 0, \dots, 0] + \dots + \begin{bmatrix} A_{1n} \\ A_{2n} \\ \vdots \\ A_{nn} \end{bmatrix} \cdot [0, 0, \dots, 1]$$

From where: $\gamma(A) \leq \sum_{i,j} |A_{i,j}| =: \|A\|_1$.

Warm-Up Exercise

Let $A \in \mathbb{R}^{n \times n}$ be a matrix and consider the following optimization problem:

$$\gamma(A) := \inf_{A = \sum_{k \geq 1} x_k y_k^T} \sum_k \|x_k\|_1 \|y_k\|_1$$

Note:

$$A = [A_{i,j}]_{i,j \in [n]} = \begin{bmatrix} A_{11} \\ A_{21} \\ \vdots \\ A_{n1} \end{bmatrix} \cdot [1, 0, \dots, 0] + \dots + \begin{bmatrix} A_{1n} \\ A_{2n} \\ \vdots \\ A_{nn} \end{bmatrix} \cdot [0, 0, \dots, 1]$$

From where: $\gamma(A) \leq \sum_{i,j} |A_{i,j}| =: \|A\|_1$.

For converse: Let $A = \sum_k x_k y_k^T$ be the optimal decomposition. Then:

$$\|A\|_1 = \left\| \sum_k x_k y_k^T \right\|_1 \leq \sum_k \|x_k y_k^T\|_1 = \sum_k \|x_k\|_1 \|y_k\|_1 = \gamma(A).$$

Projective norm

We obtained:

$$\gamma(A) := \inf_{A = \sum_{k \geq 1} x_k y_k^T} \sum_k \|x_k\|_1 \|y_k\|_1 = \sum_{i,j} |A_{i,j}| =: \|A\|_1$$

following Grothendieck, the last norm is sometime referred to as *projective norm*, $\|A\|_{\wedge}$.

Projective norm

We obtained:

$$\gamma(A) := \inf_{A = \sum_{k \geq 1} x_k y_k^T} \sum_k \|x_k\|_1 \|y_k\|_1 = \sum_{i,j} |A_{i,j}| =: \|A\|_1$$

following Grothendieck, the last norm is sometime referred to as *projective norm*, $\|A\|_{\wedge}$.

Assume now that $A = A^T$. Considered a more constrained optimization problem:

$$\gamma_0(A) := \inf_{\substack{A = \sum_{k \geq 1} \varepsilon_k x_k x_k^T \\ \varepsilon_k \in \{+1, -1\}}} \sum_k \|x_k\|_1^2$$

Projective norm

We obtained:

$$\gamma(A) := \inf_{A = \sum_{k \geq 1} x_k y_k^T} \sum_k \|x_k\|_1 \|y_k\|_1 = \sum_{i,j} |A_{i,j}| =: \|A\|_1$$

following Grothendieck, the last norm is sometime referred to as *projective norm*, $\|A\|_{\wedge}$.

Assume now that $A = A^T$. Considered a more constrained optimization problem:

$$\gamma_0(A) := \inf_{\substack{A = \sum_{k \geq 1} \varepsilon_k x_k x_k^T \\ \varepsilon_k \in \{+1, -1\}}} \sum_k \|x_k\|_1^2$$

It is not hard to show that γ_0 is a norm on $Sym(\mathbb{R}^n)$ (or $Sym(\mathbb{C}^n)$), and $\|A\|_1 \leq \gamma_0(A)$.

Projective norm

We obtained:

$$\gamma(A) := \inf_{A = \sum_{k \geq 1} x_k y_k^T} \sum_k \|x_k\|_1 \|y_k\|_1 = \sum_{i,j} |A_{i,j}| =: \|A\|_1$$

following Grothendieck, the last norm is sometime referred to as *projective norm*, $\|A\|_{\wedge}$.

Assume now that $A = A^T$. Considered a more constrained optimization problem:

$$\gamma_0(A) := \inf_{\substack{A = \sum_{k \geq 1} \varepsilon_k x_k x_k^T \\ \varepsilon_k \in \{+1, -1\}}} \sum_k \|x_k\|_1^2$$

It is not hard to show that γ_0 is a norm on $Sym(\mathbb{R}^n)$ (or $Sym(\mathbb{C}^n)$), and

$\|A\|_1 \leq \gamma_0(A)$. Leveraging the fact that

$\frac{1}{2}(xy^T + yx^T) = \frac{1}{4}((x+y)(x+y)^T - (x-y)(x-y)^T)$ one obtains:

$$\|A\|_1 \leq \gamma_0(A) := \inf_{\substack{A = \sum_{k \geq 1} \varepsilon_k x_k x_k^T \\ \varepsilon_k \in \{+1, -1\}}} \sum_k \|x_k\|_1^2 \leq 2\|A\|_1$$

Problem Formulation

Let $\text{Sym}^+(\mathbb{C}^n) = \{A \in \mathbb{C}^{n \times n}, A^* = A \geq 0\}$. For $A \in \text{Sym}^+(\mathbb{C}^n)$, denote

$$\gamma_+(A) := \inf_{A = \sum_{k \geq 1} x_k x_k^*} \sum_k \|x_k\|_1^2$$

It is obvious that $\|A\|_1 \leq \gamma_0(A) \leq \gamma_+(A)$.

Problem Formulation

Let $\text{Sym}^+(\mathbb{C}^n) = \{A \in \mathbb{C}^{n \times n}, A^* = A \geq 0\}$. For $A \in \text{Sym}^+(\mathbb{C}^n)$, denote

$$\gamma_+(A) := \inf_{A = \sum_{k \geq 1} x_k x_k^*} \sum_k \|x_k\|_1^2$$

It is obvious that $\|A\|_1 \leq \gamma_0(A) \leq \gamma_+(A)$.

The *matrix problem*: For every $n \geq 1$ find the best constant C_n such that, for every $A \in \text{Sym}^+(\mathbb{C}^n)$,

$$\gamma_+(A) \leq C_n \|A\|_1 := C_n \sum_{k,l=1}^n |A_{k,l}|$$

That is, we are interested in finding:

$$C_n = \sup_{A \geq 0} \frac{\gamma_+(A)}{\|A\|_1}$$

Properties of $\gamma_+(A)$

The infimum is achieved:

$$\gamma_+(A) := \inf_{A = \sum_{k \geq 1} x_k x_k^*} \sum_k \|x_k\|_1^2 = \min_{A = \sum_{k=1}^{n^2} x_k x_k^*} \sum_k \|x_k\|_1^2.$$

Properties of $\gamma_+(A)$

The infimum is achieved:

$$\gamma_+(A) := \inf_{A = \sum_{k \geq 1} x_k x_k^*} \sum_k \|x_k\|_1^2 = \min_{A = \sum_{k=1}^{n^2} x_k x_k^*} \sum_k \|x_k\|_1^2.$$

Upper bounds:

$$\gamma_+(A) \leq n \operatorname{trace}(A) \leq n \|A\|_1 = n \sum_{k,j} |A_{k,j}|$$

$$\gamma_+(A) \leq n \operatorname{trace}(A) \leq n^2 \|A\|_{Op}$$

Properties of $\gamma_+(A)$

The infimum is achieved:

$$\gamma_+(A) := \inf_{A = \sum_{k \geq 1} x_k x_k^*} \sum_k \|x_k\|_1^2 = \min_{A = \sum_{k=1}^{n^2} x_k x_k^*} \sum_k \|x_k\|_1^2.$$

Upper bounds:

$$\gamma_+(A) \leq n \operatorname{trace}(A) \leq n \|A\|_1 = n \sum_{k,j} |A_{k,j}|$$

$$\gamma_+(A) \leq n \operatorname{trace}(A) \leq n^2 \|A\|_{op}$$

Lower bounds:

$$\|A\|_1 = \min_{A = \sum_{k \geq 1} x_k y_k^*} \sum_k \|x_k\|_1 \|y_k\|_1 \leq \gamma_+(A)$$

Properties of $\gamma_+(A)$

The infimum is achieved:

$$\gamma_+(A) := \inf_{A=\sum_{k \geq 1} x_k x_k^*} \sum_k \|x_k\|_1^2 = \min_{A=\sum_{k=1}^{n^2} x_k x_k^*} \sum_k \|x_k\|_1^2.$$

Upper bounds:

$$\gamma_+(A) \leq n \operatorname{trace}(A) \leq n \|A\|_1 = n \sum_{k,j} |A_{k,j}|$$


$$\gamma_+(A) \leq n \operatorname{trace}(A) \leq n^2 \|A\|_{op}$$

Lower bounds:

$$\|A\|_1 = \min_{A=\sum_{k \geq 1} x_k y_k^*} \sum_k \|x_k\|_1 \|y_k\|_1 \leq \gamma_+(A)$$

Convexity: for $A, B \in \operatorname{Sym}^+(\mathbb{C}^n)$ and $t \geq 0$,

$$\gamma_+(A+B) \leq \gamma_+(A) + \gamma_+(B) \quad , \quad \gamma_+(tA) = t\gamma_+(A)$$



Maximum of $\sum_k \|x_k\|_1^2 / \|A\|_1$ over 30 random noise realizations, where x_k 's are obtained from the eigendecomposition, or the LDL factorization.

Properties of $\gamma_+(A)$

Lower bound is achieved, $\gamma_+(A) = \|A\|_1$ in the following cases:

- 1 If $A = xx^*$ is of rank one.
- 2 If $A \geq 0$ is a diagonally dominant matrix, $A_{ii} \geq \sum_{k \neq i} |A_{i,k}|$.
- 3 If $A \geq 0$ admits a Non Negative Matrix Factorization (NNMF), $A = BB^T$ with $B_{ij} \geq 0$.

Continuity, Lipschitz and linear program reformulation:

- 1 $\gamma_+ : \text{Sym}^+(\mathbb{C}^n) \rightarrow \mathbb{R}$ is continuous.
- 2 If $A, B \geq \delta I$ and $\text{trace}(A), \text{trace}(B) \leq 1$ then

$$|\gamma_+(A) - \gamma_+(B)| \leq \left(\frac{n}{\delta^2} + n^2 \right) \|A - B\|_{op}.$$

- 3 Let $S_1 = \{x \in \mathbb{C}^n, \|x\|_1 = 1\}$ denote the compact unit sphere with respect to the l^1 norm, and let $\mathcal{B}(S_1)$ denote the set of Borel measures over S_1 . Then:

$$\gamma_+(A) = \inf_{\mu \in \mathcal{B}(S_1): \int_{S_1} xx^* d\mu(x) = A} \mu(S_1), \quad \mu^*(x) = \sum_{k=1}^m \lambda_k \delta(x - g_k)$$

where $\gamma_+(A) = \sum_{k=1}^m \lambda_k$ and $A = \sum_{k=1}^m \lambda_k g_k g_k^*$ is the optimal factorization.

Primal and dual problems for γ_+

The linear program is a convex optimization problem (which is great), but it is defined on an infinite dimensional space (not so great!).

Its dual problem enjoys strong duality (this may not be obvious due to infinite dimensional technical issues):

Theorem

Assume $A \geq 0$. Its associated primal (min) & dual (max) problems are:

$$\max_{T=T^*: \langle T, x \rangle \leq 1, \forall \|x\|_1 \leq 1} \text{trace}(TA) = \min_{\mu \in \mathcal{B}(S_1): \int_{S_1} xx^* d\mu(x) = A} \mu(S_1) = \gamma_+(A)$$

Primal and dual problems for γ_+

The linear program is a convex optimization problem (which is great), but it is defined on an infinite dimensional space (not so great!).

Its dual problem enjoys strong duality (this may not be obvious due to infinite dimensional technical issues):

Theorem

Assume $A \succeq 0$. Its associated primal (min) & dual (max) problems are:

$$\max_{T=T^*: \langle T, x \rangle \leq 1, \forall \|x\|_1 \leq 1} \text{trace}(TA) = \min_{\mu \in \mathcal{B}(S_1): \int_{S_1} xx^* d\mu(x) = A} \mu(S_1) = \gamma_+(A)$$

Why? Assume the optimal support $\{x_1, \dots, x_m\}$ is known.

Construct the Lagrange function

$$L(t_1, \dots, t_m; T, \nu_1, \dots, \nu_m) = \sum_{k=1}^m t_k + \text{trace} \left(T \left(A - \sum_{k=1}^m t_k x_k x_k^* \right) \right) - \sum_{k=1}^m \nu_k t_k.$$

The dual function

$$g(T, \nu_1, \dots, \nu_m) := \inf_t L(\mathbf{t}; T, \nu) = \begin{cases} \text{trace}(TA) & \text{if } 1 - \langle T x_k, x_k \rangle - \nu_k = 0, \nu_k \geq 0 \\ -\infty & \text{if otherwise} \end{cases}$$

Note the quantity:

$$\rho_1(T) = \max_{x: \|x\|_1 \leq 1} \langle Tx, x \rangle$$

The dual problem

$$\max_{T=T^*: \langle Tx, x \rangle \leq 1, \forall \|x\|_1 \leq 1} \text{trace}(TA)$$

can be reformulated as

$$\gamma_+(A) = \max_{T=T^*: \rho_1(T) \leq 1} \text{trace}(TA)$$

The optimal constant C_n from $\gamma_+(A) \leq C_n \|A\|_1$ turns into

$$C_n = \max_{A \geq 0: \|A\|_1 \leq 1} \gamma_+(A) = \max_{\substack{A \geq 0: \\ \|A\|_1 \leq 1}} \max_{\substack{T=T^*: \\ \rho_1(T) \leq 1}} \text{trace}(TA) = \max_{A \geq 0: A \neq 0} \max_{\substack{T=T^*: \\ \rho_1(T) > 0}} \frac{\text{trace}(TA)}{\|A\|_1 \rho_1(T)}$$

The bound ρ_1

Recall, for $T = T^*$:

$$\rho_1(T) = \max_{x: \|x\|_1 \leq 1} \langle Tx, x \rangle$$

How to compute it?

Easy cases:

- ① If $T \leq 0$ then $\rho_1(T) = 0$
- ② If $T \geq 0$ then

$$\rho_1(T) = \max_k T_{k,k} = \max_{i,j} |T_{i,j}| =: \|T\|_\infty$$

This resembles the *numerical radius* of a matrix, $r(T) = \max_{\|x\|_2=1} |\langle Tx, x \rangle|$, which for hermitian matrices equals the largest singular value (operator norm). Note differences: (i) $\|\cdot\|_2 \rightarrow \|\cdot\|_1$; (ii) no absolute value $|\cdot|$.

The bound ρ_1 (2)

Assume $\lambda_{\max}(T) > 0$, i.e. T is NOT negative semi-definite. Then:

$$\rho_1(T) = \max_{x: \|x\|_1=1} \langle Tx, x \rangle = \max_{\substack{A \geq 0 : \\ \text{rank}(A) = 1 \\ \|A\|_1 = 1}} \text{trace}(TA) = \max_{\substack{A \geq 0 : \\ \text{rank}(A) = 1 \\ \|A\|_1 \leq 1}} \text{trace}(TA)$$

The bound ρ_1 (2)

Assume $\lambda_{\max}(T) > 0$, i.e. T is NOT negative semi-definite. Then:

$$\rho_1(T) = \max_{x: \|x\|_1=1} \langle Tx, x \rangle = \max_{\substack{A \geq 0 : \\ \text{rank}(A) = 1 \\ \|A\|_1 = 1}} \text{trace}(TA) = \max_{\substack{A \geq 0 : \\ \text{rank}(A) = 1 \\ \|A\|_1 \leq 1}} \text{trace}(TA)$$

Convex relaxation:

$$\pi_+(T) := \max_{\substack{A \geq 0 : \\ \|A\|_1 \leq 1}} \text{trace}(TA)$$

which is a semi-definite program (SDP). Thus:

$$\rho_1(T) \leq \pi_+(T).$$

Primal and dual problems for ρ_1

The SDP enjoys strong duality:

Theorem

Assume $T = T^*$. The primal-dual programs have strong duality:

$$\pi_+(T) = \max_{\substack{A \geq 0 : \\ \|A\|_1 \leq 1}} \text{trace}(TA) = \min_{Y \geq 0} \|T + Y\|_\infty$$

where $\|Z\|_\infty = \max_{i,j} |Z_{i,j}|$.

The proof of this theorem is based on the Von Neumann's min-max theorem:

$$\begin{aligned} \min_{Y \geq 0} \|T + Y\|_\infty &= \min_{Y \geq 0} \max_{A: \|A\|_1 \leq 1} \text{trace}((T + Y)A) \stackrel{vN}{=} \max_{A: \|A\|_1 \leq 1} \min_{Y \geq 0} \text{trace}((T + Y)A) = \\ &= \max_{A: \|A\|_1 \leq 1} \left(\text{trace}(TA) + \min_{Y \geq 0} \text{trace}(YA) \right) = \max_{A \geq 0: \|A\|_1 \leq 1} \left(\text{trace}(TA) + \min_{Y \geq 0} \text{trace}(YA) \right) = \\ &= \max_{A \geq 0: \|A\|_1 \leq 1} \text{trace}(TA) = \pi_+(T) \end{aligned}$$

Closing the loop

The final result: the connection between $\gamma_+(A)$ and C_n on one hand, and $\rho_1(T)$ and $\pi_+(T)$ on the other hand:

Theorem

$$C_n := \max_{\substack{A \geq 0 \\ A \neq 0}} \frac{\gamma_+(A)}{\|A\|_1} = \max_{\substack{T = T^* \\ \rho_1(T) \neq 0}} \frac{\pi_+(T)}{\rho_1(T)}$$

Closing the loop

The final result: the connection between $\gamma_+(A)$ and C_n on one hand, and $\rho_1(T)$ and $\pi_+(T)$ on the other hand:

Theorem

$$C_n := \max_{\substack{A \geq 0 \\ A \neq 0}} \frac{\gamma_+(A)}{\|A\|_1} = \max_{\substack{T = T^* \\ \rho_1(T) \neq 0}} \frac{\pi_+(T)}{\rho_1(T)}$$

The proof is based on the earlier derivation:

$$\begin{aligned} C_n &= \max_{\substack{A \geq 0 \\ A \neq 0}} \max_{\substack{T = T^* \\ \rho_1(T) > 0}} \frac{\text{trace}(TA)}{\|A\|_1 \rho_1(T)} = \max_{\substack{T = T^* \\ \rho_1(T) > 0}} \max_{\substack{A \geq 0 \\ A \neq 0}} \frac{\text{trace}(TA)}{\|A\|_1 \rho_1(T)} = \\ &= \max_{\substack{T = T^* \\ \rho_1(T) > 0}} \frac{1}{\rho_1(T)} \max_{\substack{A \geq 0 \\ \|A\|_1 = 1}} \text{trace}(TA) = \max_{\substack{T = T^* \\ \rho_1(T) > 0}} \frac{\pi_+(T)}{\rho_1(T)} \end{aligned}$$

Bandeira-Mixon-Steinerberger Result

Last year B-M-S announced:

Theorem (Afonso Bandeira, Dustin Mixon, Stefan Steinerberger - Oberwolfach 2024; ACHA 2024)

There are $\alpha > 0$, $N_0 > 1$ so that for any $n \geq N_0$,

$$C_n \geq \alpha\sqrt{n}$$

Consequence:

Theorem (R.B, F.Jiang)

1. There exists a PSD trace-class $A = A^* \geq 0$, $A = (A_{m,n})_{m,n \geq 0}$ with $\sum_{m,n} |A_{m,n}| < \infty$, so that for any factorization $A = \sum_{k \geq 0} f_k f_k^*$, $\sum_{k \geq 0} \|f_k\|_1^2 = \infty$.
2. There exists $K \in M^1(\mathbb{R}^2)$ so that: (i) $K(x,y) = \overline{K(y,x)}$ for all x,y ; (ii) $\int_{\mathbb{R}^2} K(x,y) f(y) \overline{f(x)} dx dy \geq 0$ for all $f \in L^2(\mathbb{R})$; and (iii) for any $(g_n)_{n \geq 0}$ with $\int_{\mathbb{R}} K(x,y) f(y) dy = \sum_{n \geq 0} \langle f, g_n \rangle g_n(x)$, $\forall f \in L^2(\mathbb{R})$, $\sum_{n \geq 0} \|g_n\|_{M^1}^2 = \infty$.

The Continuity Property

Theorem (The Continuity Property)

The map $\gamma_+ : (\text{Sym}^+(\mathbb{C}^n), \|\cdot\|) \rightarrow \mathbb{R}$ is continuous.

Remarks

- 1 This statement extends the continuity result from $\text{Sym}^{++}(\mathbb{C}^n) = \{A = A^* > 0\}$ to $\text{Sym}^+(\mathbb{C}^n) = \{A = A^* \geq 0\}$.
- 2 Proof is based on a (new?) comparison result between non-negative operators.
- 3 Global Lipschitz is still open.

The Continuity Property

The proof is based on the following two lemmas:

Lemma (L1)

Let $A \in \text{Sym}^+(\mathbb{C}^n)$ of rank $r > 0$. Let $\lambda_r > 0$ denote the r^{th} eigenvalue of A , and let $P_{A,r}$ denote the orthogonal projection onto the range of A . For any $0 < \varepsilon < 1$ and $B \in \text{Sym}^+(\mathbb{C}^n)$ such that $\|A - B\|_{op} \leq \frac{\varepsilon \lambda_r}{1 - \varepsilon}$, the following holds true:

$$A - (1 - \varepsilon)P_{A,r}BP_{A,r} \geq 0 \quad (1)$$

Lemma (L2)

Let $A \in \text{Sym}^+(\mathbb{C}^n)$ of rank $r > 0$. Let $\lambda_r > 0$ denote the r^{th} eigenvalue of A . For any $0 < \varepsilon < \frac{1}{2}$ and $B \in \text{Sym}^+(\mathbb{C}^n)$ such that $\|A - B\|_{op} \leq \varepsilon \lambda_r$, the following holds true:

$$B - (1 - \varepsilon)P_{B,r}AP_{B,r} \geq 0 \quad (2)$$

where $P_{B,r}$ denotes the orthogonal projection onto the top r eigenspace of B .

Proof of Continuity of γ_+

Fix $A \in \text{Sym}^+(\mathbb{C}^n)$. Let $(B_j)_{j \geq 1}$, $B_j \in \text{Sym}^+(\mathbb{C}^n)$, be a convergent sequence to A . We need to show $\gamma_+(B_j) \rightarrow \gamma_+(A)$.

Let $A = \sum_{k=1}^{n^2} x_k x_k^*$ be the optimal decomposition of A such that

$$\gamma_+(A) = \sum_{k=1}^{n^2} \|x_k\|_1^2.$$

If $A = 0$ then $\gamma_+(A) = 0$ and

$$0 \leq \gamma_+(B_j) \leq n \text{trace}(B_j) \leq n^2 \|B_j\|_{op}.$$

Hence $\lim_j \gamma_+(B_j) = 0$.

Assume $\text{rank}(A) = r > 0$ and let $\lambda_r > 0$ denote the smallest strictly positive eigenvalue of A . Let $\varepsilon \in (0, \frac{1}{2})$ be arbitrary. Let $J = J(\varepsilon)$ be so that

$\|A - B_j\|_{op} < \varepsilon \lambda_r$ for all $j > J$. Let $B_j = \sum_{k=1}^{n^2} y_{j,k} y_{j,k}^*$ be the optimal decomposition of B_j such that $\gamma_+(B_j) = \sum_{k=1}^{n^2} \|y_{j,k}\|_1^2$.

Let $\Delta_j = A - (1 - \varepsilon) P_{A,r} B_j P_{A,r}$. By Lemma L1, for any $j > J$,

$$\gamma_+(A) \leq (1 - \varepsilon) \gamma_+(P_{A,r} B_j P_{A,r}) + \gamma_+(\Delta_j) \leq (1 - \varepsilon) \sum_{k=1}^{n^2} \|P_{A,r} y_{j,k}\|_1^2 + n \text{trace}(\Delta_j)$$

Proof of Continuity of γ_+ (cont)

Pass to a subsequence j' of j so that $y_{j',k} \rightarrow y_k$, for every $k \in [n^2]$, and $\gamma_+(B_{j'}) \rightarrow \liminf_j \gamma_+(B_j)$. Then $\lim_{j'} P_{A,r} y_{j',k} = P_{A,r} y_k = y_k$ and

$$\lim_{j'} \sum_{k=1}^{n^2} \|P_{A,r} y_{j',k}\|_1^2 = \lim_{j'} \sum_{k=1}^{n^2} \|y_{j',k}\|_1^2 = \lim_j \inf \gamma_+(B_j)$$

On the other hand, $\lim_j \text{trace}(\Delta_j) = \varepsilon \text{trace}(A)$. Hence:

$$\gamma_+(A) \leq (1 - \varepsilon) \lim_j \inf \gamma_+(B_j) + \varepsilon \text{trace}(A)$$

Since $\varepsilon > 0$ is arbitrary, it follows $\gamma_+(A) \leq \liminf_j \gamma_+(B_j)$.

The inequality $\limsup_j \gamma_+(B_j) \leq \gamma_+(A)$ follows from Lemma L2 similarly: with

$\Delta_j = B_j - (1 - \varepsilon)P_{B_j,r}AP_{B_j,r}$ and $A = \sum_{k=1}^{n^2} x_k x_k^*$ optimal,

$$\gamma_+(B_j) \leq (1 - \varepsilon)\gamma_+(P_{B_j,r}AP_{B_j,r}) + n \text{trace}(\Delta_j) = (1 - \varepsilon) \sum_{k=1}^{n^2} \|P_{B_j,r} x_k\|_1^2 + n \text{trace}(\Delta_j).$$

Next take limsup of lhs by noticing $P_{B_j,r} \rightarrow P_{A,r}$ and $\limsup_j \|\Delta_j\|_{Op} = \varepsilon \|A\|_{Op}$:

$\limsup_j \gamma_+(B_j) \leq (1 - \varepsilon)\gamma_+(A) + n^2 \varepsilon \|A\|_{Op}$. Take $\varepsilon \rightarrow 0$ and result follows. 

Proof of Lemmas

Proof of Lemma L1

Let $P = P_{A,r}$. and $\Delta = A - (1 - \varepsilon)P_{A,r}BP_{A,r}$. For any $x \in \mathbb{C}^n$:

$$\begin{aligned} \langle \Delta x, x \rangle &= \langle APx, Px \rangle - (1 - \varepsilon)\langle BPx, Px \rangle = \langle (A - (1 - \varepsilon)B)Px, Px \rangle = \\ &= \varepsilon\langle APx, Px \rangle + (1 - \varepsilon)\langle (A - B)Px, Px \rangle \geq \varepsilon\lambda_r\|Px\|^2 - (1 - \varepsilon)\|A - B\|_{Op}\|Px\|^2 \geq 0 \end{aligned}$$

because $\|A - B\|_{Op} \leq \frac{\varepsilon\lambda_r}{1 - \varepsilon}$.

Proof of Lemma L2

Let $P = P_{B,r}$ and $\Delta = B - (1 - \varepsilon)P_{B,r}AP_{B,r}$. Let $C = B - P_{B,r}BP_{B,r} \geq 0$. Let μ_r be the r^{th} eigenvalue of B . Note $|\mu_r - \lambda_r| \leq \|A - B\|_{Op} \leq \varepsilon\lambda_r$. Thus $\mu_r \geq (1 - \varepsilon)\lambda_r$. For any $x \in \mathbb{C}^n$:

$$\begin{aligned} \langle \Delta x, x \rangle &= \langle Cx, x \rangle + \langle BPx, Px \rangle - (1 - \varepsilon)\langle APx, Px \rangle = \langle Cx, x \rangle + \varepsilon\langle BPx, Px \rangle + \\ &+ (1 - \varepsilon)\langle (B - A)Px, Px \rangle \geq \langle Cx, x \rangle + (\varepsilon\mu_r - (1 - \varepsilon)\|A - B\|_{Op})\|Px\|^2 \geq 0 \end{aligned}$$

because $\|A - B\|_{Op} \leq \varepsilon\lambda_r \leq \frac{\varepsilon\mu_r}{1 - \varepsilon}$.

The Linear Program approach

Optimal Factorization from a Measure Theoretic Viewpoint

Let $S_1 = \{x \in \mathbb{C}^n, \|x\|_1 = 1\}$ denote the compact unit sphere with respect to the l^1 norm, and let $\mathcal{B}(S_1)$ denote the set of Borel measures over S_1 . For $A \in \text{Sym}(\mathbb{C}^n)^+(\mathbb{C}^n)$ consider the optimization problem:

$$(p^*, \mu^*) = \inf_{\mu \in \mathcal{B}(S_1): \int_{S_1} xx^* d\mu(x) = A} \mu(S_1) \quad (M)$$

Theorem (Optimal Measure)

For any $A \in \text{Sym}^+(\mathbb{C}^n)$ the optimization problem (M) is convex and its global optimum (minimum) is achieved by

$$p^* = \gamma_+(A) \quad , \quad \mu^*(x) = \sum_{k=1}^m \lambda_k \delta(x - g_k)$$

where $A = \sum_{k=1}^m (\sqrt{\lambda_k} g_k)(\sqrt{\lambda_k} g_k)^*$ is an optimal decomposition that achieves $\gamma_+(A) = \sum_{k=1}^m \lambda_k$.

Super-resolution and Convex Optimizations

$$\gamma_+(A) = \min_{x_1, \dots, x_m : A = \sum_k x_k x_k^*} \sum_{k=1}^m \|x_k\|_1^2, \quad m = n^2 \quad (P)$$

$$p^* = \inf_{\mu \in \mathcal{B}(S_1) : A = \int_{S_1} x x^* d\mu(x)} \int_{S_1} d\mu(x) \quad (M)$$

Remarks

- 1 *The optimization problem (P) is non-convex, but finite-dimensional. The optimization problem (M) is convex, but infinite-dimensional.*
- 2 *If $g_1, \dots, g_m \in S_1$ in the support of μ^* are known so that $\mu^* = \sum_{k=1}^m \lambda_k \delta(x - g_k)$, then the optimal $\lambda_1, \dots, \lambda_m \geq 0$ are determined by a linear program. More general, (M) is an infinite-dimensional linear program.*
- 3 *Finding the support of μ^* is an example of a super-resolution problem. One possible approach is to choose a redundant dictionary (frame) that includes the support of μ^* , and then solve the induced linear program.*

Proof of the Optimal Measure Result

Recall: we want to show the following problems admit the same solution:

$$\gamma_+(A) = \min_{x_1, \dots, x_m : A = \sum_k x_k x_k^*} \sum_{k=1}^m \|x_k\|_1^2, \quad m = n^2 \quad (P)$$

$$p^* = \inf_{\mu \in \mathcal{B}(S_1) : A = \int_{S_1} xx^* d\mu(x)} \int_{S_1} d\mu(x) \quad (M)$$

a. Assume $A = \sum_{k=1}^m x_k x_k^*$ is a global minimum for (P). Then

$\mu(x) = \sum_{k=1}^m \|x_k\|_1^2 \delta(x - \frac{x_k}{\|x_k\|_1})$ is a feasible solution for (M). This shows

$$p^* \leq \gamma_+(A).$$

b. For reverse: Let μ^* be an optimal measure in (M). Fix $\varepsilon > 0$. Construct a disjoint partition $(U_l)_{1 \leq l \leq L}$ of S_1 so that each U_l is included in some ball $B_\varepsilon(z_l)$ of radius ε with $\|z_l\|_1 = 1$. Thus $U_l \subset B_\varepsilon(z_l) \cap S_1$.

For each l , compute $x_l = \frac{1}{\mu^*(U_l)} \int_{U_l} x d\mu^*(x) \in B_\varepsilon(z_l)$. Let $g_l = \sqrt{\mu^*(U_l)} x_l$.

Proof: The Optimal Measure Result (cont)

Key inequality:

$$0 \leq R_l := \int_{U_l} (x - x_l)(x - x_l)^* d\mu^*(x) = \int_{U_l} xx^* d\mu^*(x) - \mu^*(U_l)x_l x_l^*$$

Sum over l and with $R = \sum_{l=1}^L R_l$ get

$$A = \sum_{l=1}^L \int_{U_l} xx^* d\mu^*(x) \leq \sum_{l=1}^L g_l g_l^* + R$$

By sub-additivity and homogeneity:

$$\gamma_+(A) \leq \sum_{l=1}^L \|g_l\|_1^2 + \gamma_+(R) \leq \sum_{l=1}^L \mu^*(U_l) \|x_l\|_1^2 + n \operatorname{trace}(R)$$

But $\|x_l - z_l\|_1 \leq \varepsilon$ and $\|x - x_l\|_1 \leq 2\varepsilon$ for every $x \in U_l$. Hence $\|x_l\|_1 \leq 1 + \varepsilon$ and $\operatorname{trace}(R_l) \leq 4\mu^*(U_l)\varepsilon^2$. (In fact, $\|x_l\|_1 \leq 1$ by triangle inequality)

Proof: The Optimal Measure Result (end)

Thus:

$$\gamma_+(A) \leq \mu^*(S_1) + (2\varepsilon + \varepsilon^2 + 4n\varepsilon^2)\mu^*(S_1)$$

Since $\varepsilon > 0$ is arbitrary, it follows

$$\gamma_+(A) \leq \mu^*(S_1) = p^*$$

This ends the proof of the measure result. \square

Third new result: Strong duality for γ_+

Theorem

For every $A \geq 0$,

$$\max_{\substack{T = T^* \\ \langle Tx, x \rangle \leq 1, \forall \|x\|_1 \leq 1}} \text{trace}(TA) = \min_{\substack{\mu \in \mathcal{B}(S_1) \\ \int_{S_1} xx^* d\mu(x) = A}} \mu(S_1) = \gamma_+(A)$$

Proof [Fushuai “Black” Jiang]

The second equality was established earlier as a “super-resolution” result.

For the first equality:

1. Let $A = \sum_{k=1}^m x_k x_k^*$ be its optimal decomposition such that

$\gamma_+(A) = \sum_{k=1}^m \|x_k\|_1^2$, and let $T = T^*$ be a generic matrix so that $\langle Ty, y \rangle \leq 1$ for all $\|y\|_1 \leq 1$. Denote $y_k = \frac{x_k}{\|x_k\|_1}$. Then

$$\text{trace}(TA) = \sum_{k=1}^m \langle Tx_k, x_k \rangle = \sum_{k=1}^m \|x_k\|_1^2 \langle Ty_k, y_k \rangle \leq \sum_{k=1}^m \|y_k\|_1^2 = \gamma_+(A)$$

Proof of strong duality for γ_+ (2)

2. For the reverse inequality, let $H \subset \text{Sym}^+(\mathbb{C}^n) \times \mathbb{R}$ denote the set

$$H = \left\{ \left(\int_{S_1} zz^* d\mu(z), r + \int_{S_1} d\mu \right), \mu \in \mathcal{B}(S_1), r \geq 0 \right\}$$

Claim 1: H is closed.

Use Banach-Alaoglu theorem that the set of unit Borel measures is weak-* compact.

Claim 2: H is convex. – immediate

Let $q = \max_{T=T^*} \text{trace}(TA)$ subject to $\langle Tx, x \rangle \leq 1$ for all $\|x\|_1 \leq 1$.

Claim 3: $(A, q) \in H$, which establishes the theorem.

Assume the contrary: $(A, q) \notin H$. Then it is separated by a hyperplane from H :

$$\text{trace} \left(R \int_{S_1} xx^* d\mu(z) \right) + a \left(r + \int_{S_1} d\mu \right) \geq c_0 > \text{trace}(AR) + aq, \quad \forall \mu \in \mathcal{B}(S_1), r \geq 0$$

Deduce: $a \geq 0, c_0 \leq 0$. If $a = 0$ then contradiction for $\mu = \mu^*$. Rescale by dividing through a . Denote $T_0 = -R/a$.

Proof of strong duality for γ_+ (3)

We obtained:

$$\int_{S_1} (1 - \langle T_0 x, x \rangle) d\mu \geq c_0 > q - \text{trace}(AT_0)$$

for every Borel measure $\mu \in \mathcal{B}(S_1)$. This means $\langle T_0 x, x \rangle \leq 1$ for all $\|x\| = 1$. This also implies $\langle T_0 x, x \rangle \leq 1$ for all $\|x\|_1 \leq 1$. On the other hand $q < \text{trace}(AT_0) + c_0 \leq \text{trace}(AT_0)$ which contradicts the optimality of q . Q.E.D.