

I.

1. First note that

$$(x^4 - 1) \cdot \delta(x) = -\delta(x)$$

Hence $-\delta(x)$ is one particular solution. Thus the general solution we seek must satisfy:

$$(x^4 - 1) \cdot (f(x) + \delta(x)) = 0$$

Since $x^4 - 1 = (x^2 + 1)(x + 1)(x - 1)$ we obtain, first that

$$(x - 1)(x + 1)(f(x) + \delta(x)) = 0$$

And then

$$f(x) + \delta(x) = c_1\delta(x - 1) + c_2\delta(x + 1)$$

Hence the general solution is:

$$f(x) = -\delta(x) + c_1\delta(x - 1) + c_2\delta(x + 1)$$

2. One particular solution is δ since $(x^4 + 1) \cdot \delta(x) = \delta(x)$. Thus f must satisfy:

$$(x^4 + 1) \cdot (f(x) - \delta(x)) = 0$$

Since $x^4 + 1$ does not have real roots, the only solution (multiply both sides by $\frac{1}{1+x^4}$) is given by

$$f(x) - \delta(x) = 0$$

Hence:

$$f(x) = \delta(x)$$

is the unique solution.

II.

3. First note: $f(x) = e^{2\pi i x} \cdot \delta(x) = \delta(x)$. Thus

$$\hat{f}(s) = \hat{\delta}(s) = 1$$

4. Note $f(x) = \sin^2(\pi x) = \frac{1 - \cos(2\pi x)}{2} = \frac{1}{2} - \frac{1}{4}e^{2\pi i x} - \frac{1}{4}e^{-2\pi i x}$. Thus

$$\hat{f}(s) = \frac{1}{2}\delta(s) - \frac{1}{4}\delta(s - 1) - \frac{1}{4}\delta(s + 1)$$

5. First note $f\{\Phi\} = \delta'\{x^2\Phi\} = -\delta\{2x\Phi + x^2\Phi'\} = 0$. Thus $f = 0$. Hence

$$\hat{f}(s) = 0$$

6. Recall $\widehat{p_{-1}}(s) = -i\pi \operatorname{sign}(s)$ and $\widehat{\operatorname{sign}}(s) = \frac{1}{i\pi}p_{-1}(s)$. Thus

$$\hat{f}(s) = \widehat{\operatorname{sign}}(s - 2) = -\frac{i}{\pi}p_{-1}(s - 2)$$

7. Note $f(x) = \frac{1}{4}\frac{1}{x-2} - \frac{1}{4}\frac{1}{x+2} = \frac{1}{4}p_{-1}(x - 2) - \frac{1}{4}p_{-1}(x + 2)$. Thus

$$\begin{aligned} \hat{f}(s) &= \frac{1}{4}e^{-4\pi i s}\widehat{p_{-1}}(s) - \frac{1}{4}e^{4\pi i s}\widehat{p_{-1}}(s) = -\frac{i\pi}{4}e^{-4\pi i s}\operatorname{sign}(s) + \frac{i\pi}{4}e^{4\pi i s}\operatorname{sign}(s) = \\ &= -\frac{\pi}{2}\sin(4\pi s)\operatorname{sign}(s) = -\frac{\pi}{2}\sin(4\pi|s|) \end{aligned}$$

III

8. Fourier transform the equation and obtain

$$((2\pi i s)^4 - 1) \cdot F(s) = 1$$

Thus

$$F_0(s) = \frac{1}{(2\pi)^4 s^4 - 1}$$

is a particular solution of this algebraic equation. Since $(2\pi)^4 s^4 - 1 = (4\pi^2 s^2 + 1)(2\pi s - 1)(2\pi s + 1)$ the general solution is given by

$$F(s) = \frac{1}{16\pi^4 s^4 - 1} + c_1 \delta\left(s - \frac{1}{2\pi}\right) + c_2 \delta\left(s + \frac{1}{2\pi}\right)$$

Next we need to Fourier invert this distribution. First:

$$\begin{aligned} \frac{1}{16\pi^4 s^4 - 1} &= \frac{1}{2} \frac{1}{4\pi^2 s^2 - 1} - \frac{1}{2} \frac{1}{4\pi^2 s^2 + 1} = \frac{1}{4} \frac{1}{2\pi s - 1} - \frac{1}{4} \frac{1}{2\pi s + 1} - \frac{1}{2} \frac{1}{4\pi^2 s^2 + 1} = \\ &= \frac{1}{8\pi} p_{-1}\left(s - \frac{1}{2\pi}\right) - \frac{1}{8\pi} p_{-1}\left(s + \frac{1}{2\pi}\right) - \frac{1}{4} \frac{2}{1 + 4\pi^2 s^2} \end{aligned}$$

Thus its Fourier transform inverse is

$$f_0(x) = \frac{1}{8} e^{ix} \pi i \operatorname{sign}(x) - \frac{1}{8} e^{-ix} \pi i \operatorname{sign}(x) - \frac{1}{4} e^{-|x|} = -\frac{\pi}{4} \sin(|x|) - \frac{1}{4} e^{-|x|}$$

Then the general solution becomes

$$f(x) = -\frac{\pi}{4} \sin(|x|) - \frac{1}{4} e^{-|x|} + c_1 e^{ix} + c_2 e^{-ix}$$

9. Fourier transform the equation and obtain

$$((2\pi i s)^4 + 1) \cdot F(s) = 1$$

or

$$((16\pi^4 s^4 + 1) \cdot F(s) = 1$$

The unique solution is

$$F(s) = \frac{1}{16\pi^4 s^4 + 1} = \sum_{k=0}^3 \frac{\alpha_k}{2\pi s - z_k}$$

where $z_k = e^{i(2k+1)\pi/4}$ are the 4 roots of -1 , and α_k are obtained from:

$$\alpha_k = \lim_{2\pi s \rightarrow z_k} \frac{2\pi s - z_k}{16\pi^4 s^4 + 1} = \frac{1}{4} e^{-i3\pi(2k+1)/4} = \frac{z_k^3}{4} = -\frac{\bar{z}_k}{4}$$

Explicitly:

$$F(s) = -\frac{1}{4} \frac{\bar{z}_0}{2\pi s - z_0} - \frac{1}{4} \frac{\bar{z}_1}{2\pi s - z_1} - \frac{1}{4} \frac{\bar{z}_2}{2\pi s - z_2} - \frac{1}{4} \frac{\bar{z}_3}{2\pi s - z_3}$$

Finally we need to Fourier invert this function. Note for a real and $b > 0$,

$$F_+(s) = \frac{1}{2\pi s - (a - ib)} \rightarrow f_+(x) = -ie^{iax} e^{bx} h(-x) = -ie^{i(a-ib)x} h(-x)$$

$$F_-(s) = \frac{1}{2\pi s - (a + ib)} \rightarrow f_-(x) = ie^{iax} e^{-bx} h(x) = ie^{i(a+ib)x} h(x)$$

where $h(x)$ is the Heaviside step function (see equations (22) in Chapter 3). Note $\operatorname{imag}(z_0) = \operatorname{imag}(z_1) > 0$ and $\operatorname{imag}(z_2) = \operatorname{imag}(z_3) < 0$. Thus

$$f(x) = -\frac{\bar{z}_0}{4} e^{iz_0 x} i h(x) - \frac{\bar{z}_1}{4} e^{iz_1 x} i h(x) - \frac{\bar{z}_2}{4} e^{iz_2 x} (-i) h(-x) - \frac{\bar{z}_3}{4} e^{iz_3 x} (-i) h(-x)$$

To simplify this expression, note $z_2 = -z_0$ and $z_3 = -z_1$, hence

$$f(x) = i \frac{\bar{z}_2}{4} e^{-1z_2|x|} + i \frac{\bar{z}_3}{4} e^{-iz_3|x|} = i \frac{-\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2}}{4} e^{-i(-\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2})|x|} + i \frac{\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2}}{4} e^{-i(\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2})|x|} =$$

$$= e^{-\frac{\sqrt{2}}{2}|x|} \left[\frac{\sqrt{2}}{8}(1-i)e^{i\frac{\sqrt{2}}{2}|x|} + \frac{\sqrt{2}}{8}(1+i)e^{-i\frac{\sqrt{2}}{2}|x|} \right] = \frac{\sqrt{2}}{4} e^{-\frac{\sqrt{2}}{2}|x|} \left[\cos\left(\frac{\sqrt{2}}{2}|x|\right) + \sin\left(\frac{\sqrt{2}}{2}|x|\right) \right]$$

Note that $f(x)$ is a real valued continuous function. To test:

$$Df(x) = -\frac{1}{2} e^{-\frac{\sqrt{2}}{2}|x|} \sin\left(\frac{\sqrt{2}}{2}|x|\right) \operatorname{sign}(x)$$

$$D^2 f(x) = \frac{\sqrt{2}}{4} e^{-\frac{\sqrt{2}}{2}|x|} \left(\sin\left(\frac{\sqrt{2}}{2}|x|\right) - \cos\left(\frac{\sqrt{2}}{2}|x|\right) \right)$$

$$D^3 f(x) = \frac{1}{2} e^{-\frac{\sqrt{2}}{2}|x|} \cos\left(\frac{\sqrt{2}}{2}|x|\right) \operatorname{sign}(x)$$

$$D^4 f(x) = -\frac{\sqrt{2}}{4} e^{-\frac{\sqrt{2}}{2}|x|} \left[\cos\left(\frac{\sqrt{2}}{2}|x|\right) + \sin\left(\frac{\sqrt{2}}{2}|x|\right) \right] + \delta(x)$$

IV.

10.

First we solve the equation

$$(D^2 - 1)g(x) = \delta(x)$$

Fourier transform to obtain

$$((2\pi is)^2 - 1)G(s) = 1$$

or

$$(4\pi^2 s^2 + 1)G(s) = -1$$

The unique solution is

$$G(s) = -\frac{1}{4\pi s^2 + 1}$$

whose inverse Fourier transform is

$$g(x) = -\frac{1}{2} e^{-|x|}$$

Then the solution of $(D^2 - 1)f = \sin(x)$ is $f = g * \sin$:

$$f(x) = -\frac{1}{2} \int_{-\infty}^{\infty} e^{-|y|} \sin(x-y) dy = -\frac{1}{2} \sin(x) \int_{-\infty}^{\infty} e^{-|y|} \cos(y) dy + \frac{1}{2} \cos(x) \int_{-\infty}^{\infty} e^{-|y|} \sin(y) dy$$

The two integrals are:

$$\int_{-\infty}^{\infty} e^{-|y|} \sin(y) dy = 0$$

because the integrand is odd, and

$$\int_{-\infty}^{\infty} e^{-|y|} \cos(y) dy = 2 \int_0^{\infty} e^{-y} \cos(y) dy = 2I$$

where, integrating by parts,

$$I = \int_0^{\infty} e^{-y} \cos(y) dy = 1 - \int_0^{\infty} e^{-y} \sin(y) dy = 1 - (0 + \int_0^{\infty} e^{-y} \cos(y) dy) = 1 - I$$

Hence $I = 1/2$ and

$$\int_{-\infty}^{\infty} e^{-|y|} \cos(y) dy = 1$$

Assembling the result we get:

$$f(x) = -\frac{1}{2} \sin(x)$$