

A Stochastic Galerkin Method for the Boltzmann Equation with High Dimensional Random Inputs Using Sparse Grids

Ruiwen Shu*, Jingwei Hu†, Shi Jin‡

August 20, 2016

Abstract

We propose a stochastic Galerkin method using sparse grids for the Boltzmann equation with high dimensional random inputs. The method uses locally supported piecewise polynomials as an orthonormal basis of the random space. By a sparse grid technique, only a moderate number of basis functions are required to achieve good accuracy in high dimensional random spaces. We discover a sparse structure of a set of basis-related coefficients, which allows us to accelerate the computation of the collision operator. Regularity of the solution of the Boltzmann equation in the random space and an accuracy result of the stochastic Galerkin method are proved in multidimensional case. The efficiency of the method is illustrated by numerical examples with uncertainties from the initial data, boundary data and collision kernel.

Key words. Uncertainty quantification, Boltzmann equation, stochastic Galerkin methods, sparse grids

1 Introduction

The Boltzmann equation plays an essential role in kinetic theory [8]. It describes the time evolution of the density distribution of dilute gases, where fluid dynamics equations, such as the Euler equations and the Navier-Stokes equations, fail to provide reliable information. It is an indispensable tool in fields concerning non-equilibrium statistical mechanics, such as rarefied gas dynamics and astronautical engineering.

*Department of Mathematics, University of Wisconsin-Madison, Madison, WI 53706, USA (rshu2@math.wisc.edu).

†Department of Mathematics, Purdue University, West Lafayette, IN 47907, USA (jingwei.hu@purdue.edu). This author's research was partially supported by NSF grant DMS-1620250 and a startup grant from Purdue University.

‡Department of Mathematics, University of Wisconsin-Madison, Madison, WI 53706, USA (sjin@wisc.edu) and Institute of Natural Sciences, School of Mathematical Science, MOELSEC and SHL-MAC, Shanghai Jiao Tong University, Shanghai 200240, China. This author's research was supported by NSF grants DMS-1522184 and DMS-1107291: RNMS KI-Net, by NSFC grant No. 91330203, and by the Office of the Vice Chancellor for Research and Graduate Education at the University of Wisconsin-Madison with funding from the Wisconsin Alumni Research Foundation.

For most applications of the Boltzmann equation, the initial and boundary data are given by physical measurements, which inevitably bring measurement errors. Furthermore, due to the difficulty of deriving the collision kernels from first principles, empirical collision kernels are often used. Such kernels contain adjustable parameters, which are determined by matching with experimental data [5]. This procedure involves uncertainty on the parameters in the collision kernel. To understand the impact of these random inputs on the solution of the Boltzmann equation, it is imperative to incorporate the uncertainties into the equation, and design numerical methods to solve the resulting system [24]. A proper quantification of uncertainty will provide reliable predictions and a guidance for improving the models. Since the uncertainties of the Boltzmann equations come from many independent sources, it is necessary to use a high dimensional random space to incorporate all the uncertainties. Moreover, a Karhunen-Loeve expansion of a random field will result in a high dimensional random space.

Various numerical methods have been developed to solve the problem of uncertainty quantification (UQ) [24, 11, 19, 25]. Monte-Carlo methods [20] use statistical sampling in the random space, which give halft order convergence in any dimension. Stochastic collocation methods [2, 4] take sampling points on a well-designed grid, usually according to a quadrature rule, and the statistical moments are computed by numerical quadrature. Stochastic Galerkin methods [4, 3] use an orthonormal basis expansion in the random space. By a truncation of the expansion and Galerkin projection, one is led to a deterministic system of expansion coefficients. Both methods can achieve spectral accuracy in one-dimensional random space if the quadrature rule (orthonormal basis) is well chosen.

Hu and Jin [16] gave a first numerical method to solve the Boltzmann equation with uncertainty by a general polynomial chaos based stochastic Galerkin method. By a singular value decomposition on a set of coefficients related to the basis functions, the computational cost of the collision kernel is decreased dramatically. However, their work focuses on low dimensional random spaces, and a direct extension of their method to high dimensional random spaces will suffer from the curse of dimensionality, which means K , the number of basis functions, will grow like $K = \binom{K_1+d}{K_1}$, where K_1 is the number of basis in one dimension, and d is the dimension of the random space. This cost is not affordable if both K_1 and d are large. Monte-Carlo methods are feasible, but a halft order convergence rate can be unsatisfactory in many applications. Therefore it is desirable to have an efficient and accurate method to solve the Boltzmann equation with high dimensional random inputs.

In this work, we adopt a sparse grid approach [15, 10] for the stochastic Galerkin method to circumvent the curse of dimensionality. The idea of sparse grids traces back to Smolyak [22]. In recent years, sparse grids have become a major approach to break the curse of dimensionality in various contexts, for example in Galerkin finite element methods [15, 27], finite difference methods [12, 13] and high-dimensional stochastic differential equations [26, 21]. We adopt the sparse grid method proposed by Guo and Cheng [14], who use the method for a discrete Galerkin method for transport equations. Simply speaking, we start from a hierarchical basis in one dimension. To construct the sparse grid basis in multi-dimension, we take the tensor grid and discard those basis functions that are in deep levels in most dimensions. In this way only a small number of basis functions are kept, yet it can be proved that the accuracy is still as good as the corresponding tensor grid, if the function to approximate is smooth enough. With a hierarchical basis with N levels and piecewise polynomials of degree at most m , our method can achieve an

accuracy of $O(N^d 2^{-N(m+1)})$ with number of basis $K = O((m+1)^d 2^N N^{d-1})$ for d -dimensional random spaces. This accuracy is $O(K^{-(m+1)}(\log K)^{(m+2)(d-1)})$ in terms of K . It is algebraically accurate, but as d increases, the accuracy deteriorates very slowly. Furthermore, we discover a sparse structure of a set of basis related coefficients, S_{ijk} , which greatly reduces the cost of the expensive collision operator evaluation.

The paper is organized as follows: in Section 2 we introduce the Boltzmann equation with uncertainty and the framework of stochastic Galerkin (sG) method; in Section 3 we introduce our sparse grid method with multi-wavelet functions; in Section 4 we give an estimate of the sparsity of the coefficients S_{ijk} ; in Section 5 we prove the random space regularity of the solution of the Boltzmann equation with uncertainty, as well as the accuracy of the sG method with sparse grid; in Section 6 we give some numerical results.

2 The Boltzmann Equation with Uncertainty

The classical (deterministic) Boltzmann equation in its dimensionless form reads

$$\partial_t f + \mathbf{v} \cdot \nabla_{\mathbf{x}} f = \frac{1}{\text{Kn}} Q(f, f), \quad (2.1)$$

where $f = f(t, \mathbf{x}, \mathbf{v})$ is the density distribution function of a dilute gas at time $t \in \mathbb{R}^+$, position $\mathbf{x} \in \Omega \subset \mathbb{R}^{d_x}$, and with particle velocity $\mathbf{v} \in \mathbb{R}^{d_v}$. Kn is the Knudsen number, a dimensionless number defined as the ratio of the mean free path and a typical length scale, such as the size of the spatial domain. The collision operator $Q(f, f)$ is given by

$$Q(f, f) = \int_{\mathbb{R}^{d_v}} \int_{S^{d_v-1}} B(\mathbf{v}, \mathbf{v}_*, \sigma) [f(\mathbf{v}')f(\mathbf{v}'_*) - f(\mathbf{v})f(\mathbf{v}_*)] d\sigma d\mathbf{v}_*, \quad (2.2)$$

which is a quadratic integral operator modeling the binary elastic collision between particles. $(\mathbf{v}, \mathbf{v}_*)$ and $(\mathbf{v}', \mathbf{v}'_*)$ are the particle velocities before and after a collision, which are given by

$$\begin{cases} \mathbf{v}' = \frac{\mathbf{v} + \mathbf{v}_*}{2} + \frac{|\mathbf{v} - \mathbf{v}_*|}{2} \sigma, \\ \mathbf{v}'_* = \frac{\mathbf{v} + \mathbf{v}_*}{2} - \frac{|\mathbf{v} - \mathbf{v}_*|}{2} \sigma, \end{cases} \quad (2.3)$$

with a vector σ varying on the unit sphere. The collision kernel B is a non-negative function of the form $B(\mathbf{v}, \mathbf{v}_*, \sigma) = B(|\mathbf{v} - \mathbf{v}_*|, \cos \theta)$, where $\theta = \arccos \frac{\sigma \cdot (\mathbf{v} - \mathbf{v}_*)}{|\mathbf{v} - \mathbf{v}_*|}$ is the deviation angle. A commonly used model for the collision kernel is the variable hard sphere (VHS) model [5], which takes the form

$$B = b|\mathbf{v} - \mathbf{v}_*|^\lambda, \quad (2.4)$$

where b and λ are some constants, whose values are usually determined by matching with the experimental data to reproduce the correct transport coefficients such as the viscosity.

The Boltzmann collision operator satisfies the conservation laws

$$\int Q(f, f) \begin{pmatrix} 1 \\ \mathbf{v} \\ |\mathbf{v}|^2 \end{pmatrix} d\mathbf{v} = 0, \quad (2.5)$$

as well as the H -theorem

$$- \int Q(f, f) \ln f \, d\mathbf{v} \geq 0. \quad (2.6)$$

The equality is achieved if and only if f takes the form

$$M(\mathbf{v})_{(\rho, \mathbf{u}, T)} = \frac{\rho}{(2\pi T)^{d_v/2}} e^{-\frac{(\mathbf{v}-\mathbf{u})^2}{2T}}, \quad (2.7)$$

which is called the Maxwellian. ρ , \mathbf{u} and T are the density, bulk velocity and temperature, given by

$$\rho = \int f \, d\mathbf{v}, \quad \mathbf{u} = \frac{1}{\rho} \int f \mathbf{v} \, d\mathbf{v}, \quad T = \frac{1}{d_v \rho} \int f |\mathbf{v} - \mathbf{u}|^2 \, d\mathbf{v}. \quad (2.8)$$

The initial condition of the Boltzmann equation is given by

$$f(0, \mathbf{x}, \mathbf{v}) = f^0(\mathbf{x}, \mathbf{v}), \quad (2.9)$$

and a boundary condition is needed if the spatial domain Ω is a proper subset of \mathbb{R}^{d_x} . We adopt the Maxwell boundary condition, which takes the form

$$f(t, \mathbf{x}, \mathbf{v}) = g(t, \mathbf{x}, \mathbf{v}), \quad \mathbf{x} \in \partial\Omega, \quad \mathbf{v} \cdot \mathbf{n} > 0, \quad (2.10)$$

with

$$\begin{aligned} g(t, \mathbf{x}, \mathbf{v}) = & (1 - \alpha) f(t, \mathbf{x}, \mathbf{v} - 2(\mathbf{v} \cdot \mathbf{n})\mathbf{n}) \\ & + \frac{\alpha}{(2\pi)^{(d_v-1)/2} T_w^{(d_v+1)/2}} e^{-\frac{|\mathbf{v}|^2}{2T_w}} \int_{\mathbf{v} \cdot \mathbf{n} < 0} f(t, \mathbf{x}, \mathbf{v}) |\mathbf{v} \cdot \mathbf{n}| \, d\mathbf{v}, \end{aligned} \quad (2.11)$$

where T_w is the temperature of the wall, and \mathbf{n} is the inner normal unit vector of the wall. The first term is the specular reflective part, and the second term is the diffusive part. α is the accommodation coefficient. $\alpha = 1$ implies purely diffusive boundary, while $\alpha = 0$ implies purely reflective boundary. For simplicity we only consider the case where the wall is static.

As mentioned before, there are many sources of uncertainties in the Boltzmann equation, such as the initial data, boundary data, and collision kernel. To quantify these uncertainties we introduce the Boltzmann equation with uncertainty

$$\begin{cases} \partial_t f(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) + \mathbf{v} \cdot \nabla_{\mathbf{x}} f(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) = \frac{1}{\text{Kn}} Q_{\mathbf{z}}(f, f), & t \in \mathbb{R}_+, \mathbf{x} \in \Omega, \mathbf{v} \in \mathbb{R}^{d_v}, \mathbf{z} \in I_{\mathbf{z}} \subset \mathbb{R}^d, \\ f(0, \mathbf{x}, \mathbf{v}, \mathbf{z}) = f^0(\mathbf{x}, \mathbf{v}, \mathbf{z}), \\ f(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) = g(t, \mathbf{x}, \mathbf{v}, \mathbf{z}), & \mathbf{x} \in \partial\Omega. \end{cases} \quad (2.12)$$

Here $\mathbf{z} \in I_{\mathbf{z}}$ is a d -dimensional random vector with probability distribution $\pi(\mathbf{z})$ characterizing the uncertainty in the system. We assume that the collision kernel has the form

$$B(\mathbf{v}, \mathbf{v}_*, \sigma, \mathbf{z}) = b(\mathbf{z}) B_0(\mathbf{v}, \mathbf{v}_*, \sigma),$$

which means that $Q_{\mathbf{z}}$ can be written as

$$Q_{\mathbf{z}}(f, f) = b(\mathbf{z}) Q(f, f).$$

The Maxwell boundary data $g(t, \mathbf{x}, \mathbf{v}, \mathbf{z})$ is given by

$$\begin{aligned} g(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) = & (1 - \alpha(\mathbf{z})) f(t, \mathbf{x}, \mathbf{v} - 2(\mathbf{v} \cdot \mathbf{n})\mathbf{n}, \mathbf{z}) \\ & + \frac{\alpha(\mathbf{z})}{(2\pi)^{(d_v-1)/2} T_w(\mathbf{z})^{(d_v+1)/2}} e^{-\frac{|\mathbf{v}|^2}{2T_w(\mathbf{z})}} \int_{\mathbf{v} \cdot \mathbf{n} < 0} f(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) |\mathbf{v} \cdot \mathbf{n}| \, d\mathbf{v}. \end{aligned} \quad (2.13)$$

To solve the stochastic system (2.12), Hu and Jin [16] proposed a stochastic Galerkin method (sG). The idea is to approximate f by a truncated Galerkin expansion:

$$f(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) \approx f^K(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) = \sum_{k=0}^K \hat{f}_k(t, \mathbf{x}, \mathbf{v}) \Phi_k(\mathbf{z}), \quad (2.14)$$

where $\{\Phi_k(\mathbf{z})\}$ are an orthonormal polynomial basis, with $\Phi_k(\mathbf{z})$ being the polynomial of degree k in the random domain, which satisfies

$$\int_{I_{\mathbf{z}}} \Phi_i(\mathbf{z}) \Phi_j(\mathbf{z}) \pi(\mathbf{z}) \, d\mathbf{z} = \delta_{ij}.$$

Substituting (2.14) into (2.12) and conducting a standard Galerkin projection, one gets

$$\partial_t \hat{f}_k(t, \mathbf{x}, \mathbf{v}) + \mathbf{v} \cdot \nabla_{\mathbf{x}} \hat{f}_k(t, \mathbf{x}, \mathbf{v}) = Q_k(f^K, f^K), \quad (2.15)$$

$$\hat{f}_k(0, \mathbf{x}, \mathbf{v}) = \hat{f}_k^0(\mathbf{x}, \mathbf{v}), \quad (2.16)$$

$$Q_k(f^K, f^K) = \sum_{i,j=0}^K S_{ijk} Q(\hat{f}_i, \hat{f}_j), \quad (2.17)$$

where

$$S_{ijk} = \int_{I_{\mathbf{z}}} b(\mathbf{z}) \Phi_i(\mathbf{z}) \Phi_j(\mathbf{z}) \Phi_k(\mathbf{z}) \pi(\mathbf{z}) \, d\mathbf{z}. \quad (2.18)$$

The boundary condition is given by

$$\begin{aligned} \hat{g}_k &= \sum_{i,j=0}^K S_{0,ijk} (1 - \hat{\alpha}_i) \hat{f}_j(t, \mathbf{x}, \mathbf{v} - 2(\mathbf{v} \cdot \mathbf{n})\mathbf{n}) \\ &\quad + \sum_{j=0}^K D_{kj}(\mathbf{x}, \mathbf{v}) \int_{(\mathbf{v} \cdot \mathbf{n} < 0} \hat{f}(t, \mathbf{x}, \mathbf{v}, \mathbf{z}) |\mathbf{v} \cdot \mathbf{n}| \, d\mathbf{v}, \end{aligned} \quad (2.19)$$

where

$$D_{kj}(\mathbf{x}, \mathbf{v}) = \int_{I_{\mathbf{z}}} \frac{\alpha(\mathbf{z})}{(2\pi)^{(d_v-1)/2} T_w(\mathbf{z})^{(d_v+1)/2}} e^{-\frac{|\mathbf{v}|^2}{2T_w(\mathbf{z})}} \Phi_k(\mathbf{z}) \Phi_j(\mathbf{z}) \pi(\mathbf{z}) \, d\mathbf{z}, \quad (2.20)$$

is a matrix that is time independent hence can be pre-computed.

This gPC-sG method works well for low dimensional random inputs, but for high dimensional ones, it might require a very large number of basis functions (K large) to approximate f to a given accuracy. If one takes K_1 basis functions in one dimensional random space, then a direct extension of the gPC-sG method will require $\binom{K_1+d}{K_1}$ basis functions, which is prohibitively expensive if both K_1 and d are large. Furthermore, since the computation of the $\{Q_k\}_{k=0}^K$ requires $O(K^2)$ times evaluation of the deterministic collision operator in general, one has to choose a relatively small K in order to afford the computation of $\{Q_k\}_{k=0}^K$. Also, [16] uses a singular value decomposition of size K as a pre-computation for the collision operator, which reduces the computational cost by one order of magnitude, but it is still too expensive for high-dimensional random inputs. In the following sections we propose a stochastic Galerkin method with sparse grid basis functions, which requires much fewer basis functions for high dimensional random space.

3 A Sparse Grid Method with Multi-wavelet Functions

3.1 The Sparse Grid Construction

For simplicity we restrict to the case $I_{\mathbf{z}} = [-1, 1]^d$, and $\pi(\mathbf{z}) = \frac{1}{2^d}$ is the uniform distribution. We follow the notation by Guo and Cheng [14]. We start by constructing a hierarchical decomposition of the space consisting of piecewise polynomials of order at most m . Let $P^m(a, b)$ be the space of polynomials of order at most m on the interval (a, b) , and for every $N \geq 0$,

$$V_N^m = \{\phi : \phi \in P^m(-1 + 2^{-N+1}j, -1 + 2^{-N+1}(j+1)), j = 0, 1, \dots, 2^N - 1\}. \quad (3.1)$$

Then define the wavelet space $W_N^m, N = 1, 2, \dots$ as the orthogonal complement of V_{N-1}^m inside V_N^m . For convenience we define $W_0^m = V_0^m$. Then one gets the hierarchical decomposition $V_N^m = \bigoplus_{0 \leq j \leq N} W_j^m$.

Then a standard sparse grid trick can be applied. For simplicity we introduce the following vector notations:

If $\mathbf{i} = (i_1, \dots, i_d)$, $\mathbf{j} = (j_1, \dots, j_d)$ then

$$\mathbf{i} \leq \mathbf{j} \text{ means } i_1 \leq j_1, \dots, i_d \leq j_d,$$

$$\binom{\mathbf{j}}{\mathbf{i}} := \binom{j_1}{i_1} \times \dots \times \binom{j_d}{i_d},$$

$\mathbf{1}_m$ is the vector with 1 at m -th component and 0 at others,

$$|\mathbf{i}|_\infty = \max_m \{|i_1|, \dots, |i_d|\}, \quad |\mathbf{i}|_1 = |i_1| + \dots + |i_d|.$$

Define the d -fold tensor product of V_N^m by

$$\mathbf{V}_{N,\mathbf{z}}^m = V_{N,z_1}^m \times \dots \times V_{N,z_d}^m. \quad (3.2)$$

Similarly define the d -fold tensor product of W_j^m by

$$\mathbf{W}_{\mathbf{j},\mathbf{z}}^m = W_{j_1,z_1}^m \times \dots \times W_{j_d,z_d}^m. \quad (3.3)$$

Then

$$\mathbf{V}_{N,\mathbf{z}}^m = \bigoplus_{0 \leq |\mathbf{j}|_\infty \leq N} \mathbf{W}_{\mathbf{j},\mathbf{z}}^m.$$

The sparse grid trick is to replace the l^∞ norm on \mathbf{j} by the l^1 norm. In this way we define the sparse grid space

$$\hat{\mathbf{V}}_{N,\mathbf{z}}^m = \bigoplus_{0 \leq |\mathbf{j}|_1 \leq N} \mathbf{W}_{\mathbf{j},\mathbf{z}}^m. \quad (3.4)$$

From now on we will omit the subscript \mathbf{z} for these spaces.

3.2 Construction of the Basis Functions

We adopt the basis functions of W_j^m constructed by Alpert [1]. The basis functions of W_j^m are denoted by $\psi_{j,l}^{m'}$, $m' = 0, 1, \dots, m$, $l = 0, 1, \dots, 2^{j-1} - 1$ for $j \geq 1$ and $l = 0$ for $j = 0$. $\psi_{0,0}^{m'}$ are the orthonormal Legendre polynomials of degree m' on $[-1, 1]$, and $\psi_{1,0}^{m'}$ are piecewise polynomials on $[-1, 0]$ and $[0, 1]$ that are orthogonal to those Legendre polynomials, which can

be constructed by a procedure similar to the Gram-Schmidt orthogonalization. Other $\psi_{j,l}^{m'}$ are defined by dilation and translation of the $\psi_{1,0}^{m'}$:

$$\psi_{j,l}^{m'}(y) = 2^{(j-1)/2} \psi_{1,0}^{m'}(2^{j-1}y + 2^{j-1} - 1 - 2l), \quad j = 2, 3, \dots, \quad l = 0, 1, \dots, 2^{j-1} - 1,$$

which has support on the interval $[-1 + 2^{2-j}l, -1 + 2^{2-j}(l+1)]$.

The basis functions of $\mathbf{W}_{\mathbf{j}}^m$ are tensor products of the one dimensional basis functions:

$$\psi_{\mathbf{j},\mathbf{l}}^{\mathbf{m}'}(\mathbf{z}) = \psi_{j_1,l_1}^{m'_1}(z_1) \times \dots \times \psi_{j_d,l_d}^{m'_d}(z_d), \quad 0 \leq |\mathbf{m}'|_\infty \leq m, 0 \leq l_1 \leq 2^{j_1-1} - 1, \dots, 0 \leq l_d \leq 2^{j_d-1} - 1,$$

and the basis functions of $\hat{\mathbf{V}}_N^m$ consist of all the above functions for $0 \leq |\mathbf{j}|_1 \leq N$. By reordering the basis functions for $\hat{\mathbf{V}}_N^m$ we make them $\Phi_0(\mathbf{z}), \dots, \Phi_K(\mathbf{z})$, where $K = K(m, N, d)$ is the total number of basis functions minus 1. It is proved in Lemma 2.3 of [23] that

$$K = O((m+1)^d 2^N N^{d-1}). \quad (3.5)$$

4 Estimate of the Sparsity of S_{ijk}

Recall the triple product tensor S_{ijk} defined in (2.18). Due to the local support of the sparse grid basis functions Φ_k , this tensor is sparse, especially when N and d are large. Due to this sparsity, when one computes $Q_k = \sum_{i,j=0}^K S_{ijk} Q(\hat{f}_i, \hat{f}_j)$, one only needs to compute those $Q(\hat{f}_i, \hat{f}_j)$ where there is at least one k with $S_{ijk} \neq 0$. Now we prove some results on its sparsity. We focus on the dependence on N , so every $O(\cdot)$ notation means multiplication by a constant that may depend on d .

Recall that when one takes the sparse grid space $\hat{\mathbf{V}}_N^m$, the basis functions are

$$\begin{aligned} \psi_{\mathbf{j},\mathbf{l}}^{\mathbf{m}'}(\mathbf{z}) &= \psi_{j_1,l_1}^{m'_1}(z_1) \times \dots \times \psi_{j_d,l_d}^{m'_d}(z_d), \quad 0 \leq |\mathbf{m}'|_\infty \leq m, \\ 0 \leq l_1 \leq 2^{j_1-1} - 1, \dots, 0 \leq l_d \leq 2^{j_d-1} - 1, \quad &|\mathbf{j}|_1 \leq N. \end{aligned} \quad (4.1)$$

The function $\psi_{j,l}^{m'}(z)$ is supported on the interval $[-1 + 2^{2-j}l, -1 + 2^{2-j}(l+1)]$ for $j \geq 1$. Since this support is independent of m' , we omit the m' index in the following consideration. If ψ_{j^1,l^1} and ψ_{j^2,l^2} have non-intersecting supports, then

$$\int_{I_{\mathbf{z}}} b(\mathbf{z}) \psi_{j^1,l^1}(\mathbf{z}) \psi_{j^2,l^2}(\mathbf{z}) \psi_{j^3,l^3}(\mathbf{z}) \pi(\mathbf{z}) d\mathbf{z} = 0, \quad \forall \mathbf{j}_3, \mathbf{l}_3.$$

Recall that the number of basis functions, in $\hat{\mathbf{V}}_N^m$, which includes those $\psi_{\mathbf{j},\mathbf{l}}$ with $|\mathbf{j}|_1 \leq N$ and $0 \leq l_1 \leq 2^{j_1-1} - 1, \dots, 0 \leq l_d \leq 2^{j_d-1} - 1$, is $O((m+1)^d 2^N N^{d-1})$. Thus the number of the pairs of such functions is $O((m+1)^{2d} 2^{2N} N^{2d-2})$. Now we state our result:

Theorem 4.1. *The pairs of basis functions of $\hat{\mathbf{V}}_N^m$ with intersecting supports have a total number at most $O((m+1)^{2d} 2^{2N} N^{d+1})$.*

Proof. The number of $\phi_{j,l}$ for a fixed j is $(m+1)2^{j-1}$ for $j \geq 1$, and $m+1$ if $j = 0$. Thus it is less than or equal to $(m+1)2^j$ for all j . For fixed j^1, j^2 , suppose $j^1 \geq j^2$, then ϕ_{j^1,l^1} and ϕ_{j^2,l^2} have intersecting supports if and only if the support of ϕ_{j^1,l^1} is a subinterval of the support of ϕ_{j^2,l^2} . For every l^1 , there is one and only one such l^2 . Thus the number of pairs l^1, l^2 such that ϕ_{j^1,l^1} and ϕ_{j^2,l^2} have intersecting supports is 2^{j^1} , which is $2^{\max\{j^1, j^2\}}$ in general.

Thus the desired number is

$$S = (m+1)^{2d} \sum_{0 \leq |\mathbf{j}^1|_1 \leq N, 0 \leq |\mathbf{j}^2|_1 \leq N} 2^{\max\{j_1^1, j_1^2\} + \dots + \max\{j_d^1, j_d^2\}}. \quad (4.2)$$

Let $\mathbf{k}^1 = \max\{\mathbf{j}^1, \mathbf{j}^2\}$, where the maximum acts on each component of vectors. Similarly let $\mathbf{k}^2 = \min\{\mathbf{j}^1, \mathbf{j}^2\}$. Then $|\mathbf{k}^1 + \mathbf{k}^2|_1 = |\mathbf{j}^1 + \mathbf{j}^2|_1 = |\mathbf{j}^1|_1 + |\mathbf{j}^2|_1 \leq 2N$, and for each fixed $\mathbf{k}^1, \mathbf{k}^2$, there are at most 2^d pairs of $\mathbf{j}^1, \mathbf{j}^2$ satisfying the conditions $\mathbf{k}^1 = \max\{\mathbf{j}^1, \mathbf{j}^2\}$ and $\mathbf{k}^2 = \min\{\mathbf{j}^1, \mathbf{j}^2\}$. Thus

$$S \leq C(d)(m+1)^{2d} \sum_{0 \leq |\mathbf{k}^1|_1 + |\mathbf{k}^2|_1 \leq 2N} 2^{|\mathbf{k}^1|_1} \quad (4.3)$$

$$= C(d)(m+1)^{2d} \sum_{k=0}^{2N} 2^k \binom{k+d-1}{d-1} \sum_{l=0}^{2N-k} \binom{l+d-1}{d-1} \quad (4.4)$$

$$= C(d)(m+1)^{2d} N \sum_{k=0}^{2N} 2^k (k+1)^{d-1} (2N-k+1)^{d-1}. \quad (4.5)$$

The first equality is because there are $\binom{k+d}{d-1}$ choices of \mathbf{k}^1 with $|\mathbf{k}^1|_1 = k$, and similarly for \mathbf{k}^2 . The second equality is because $\binom{k+d-1}{d-1} = \frac{k+1}{1} \frac{k+2}{2} \dots \frac{k+d-1}{d-1} \leq (k+1)^{d-1}$, and taking the largest term in the l summation.

Then by taking derivative with respect of k , it is easy to see that the previous summation is optimized at $k_{max} = 2N - O(d)$. Thus

$$S \leq C(d)(m+1)^{2d} N^2 2^{k_{max}} (k_{max} + 1)^{d-1} (2N - k_{max} + 1)^{d-1} \quad (4.6)$$

$$\leq C(d)(m+1)^{2d} 2^{2N} N^{d+1}, \quad (4.7)$$

which finishes the proof. \square

Remark 4.2. When $d \geq 4$, one has $2^{2N} N^{2d-2} \geq 2^{2N} N^{d+1}$, thus in this case the number of $Q(\hat{f}_i, \hat{f}_j)$ needed to be computed is much less than the total number of pairs of \hat{f}_i, \hat{f}_j . And the bigger d is, the more saving one will gain.

Notice that if one is more careful when choosing the maximum to replace a summation (first on l and then on k), by using some kind of exponential decay of the summation with respect to the summation index, one should be able to get an estimate of order $O(2^{2N} N^{d-1})$.

For the case $m = 0$, all basis functions are piecewise constant. It is easy to see that if the collision kernel is deterministic, i.e. $b(\mathbf{z}) = 1$, then $\int_{I_{\mathbf{z}}} \psi_{\mathbf{j}^1, 1^1}(\mathbf{z}) \psi_{\mathbf{j}^2, 1^2}(\mathbf{z}) \psi_{\mathbf{j}^3, 1^3}(\mathbf{z}) \pi(\mathbf{z}) d\mathbf{z} = 0$ if and only if two of the functions are the same, and the third one has support properly including that of the previous two functions. In principle this should lead to a better sparsity result.

The sparsity of the tensor S_{ijk} can be estimated by the same method.

5 Regularity and Accuracy

We prove the regularity of the solution of the Boltzmann equation in the random space, and the accuracy of the stochastic Galerkin method with sparse grids. These are straightforward multi-dimensional extension of the corresponding results in [16]. We consider a d -dimensional random domain $I_{\mathbf{z}}$, $\mathbf{z} = (z_1, \dots, z_d)$. We also assume that the random collision kernel depends

linearly on \mathbf{z} . This is a reasonable assumption because when one uses the Karhunen-Loeve expansion to approximate a random field, the resulting dependence on \mathbf{z} is linear.

5.1 Regularity in the random space for the classical Boltzmann equation

In this subsection, we consider the spatially homogeneous classical Boltzmann equation

$$\frac{\partial f}{\partial t} = Q(f, f), \quad (5.1)$$

subject to random initial data and random collision kernel

$$f(0, \mathbf{v}, \mathbf{z}) = f^0(\mathbf{v}, \mathbf{z}), \quad B = B(\mathbf{v}, \mathbf{v}_*, \sigma, \mathbf{z}), \quad \mathbf{z} \in I_{\mathbf{z}}.$$

We define the norms and operators:

$$\|f(t, \cdot, \mathbf{z})\|_{L_{\mathbf{v}}^p} = \left(\int_{\mathbb{R}^d} |f(t, \mathbf{v}, \mathbf{z})|^p d\mathbf{v} \right)^{1/p}, \quad \|f(t, \mathbf{v}, \cdot)\|_{L_{\mathbf{z}}^2} = \left(\int_{I_{\mathbf{z}}} f(t, \mathbf{v}, \mathbf{z})^2 \pi(\mathbf{z}) d\mathbf{z} \right)^{1/2},$$

$$\|f(t, \cdot, \cdot)\|_k = \sup_{\mathbf{z} \in I_{\mathbf{z}}} \left(\sum_{|l|=0}^k \|\partial_{\mathbf{z}}^l f(t, \mathbf{v}, \mathbf{z})\|_{L_{\mathbf{v}}^2}^2 \right)^{1/2},$$

$$Q(g, h)(\mathbf{v}) = \int_{\mathbb{R}^d} \int_{S^{d-1}} B(\mathbf{v}, \mathbf{v}_*, \omega, \mathbf{z}) [g(\mathbf{v}')h(\mathbf{v}'_*) - g(\mathbf{v})h(\mathbf{v}_*)] d\omega d\mathbf{v}_*,$$

$$Q_{1,j}(g, h)(\mathbf{v}) = \int_{\mathbb{R}^d} \int_{S^{d-1}} \partial_{z_j} B(\mathbf{v}, \mathbf{v}_*, \omega, \mathbf{z}) [g(\mathbf{v}')h(\mathbf{v}'_*) - g(\mathbf{v})h(\mathbf{v}_*)] d\omega d\mathbf{v}_*.$$

We first state the following estimates of $Q(g, h)$ and $Q_{1,j}(g, h)$, which are standard results proved in [18, 7] and its extension to the uncertain case is straightforward:

Lemma 5.1. *Assume the collision kernel B depends on \mathbf{z} linearly, B and $\partial_{\mathbf{z}} B$ are locally integrable and bounded in \mathbf{z} . If $g, h \in L_{\mathbf{v}}^1 \cap L_{\mathbf{v}}^2$, then*

$$\|Q(g, h)\|_{L_{\mathbf{v}}^2}, \|Q_{1,j}(g, h)\|_{L_{\mathbf{v}}^2} \leq C_B \|g\|_{L_{\mathbf{v}}^1} \|h\|_{L_{\mathbf{v}}^2}, \quad (5.2)$$

$$\|Q(g, h)\|_{L_{\mathbf{v}}^2}, \|Q_{1,j}(g, h)\|_{L_{\mathbf{v}}^2} \leq C_B \|g\|_{L_{\mathbf{v}}^2} \|h\|_{L_{\mathbf{v}}^2}, \quad (5.3)$$

where the constant $C_B > 0$ depends only on B and $\partial_{z_j} B, j = 1, \dots, d$.

Now we state our estimate on $\|f\|_k$.

Theorem 5.2. *Assume that B satisfies the assumption in Lemma 5.1, and $\sup_{\mathbf{z} \in I_{\mathbf{z}}} \|f^0\|_{L_{\mathbf{v}}^1} \leq M$, $\|f^0\|_k < \infty$ for some integer $k \geq 0$. Then there exists a constant $C_k > 0$, depending only on C_B, M, T , and $\|f^0\|_k$ such that*

$$\|f\|_k \leq C_k, \quad \text{for any } t \in [0, T]. \quad (5.4)$$

The proof of the theorem is provided in the Appendix.

5.2 Accuracy analysis

In this subsection, we will prove the convergence rate of the stochastic Galerkin method using the previous established regularity. As in section 5.1, we will still restrict to the spatially homogeneous equation (5.1).

We use the sparse grid space $\hat{\mathbf{V}}_N^m$ with parameters m, N . For this space, the number of basis functions $K = O((m+1)^d 2^N N^{d-1})$.

Define the space $\mathcal{H}^m(I_{\mathbf{z}})$ by

$$\|f\|_{\mathcal{H}^m(I_{\mathbf{z}})} = \max \|\partial_{z_{i_1}}^m \cdots \partial_{z_{i_r}}^m\|_{L^2(I_{\mathbf{z}})},$$

where the maximum is taken over all subsets $\{i_1, \dots, i_r\} \subset \{1, \dots, d\}$. Using the orthonormal basis $\{\Phi_k(z)\}$, the solution f to (5.1) can be represented as

$$f(t, \mathbf{v}, \mathbf{z}) = \sum_{k=0}^{\infty} \hat{f}_k(t, \mathbf{v}) \Phi_k(\mathbf{z}), \quad \text{where} \quad \hat{f}_k(t, \mathbf{v}) = \int_{I_{\mathbf{z}}} f(t, \mathbf{v}, \mathbf{z}) \Phi_k(\mathbf{z}) \pi(\mathbf{z}) \, d\mathbf{z}. \quad (5.5)$$

Let P_K be the projection operator defined as

$$P_K f(t, \mathbf{v}, \mathbf{z}) = \sum_{k=0}^K \hat{f}_k(t, \mathbf{v}) \Phi_k(\mathbf{z}).$$

Then one has the following projection error estimate (Lemma 3.2 in Guo and Cheng [14] by taking $s = 0, p = q = k = m$):

Lemma 5.3. *For any $f \in \mathcal{H}^{m+1}(I_{\mathbf{z}})$, $N \geq 1$, we have*

$$\|P_K f - f\|_{L^2(I_{\mathbf{z}})} \leq (C(m)N)^d 2^{-N(m+1)} \|f\|_{\mathcal{H}^{m+1}(I_{\mathbf{z}})}. \quad (5.6)$$

This lemma implies that the projection error

$$\|P_K f - f\|_{L^2(I_{\mathbf{z}})} \leq C(m, d) K^{-(m+1)} (\log K)^{(m+2)(d-1)} \|f\|_{\mathcal{H}^{m+1}(I_{\mathbf{z}})}. \quad (5.7)$$

Define the norms

$$\|f(t, \mathbf{v}, \cdot)\|_{H_{\mathbf{z}}^k} = \left(\sum_{|\mathbf{l}|=0}^k \|\partial_{\mathbf{z}}^{\mathbf{l}} f(t, \mathbf{v}, \mathbf{z})\|_{L_{\mathbf{z}}^2}^2 \right)^{1/2}, \quad \|f(t, \cdot, \cdot)\|_{L_{\mathbf{v}, \mathbf{z}}^2} = \left(\int_{I_{\mathbf{z}}} \int_{\mathbb{R}^d} f(t, \mathbf{v}, \mathbf{z})^2 \, d\mathbf{v} \pi(\mathbf{z}) \, d\mathbf{z} \right)^{1/2}, \quad (5.8)$$

then we have the following:

Lemma 5.4. *Assume \mathbf{z} obeys the uniform distribution, i.e., $\mathbf{z} \in I_{\mathbf{z}} = [-1, 1]^d$ and $\pi(\mathbf{z}) = 1/2^d$. If $\|f^0\|_{d(m+1)}$ is bounded, then*

$$\|P_K f - f\|_{L_{\mathbf{v}, \mathbf{z}}^2} \leq C(m, d) K^{-(m+1)} (\log K)^{(m+2)(d-1)} \|f\|_{\mathcal{H}^{m+1}(I_{\mathbf{z}})}, \quad (5.9)$$

where $C(m, d)$ is a constant depending on m and d .

Give the gPC approximation of f :

$$f^K(t, \mathbf{v}, \mathbf{z}) = \sum_{k=0}^K f_k(t, \mathbf{x}, \mathbf{v}) \Phi_k(\mathbf{z}), \quad (5.10)$$

(a) $m = 0$				(b) $m = 1$			
	$N = 3$	$N = 4$	$N = 5$		$N = 3$	$N = 4$	$N = 5$
$d = 1$	8,8	16,16	32,32	$d = 1$	16,16	32,32	64,64
$d = 2$	20,64	48,256	112,1024	$d = 2$	80,256	192,1024	448,4096
$d = 3$	38,512	104,4096	272,32768	$d = 3$	304,4096	832,32768	2176,262144
$d = 4$	63,4096	192,65536	552,1048576				

Table 1: Comparison of number of basis functions: m is the maximal degree of polynomials. d is the dimension; in each cell, the left number is the number of basis of functions of $\hat{\mathbf{V}}_N^m$; the right number is the number of basis of functions of \mathbf{V}_N^m .

we now define the error function

$$e^K(t, \mathbf{v}, \mathbf{z}) = P_K f(t, \mathbf{v}, \mathbf{z}) - f^K(t, \mathbf{v}, \mathbf{z}) := \sum_{k=0}^K e_k(t, \mathbf{v}) \Phi_k(\mathbf{z}),$$

where $e_k = \hat{f}_k - f_k$. Then we have

Theorem 5.5. *Assume the random variable z and initial data f^0 satisfy the assumption in Lemma 5.4, and the gPC approximation f^K is uniformly bounded in K , then*

$$\|f - f^K\|_{L^2_{\mathbf{v},z}} \leq C(t) \left\{ C(m, d) K^{-(m+1)} (\log K)^{(m+2)(d-1)} + \|e^K(0)\|_{L^2_{\mathbf{v},z}} \right\}.$$

The proof of Lemma 5.4 and Theorem 5.5 can be proved in the same way as Section 4.2 in Hu and Jin [16], in view of Lemma 5.3. We omit the details.

6 Numerical results

In this section we give some numerical results of the stochastic Galerkin method with sparse grid technique. We first demonstrate the efficiency of the sparse grid basis, and then show its application to the Boltzmann equation with uncertainty.

6.1 The sparse grid basis

6.1.1 Number of basis functions

We first give a comparison of number of basis functions between our sparse grid function space $\hat{\mathbf{V}}_N^m$ and the tensor grid basis \mathbf{V}_N^m . The result is shown in Table 1. It is clear that the sparse grid technique saves a great number of basis functions, especially in high dimensional random spaces.

6.1.2 Efficiency of the sparse grid function space

We give a comparison of the L^2 approximation error of $\hat{\mathbf{V}}_N^m$ and \mathbf{V}_N^m . For each random dimension $d = 2, 3, 4$ we pick a smooth test function as follows:

$$f(\mathbf{z}) = \frac{1}{2\pi\mathcal{K}(\mathbf{z})^2} \exp\left(-\frac{1}{2\mathcal{K}(\mathbf{z})}\right) \left(2\mathcal{K}(\mathbf{z}) - 1 + \frac{1 - \mathcal{K}(\mathbf{z})}{2\mathcal{K}(\mathbf{z})}\right), \quad (6.1)$$




Figure 1: Comparison of approximation error of both sparse grid and full grid for $d = 2, 3, 4$. For $d = 4$ we do not give the result by tensor grid because the number of basis functions is too large.

where

$$\begin{aligned}
 \mathcal{K}_{d=2}(\mathbf{z}) &= 1 - 0.5(0.5 + 0.1 \sin(z_1) + 0.1 \sin(2z_2)), \\
 \mathcal{K}_{d=3}(\mathbf{z}) &= 1 - 0.5(0.5 + 0.1 \sin(z_1) + 0.1 \sin(2z_2) + 0.1 \cos(z_3)), \\
 \mathcal{K}_{d=4}(\mathbf{z}) &= 1 - 0.5(0.5 + 0.1 \sin(z_1) + 0.1 \sin(2z_2) + 0.1 \cos(z_3) + 0.1 \cos(2z_4)).
 \end{aligned} \tag{6.2}$$

We use the function spaces $\hat{\mathbf{V}}_N^m$ and \mathbf{V}_N^m with different m, N values to approximate these functions, and compute their relative L^2 error $\frac{\|f - P_K f\|_{L^2}}{\|f\|_{L^2}}$, where P_K is the projection operator onto the corresponding function space. The result is shown in Figure 1. It can be seen that the sparse grid method performs much better than the tensor grid method.




Figure 2: Sparsity of S_{ijk} and the number of $Q(\hat{f}_i, \hat{f}_j)$ needed to compute, $d = 2, 3, 4$, $m = 0$.

6.1.3 Sparsity of S_{ijk}

We give a test of the sparsity of the tensor S_{ijk} , as well as the number of $Q(\hat{f}_i, \hat{f}_j)$ needed to compute. We take a random collision kernel $b(\mathbf{z}) = 1 + 0.2z_1$. For simplicity we only show the results with $m = 0$, since the sparsity of S_{ijk} with larger m is similar. The result is shown in Figure 2. One can clearly see an exponential decay of the percentage of nonzeros in S_{ijk} , as well as the percentage of $Q(\hat{f}_i, \hat{f}_j)$ needed to compute, as N or d increase. This is even better than what we have proved.

To further demonstrate the sparsity of S_{ijk} we give a graph of nonzero elements of S_{ijk} for $m = 0$, $N = 4$, $d = 3$, shown in Figure 3. The points in the first graph represent nonzero elements in S_{ijk} . The second graph is the projection of the first graph onto i, j coordinates, and the points in it represent those $Q(\hat{f}_i, \hat{f}_j)$ needed to compute.

6.2 Application to the Boltzmann equation with uncertainty

In this subsection, the velocity space is assumed to be two-dimensional and its discretization is always given by $N_v = 32$. The time discretization is given by 0.8 times the CFL condition for spatial inhomogeneous problems.

6.2.1 Accuracy of the approximation of the collision operator

We first check the accuracy of the collision operator $Q(f, f)$ computed by the sparse grid stochastic Galerkin method. The function f is given by the Bobylev-Krook-Wu [6, 17] solution with uncertainty:

$$f(\mathbf{v}, \mathbf{z}) = \frac{1}{2\pi\mathcal{K}(\mathbf{z})^2} \exp\left(-\frac{|\mathbf{v}|^2}{2\mathcal{K}(\mathbf{z})}\right) \left(2\mathcal{K}(\mathbf{z}) - 1 + \frac{1 - \mathcal{K}(\mathbf{z})}{2\mathcal{K}(\mathbf{z})} \mathbf{v}^2\right), \quad (6.3)$$




Figure 3: Demonstration of sparsity of S_{ijk} : $m = 0, N = 4, d = 3$.

where

$$\begin{aligned}
 \mathcal{K}_{d=2}(\mathbf{z}) &= 1 - 0.5(0.5 + 0.1 \sin(z_1) + 0.1 \sin(2z_2)), \\
 \mathcal{K}_{d=3}(\mathbf{z}) &= 1 - 0.5(0.5 + 0.1 \sin(z_1) + 0.1 \sin(2z_2) + 0.1 \cos(z_3)), \\
 \mathcal{K}_{d=4}(\mathbf{z}) &= 1 - 0.5(0.5 + 0.1 \sin(z_1) + 0.1 \sin(2z_2) + 0.1 \cos(z_3) + 0.1 \cos(2z_4)).
 \end{aligned} \tag{6.4}$$

For this f , $Q(f, f)$ with collision kernel $B = \frac{1}{2\pi}$ is given explicitly by

$$\begin{aligned}
 Q(f, f)(\mathbf{v}, \mathbf{z}) &= \left(\left(-\frac{2}{\mathcal{K}(\mathbf{z})} + \frac{|\mathbf{v}|^2}{2\mathcal{K}(\mathbf{z})^2} \right) f \right. \\
 &\quad \left. + \frac{1}{2\pi\mathcal{K}(\mathbf{z})^2} \exp\left(-\frac{|\mathbf{v}|^2}{2\mathcal{K}(\mathbf{z})}\right) \left(2 - \frac{1}{2\mathcal{K}(\mathbf{z})^2} |\mathbf{v}|^2 \right) \right) \frac{1 - \mathcal{K}(\mathbf{z})}{8}.
 \end{aligned} \tag{6.5}$$

The numerical solution is given by

$$\tilde{Q}(f, f)(\mathbf{v}, \mathbf{z}) = \sum_{k=0}^K Q_k(\mathbf{v}) \Phi_k(\mathbf{z}), \quad \text{where } Q_k(\mathbf{v}) = \sum_{i,j=0}^K S_{ijk} Q(\hat{f}_i, \hat{f}_j)(\mathbf{v}).$$

We compare the relative L^2 error for $d = 2, 3, 4$ and sparse grid $\hat{\mathbf{V}}_N^m$ with different m, N . The result is shown in Figure 4. One can clearly see the error is a little worse than $O(K^{-(m+1)})$, and it becomes a little worse as d increases. This is caused by the $\log K$ factor in the error estimate.

6.2.2 The homogeneous Boltzmann equation with uncertainty on the collision kernel

We solve the homogeneous Boltzmann equation with deterministic initial data and a random collision kernel. We take the dimension of the random space $d = 2, 3$, and the collision kernels are

$$\begin{aligned}
 b(\mathbf{z}) &= 1 + 0.2z_1 + 0.1z_2, & d = 2, \\
 b(\mathbf{z}) &= 1 + 0.2z_1 + 0.1z_2 + 0.07z_3, & d = 3.
 \end{aligned} \tag{6.6}$$




Figure 4: Accuracy of the approximation of the collision operator for $d = 2, 3, 4$.

The initial data is the BKW solution

$$f_0(\mathbf{v}, \mathbf{z}) = \frac{1}{\pi} \exp(-|\mathbf{v}|^2) \frac{|\mathbf{v}|^2}{2}, \quad (6.7)$$

and the exact solution is given by

$$f(t, \mathbf{v}, \mathbf{z}) = \frac{1}{2\pi\mathcal{K}^2} \exp\left(-\frac{|\mathbf{v}|^2}{2\mathcal{K}}\right) \left(2\mathcal{K} - 1 + \frac{1 - \mathcal{K}}{2\mathcal{K}} |\mathbf{v}|^2\right), \quad (6.8)$$

with

$$\mathcal{K}(t, \mathbf{z}) = 1 - \exp(-b(\mathbf{z})t/8)/2. \quad (6.9)$$

We solve this equation by the sparse grid sG method with $m = 0$, time step $\Delta t = 0.01$ and final time $t = 1$, and check the relative L^2 error with the exact solution. The result is shown in Figure 5. The phenomenon is similar to the previous accuracy test.

6.2.3 The Boltzmann equation with random initial data

We test our method on the (inhomogeneous) Boltzmann equation with uncertainty. The random space is 4-dimensional. We take the x -domain to be $[0, 1]$ with the periodic boundary condition. We use the following random initial data to mimic the Karhunen-Loeve expansion

$$\begin{cases} \rho_0 = \frac{1}{3} (2 + \sin(2\pi x) + \sin(4\pi x)z_1/2 + \sin(6\pi x)z_2/4 + \sin(8\pi x)z_3/6 + \sin(10\pi x)z_4/7), \\ \mathbf{u}_0 = (0.2, 0), \\ T_0 = \frac{1}{4} (3 + \cos(2\pi x) + \cos(4\pi x)z_1/2 + \cos(6\pi x)z_2/4 + \cos(8\pi x)z_3/6 + \cos(10\pi x)z_4/7), \\ f = \frac{\rho_0}{4\pi T_0} \left(\exp\left(-\frac{|\mathbf{v} - \mathbf{u}_0|^2}{2T_0}\right) + \exp\left(-\frac{|\mathbf{v} + \mathbf{u}_0|^2}{2T_0}\right) \right). \end{cases} \quad (6.10)$$

The x -domain is discretized into $N_x = 50$ mesh points, and we compare the solution by the sparse grid stochastic Galerkin method with $m = 0$, $N = 3$ and a stochastic collocation method




Figure 5: The homogeneous Boltzmann equation with a random collision kernel: accuracy result. $m = 0$, $\Delta t = 0.01$, $t = 1$.

with full grid in random space at time $t = 0.1$. The collocation method is implemented by solving the deterministic problem at points of the form $\mathbf{z} = (z_1, \dots, z_d)$ where each z_i is one of the $M_z = 8$ Gauss-Legendre quadrature points (thus one needs to solve M_z^d deterministic problems). And then the mean and standard deviation are computed by numerical quadrature. The comparison result is shown in Figure 6. We see the results by the two methods agree well.

6.2.4 The Boltzmann equation with randomness on initial data, boundary data, and collision kernel

We finally solve the inhomogeneous Boltzmann equation with uncertainty on initial data, boundary data, and collision kernel. The random domain is taken to be 6-dimensional. We take the initial data to be the equilibrium with

$$\rho(x, \mathbf{z}) = 1, \quad \mathbf{u}(x, \mathbf{z}) = 0, \quad T = 1 + 0.5(1 + 0.2z_2) \exp(-100(1 + 0.1z_3)(x - 0.4 - 0.01z_1)^2), \quad (6.11)$$

and the boundary data is given by the Maxwellian boundary condition with random parameters

$$T_w = 1 + 0.2z_4, \quad \alpha = 0.5 + 0.3z_5. \quad (6.12)$$

The collision kernel is given by

$$b(\mathbf{z}) = 1 + 0.2z_6. \quad (6.13)$$

The spatial discretization is given by $N_x = 100$ to better capture the details near the boundary. We compare the result by the stochastic Galerkin method with sparse grid technique with the stochastic collocation method with full grid at time $t = 0.04$. The Galerkin method has parameters $m = 0$, $N = 3$, and the collocation method is as described in the previous numerical result with $M_z = 4$ collocation points in each dimension. The result is shown in Figure 7. One can see that the two results agree well.




Figure 6: The Boltzmann equation with random initial data. $N_x = 50$, $t = 0.1$. Curve: collocation with $M_z = 8$; asterisks: Galerkin with $m = 0, N = 3$. Left column: mean of density, first component of bulk velocity, and temperature. Right column: standard deviation of density, first component of bulk velocity, and temperature.




Figure 7: The Boltzmann equation with randomness on initial data, boundary data, and collision kernel ($d = 6$). $N_x = 100$, $t = 0.04$. Curve: collocation with $M_z = 4$; asterisks: Galerkin with $m = 0$, $N = 3$. Left column: mean of density, first component of bulk velocity, and temperature. Right column: standard deviation of density, first component of bulk velocity, and temperature.

7 Conclusion

In this paper we developed a sparse grid based stochastic Galerkin method for the Boltzmann equation with uncertainty. The uncertainty could come from initial data, boundary data, and collision kernel. This method enables us to quantify the uncertainty from truly high dimensional random inputs, which is previously infeasible using the global gPC basis. We proved and numerically demonstrated the sparsity of the basis related coefficient, S_{ijk} , which allows us to dramatically accelerate the computation of the collision operator under the Galerkin projection. Regularity of the solution of the Boltzmann equation in the random space and an accuracy result of the stochastic Galerkin method are proved.

Many related problems are still open, for example, asymptotic-preserving schemes [9] for the Boltzmann equation with uncertainty, adaptive mesh techniques to capture discontinuities in the random space, quantification of nonlinear uncertainties on the collision kernel, etc.

Appendix: Proof of Theorem 5.2

Proof. First, from the conservation property of Q , one has

$$\|f(t, \cdot, \mathbf{z})\|_{L_v^1} = \|f^0(\cdot, \mathbf{z})\|_{L_v^1} \leq M.$$

Then we use mathematical induction on k . For $k = 0$, multiplying (5.1) by f and integrating on \mathbf{v} , by the Cauchy-Schwarz inequality and (5.2), one obtains

$$\frac{1}{2} \partial_t \int_{\mathbb{R}^d} f^2 \, d\mathbf{v} = \int_{\mathbb{R}^d} f Q(f, f) \, d\mathbf{v} \leq \|f\|_{L_v^2} \|Q(f, f)\|_{L_v^2} \leq C_B \|f\|_{L_v^1} \|f\|_{L_v^2}^2 \leq C_B M \|f\|_{L_v^2}^2.$$

Now Gronwall's inequality implies that there is a positive constant C_0 such that (5.4) is true for $k = 0$.

Now for some $k \geq 0$ assume (5.4) holds. Take any multi-index \mathbf{j} with $|\mathbf{j}| = k + 1$. Taking \mathbf{j} -th derivative of z on (5.1) gives

$$\partial_t \partial_{\mathbf{z}}^{\mathbf{j}} f = \sum_{\mathbf{l}=0}^{\mathbf{j}} \binom{\mathbf{j}}{\mathbf{l}} Q(\partial_{\mathbf{z}}^{\mathbf{l}} f, \partial_{\mathbf{z}}^{\mathbf{j}-\mathbf{l}} f) + \sum_{m=1}^d j_m \sum_{\mathbf{l}=0}^{\mathbf{j}-\mathbf{1}_m} \binom{\mathbf{j}-\mathbf{1}_m}{\mathbf{l}} Q_{1,m}(\partial_{\mathbf{z}}^{\mathbf{l}} f, \partial_{\mathbf{z}}^{\mathbf{j}-\mathbf{1}_m-\mathbf{l}} f), \quad (\text{A.1})$$

where we used the bilinearity of the collision operator and the assumption that B is linear in z .

Multiplying (A.1) by $\partial_{\mathbf{z}}^{\mathbf{j}} f$ and integrating over \mathbf{v} yields

$$\begin{aligned} & \frac{1}{2} \partial_t \int_{\mathbb{R}^d} (\partial_{\mathbf{z}}^{\mathbf{j}} f)^2 \, d\mathbf{v} \\ & \leq \sum_{\mathbf{l}=0}^{\mathbf{j}} \binom{\mathbf{j}}{\mathbf{l}} \|\partial_{\mathbf{z}}^{\mathbf{l}} f\|_{L_v^2} \|Q(\partial_{\mathbf{z}}^{\mathbf{l}} f, \partial_{\mathbf{z}}^{\mathbf{j}-\mathbf{l}} f)\|_{L_v^2} + \sum_{m=1}^d j_m \sum_{\mathbf{l}=0}^{\mathbf{j}-\mathbf{1}_m} \binom{\mathbf{j}-\mathbf{1}_m}{\mathbf{l}} \|\partial_{\mathbf{z}}^{\mathbf{l}} f\|_{L_v^2} \|Q_{1,m}(\partial_{\mathbf{z}}^{\mathbf{l}} f, \partial_{\mathbf{z}}^{\mathbf{j}-\mathbf{1}_m-\mathbf{l}} f)\|_{L_v^2} \\ & \leq \sum_{\mathbf{l}=0}^{\mathbf{j}} \binom{\mathbf{j}}{\mathbf{l}} C_B \|\partial_{\mathbf{z}}^{\mathbf{l}} f\|_{L_v^2} \|\partial_{\mathbf{z}}^{\mathbf{l}} f\|_{L_v^2} \|\partial_{\mathbf{z}}^{\mathbf{j}-\mathbf{l}} f\|_{L_v^2} + \sum_{m=1}^d j_m \sum_{\mathbf{l}=0}^{\mathbf{j}-\mathbf{1}_m} \binom{\mathbf{j}-\mathbf{1}_m}{\mathbf{l}} C_B \|\partial_{\mathbf{z}}^{\mathbf{l}} f\|_{L_v^2} \|\partial_{\mathbf{z}}^{\mathbf{l}} f\|_{L_v^2} \|\partial_{\mathbf{z}}^{\mathbf{j}-\mathbf{1}_m-\mathbf{l}} f\|_{L_v^2} \\ & \leq C_B C_k^2 \|\partial_{\mathbf{z}}^{\mathbf{j}} f\|_{L_v^2} \sum_{\mathbf{0} \leq \mathbf{l} \leq \mathbf{j}, \mathbf{l} \neq \mathbf{0}, \mathbf{j}} \binom{\mathbf{j}}{\mathbf{l}} + 2C_B C_0 \|\partial_{\mathbf{z}}^{\mathbf{j}} f\|_{L_v^2}^2 + C_B C_k^2 \|\partial_{\mathbf{z}}^{\mathbf{j}} f\|_{L_v^2} \sum_{m=1}^d j_m \sum_{\mathbf{l}=0}^{\mathbf{j}-\mathbf{1}_m} \binom{\mathbf{j}-\mathbf{1}_m}{\mathbf{l}} \\ & = (2^{k+1} - 2) C_B C_k^2 \|\partial_{\mathbf{z}}^{\mathbf{j}} f\|_{L_v^2} + 2C_B C_0 \|\partial_{\mathbf{z}}^{\mathbf{j}} f\|_{L_v^2}^2 + 2^k (k+1) C_B C_k^2 \|\partial_{\mathbf{z}}^{\mathbf{j}} f\|_{L_v^2}. \end{aligned} \quad (\text{A.2})$$

In the first inequality we used the Cauchy-Schwarz inequality, and in the second inequality we used (5.3). In the third inequality the induction assumption is used for the second sum, since the indexes \mathbf{l} and $\mathbf{j} - \mathbf{1}_m - \mathbf{l}$ appeared there have order less than or equal to k . Every term in the first sum can be treated similarly except terms corresponding to the cases of $\mathbf{l} = \mathbf{0}$ and $\mathbf{l} = \mathbf{j}$, which are treated separately. In the final equality, we used the identity $\sum_{l=0}^L \binom{L}{l} = (1+1)^L = 2^L$.

Then we apply Gronwall's inequality to (A.2) and get the control

$$\sup_{\mathbf{z} \in I_{\mathbf{z}}} \left(\|\partial_{\mathbf{z}}^{\mathbf{j}} f(t, \mathbf{v}, \mathbf{z})\|_{L_v^2}^2 \right)^{1/2} \leq C_{k+1},$$

with a positive constant C_{k+1} . Sum over all \mathbf{j} with $|\mathbf{j}| = k + 1$ we get (5.4) for $k + 1$. This completes the mathematical induction and the proof. \square

References

- [1] B. Alpert. A class of bases in L^2 for the sparse representation of integral operators. *SIAM Journal on Mathematical Analysis*, 24(1):246–262, 1993.
- [2] I. Babuska, F. Nobile, and R. Tempone. A stochastic collocation method for elliptic partial differential equations with random input data. *SIAM Journal on Numerical Analysis*, 45(3):1005–1034, 2007.
- [3] I. Babuska, R. Tempone, and G. E. Zouraris. Galerkin finite element approximations of stochastic elliptic partial differential equations. *SIAM Journal on Numerical Analysis*, 42(2):800–825, 2004.
- [4] J. Back, F. Nobile, L. Tamellini, and R. Tempone. Stochastic spectral galerkin and collocation methods for pdes with random coefficients: a numerical comparison. In E. M. Rønquist J. S. Hesthaven, editor, *Spectral and High Order Methods for Partial Differential Equations*. Springer-Verlag Berlin Heidelberg, 2011.
- [5] G. A. Bird. *Molecular Gas Dynamics and the Direct Simulation of Gas Flows*. Clarendon Press, Oxford, 1994.
- [6] A. V. Bobylev. One class of invariant solutions of the Boltzmann equation. *Akademiia Nauk SSSR, Doklady*, 231:571–574, 1976.
- [7] F. Bouchut and L. Desvillettes. A proof of the smoothing properties of the positive part of Boltzmann’s kernel. *Revista Matemática Iberoamericana*, 14:47–61, 1998.
- [8] C. Cercignani. *The Boltzmann Equation and Its Applications*. Springer-Verlag, New York, 1988.
- [9] F. Filbet and S. Jin. A class of asymptotic-preserving schemes for kinetic equations and related problems with stiff sources. *Journal of Computational Physics*, 229:7625–7648, 2010.
- [10] J. Garcke and M. Griebel. *Sparse Grids and Applications*. Springer, 2013.
- [11] R. G. Ghanem and P. D. Spanos. *Stochastic Finite Elements: A Spectral Approach*. Springer-Verlag, New York, 1991.
- [12] M. Griebel. Adaptive sparse grid multilevel methods for elliptic PDEs based on finite differences. *Computing*, 61(2):151–179, 1998.
- [13] M. Griebel and G. Zumbusch. Adaptive sparse grids for hyperbolic conservation laws. In *Hyperbolic Problems: Theory, Numerics, Applications*, pages 411–422. Springer, 1999.
- [14] W. Guo and Y. Cheng. A sparse grid discontinuous Galerkin method for high-dimensional transport equations and its application to kinetic simulations. *SIAM Journal on Scientific Computing*, accepted.
- [15] Bungartz H.-J. and M. Griebel. Sparse grids. *Acta Numerica*, 13:147–269, 2004.

- [16] J. Hu and S. Jin. A stochastic Galerkin method for the Boltzmann equation with uncertainty. *Journal of Computational Physics*, 315:150–168, 2016.
- [17] M. Krook and T. T. Wu. Formation of Maxwellian tails. *Physics of Fluids*, 20:1589–1595, 1977.
- [18] P.-L. Lions. Compactness in Boltzmann’s equation via Fourier integral operators and applications. I, II. *Journal of Mathematics of Kyoto University*, 34(2):391–427,429–461, 1994.
- [19] O. P. Le Maître and O. M. Knio. *Spectral Methods for Uncertainty Quantification, Scientific Computation, with Applications to Computational Fluid Dynamics*. Springer, New York, 2010.
- [20] H. Niederreiter, P. Hellekalek, G. Larcher, and P. Zinterhof. *Monte Carlo and Quasi-Monte Carlo Methods 1996*. Springer-Verlag, 1998.
- [21] F. Nobile, R. Tempone, and C. Webster. A sparse grid stochastic collocation method for partial differential equations with random input data. *SIAM Journal on Numerical Analysis*, 46(5):2309–2345, 2008.
- [22] S. Smolyak. Quadrature and interpolation formulas for tensor products of certain classes of functions. *Doklady Akademii Nauk SSSR*, 4:240–243, 1963.
- [23] Z. Wang, Q. Tang, W. Guo, and Y. Cheng. Sparse grid discontinuous Galerkin methods for high-dimensional elliptic equations. *Journal of Computational Physics*, accepted.
- [24] D. Xiu. Fast numerical methods for stochastic computations: a review. *Communications in Computational Physics*, 5(2-4):242–272, 2009.
- [25] D. Xiu. *Numerical Methods for Stochastic Computation*. Princeton University Press, Princeton, New Jersey, 2010.
- [26] D. Xiu and J. Hesthaven. High-order collocation methods for differential equations with random inputs. *SIAM Journal on Scientific Computing*, 27(3):1118–1139, 2005.
- [27] C. Zenger. Sparse grids. In *Parallel Algorithms for Partial Differential Equations, Proceedings of the Sixth GAMM-Seminar*, volume 31. 1990.