

Introduction to Shimura Varieties with Bad Reduction of Parahoric Type

Thomas J. Haines

CONTENTS

1. Introduction	583
2. Notation	584
3. Iwahori and parahoric subgroups	587
4. Local models	589
5. Some PEL Shimura varieties with parahoric level structure at p	593
6. Relating Shimura varieties and their local models	602
7. Flatness	608
8. The Kottwitz-Rapoport stratification	609
9. Langlands' strategy for computing local L -factors	614
10. Nearby cycles	618
11. The semi-simple local zeta function for “fake” unitary Shimura varieties	621
12. The Newton stratification on Shimura varieties over finite fields	628
13. The number of irreducible components in $Sh_{\overline{\mathbb{F}}_p}$	634
14. Appendix: Summary of Dieudonné theory and de Rham and crystalline cohomology for abelian varieties	636
References	640

1. Introduction

This survey article is intended to introduce the reader to several important concepts relating to Shimura varieties with parahoric level structure at p . The main tool is the Rapoport-Zink local model **[RZ]**, which plays an important role in several aspects of the theory. We discuss local models attached to general linear and symplectic groups, and we illustrate their relation to Shimura varieties in two examples: the *simple* or “fake” unitary Shimura varieties with parahoric level structure, and the Siegel modular varieties with $\Gamma_0(p)$ -level structure. In addition, we present some applications of local models to questions of flatness, stratifications

Research partially supported by NSF grant DMS 0303605.

of the special fiber, and the determination of the semi-simple local zeta functions for simple Shimura varieties.

There are several good references for material of this sort that already exist in the literature. This survey has a great deal of overlap with two articles of Rapoport: [R1] and [R2]. A main goal of this paper is simply to make more explicit some of the ideas expressed very abstractly in those papers. Hopefully it will shed some new light on the earlier seminal works of Rapoport-Zink [RZ82], and Zink [Z]. This article is also closely related to some recent work of H. Reimann [Re1], [Re2], [Re3].

Good general introductions to aspects of the Langlands program which might be consulted while reading parts of this report are those of Blasius-Rogawski [BR], and T. Wedhorn [W2].

Several very important developments have taken place in the theory of Shimura varieties with bad reduction, which are completely ignored in this report. In particular, we mention the book of Harris-Taylor [HT] which uses bad reduction of Shimura varieties to prove the local Langlands conjecture for $GL_n(\mathbb{Q}_p)$, and the recent work of L. Fargues and E. Mantovan [FM].

Most of the results stated here are well-known by now (although scattered around the literature, with differing systems of conventions). However, the author took this opportunity to present a few new results (and some new proofs of old results). For example, there is the proof of the non-emptiness of the Kottwitz-Rapoport strata in any connected component of the Siegel modular and “fake” unitary Shimura varieties with Iwahori-level structure (Lemmas 13.1, 13.2), some foundational relations between Newton strata, Kottwitz-Rapoport strata, and affine Deligne-Lusztig varieties (Prop. 12.6), and the verification of the conjectural non-emptiness of the basic locus in the “fake” unitary case (Cor. 12.12). The main new proofs relate to topological flatness of local models attached to Iwahori subgroups of unramified groups (see §7) and to the description of the nonsingular locus of Shimura varieties with Iwahori-level structure (see §8.4). Finally, some of the results explained here (especially in §11) are background material necessary for the author’s as yet unpublished joint work with B.C. Ngô [HN3].

I am grateful to U. Görtz, R. Kottwitz, B.C. Ngô, G. Pappas, and M. Rapoport for all they have taught me about this subject over the years. I thank them for their various comments and suggestions on an early version of this article. Also, I heartily thank U. Görtz for providing the figures. Finally, I thank the Clay Mathematics Institute for sponsoring the June 2003 *Summer School on Harmonic Analysis, the Trace Formula, and Shimura Varieties*, which provided the opportunity for me to write this survey article.

2. Notation

2.1. Some field-theoretic notation. Fix a rational prime p . We let F denote a non-Archimedean local field of residual characteristic p , with ring of integers \mathcal{O} . Let $\mathfrak{p} \subset \mathcal{O}$ denote the maximal ideal, and fix a uniformizer $\pi \in \mathfrak{p}$. The residue field \mathcal{O}/\mathfrak{p} has cardinality q , a power of p .

We will fix an algebraic closure k of the finite field \mathbb{F}_p . The Galois group $\text{Gal}(k/\mathbb{F}_p)$ has a canonical generator (the Frobenius automorphism), given by $\sigma(x) = x^p$. For each positive integer r , we denote by k_r the fixed field of σ^r . Let $W(k_r)$ (resp. $W(k)$) denote the ring of Witt vectors of k_r (resp. k), with fraction field L_r

(resp. L). We also use the symbol σ to denote the Frobenius automorphism of L induced by that on k .

We fix throughout a rational prime $\ell \neq p$, and a choice of algebraic closure $\mathbb{Q}_\ell \subset \overline{\mathbb{Q}_\ell}$.

2.2. Some group-theoretic notation. The symbol \mathbf{G} will always denote a connected reductive group over \mathbb{Q} (sometimes defined over \mathbb{Z}). Unless otherwise indicated, G will denote the base-change \mathbf{G}_F , where F is a suitable local field (usually, $G = \mathbf{G}_{\mathbb{Q}_p}$).

Now let G denote a connected reductive F -group. Fix once and for all a Borel subgroup B and a maximal torus T contained in B . We will usually assume G is split over F , in which case we can even assume G, B and T are defined and split over the ring \mathcal{O} . For GL_n or GSp_{2n} , T will denote the usual “diagonal” torus, and B will denote the “upper triangular” Borel subgroup.

We will often refer to “standard” parahoric and “standard” Iwahori subgroups. For the group $G = \mathrm{GL}_n$ (resp. $G = \mathrm{GSp}_{2n}$), the “standard” hyperspecial maximal compact subgroup of $G(F)$ will be the subgroup $G(\mathcal{O})$. The “standard” Iwahori subgroup will be inverse image of $B(\mathcal{O}/\mathfrak{p})$ under the reduction modulo \mathfrak{p} homomorphism

$$G(\mathcal{O}) \rightarrow G(\mathcal{O}/\mathfrak{p}).$$

A “standard” parahoric will be defined similarly as the inverse image of a standard (= upper triangular) parabolic subgroup.

For $\mathrm{GL}_n(F)$, the standard Iwahori is the subgroup stabilizing the standard lattice chain (defined in §3). The standard parahorics are stabilizers of standard partial lattice chains. Similar remarks apply to the group $\mathrm{GSp}_{2n}(F)$. The symbols I or I_r or K_p^a will always denote a standard Iwahori subgroup of $G(F)$ defined in terms of our fixed choices of $B \supset T$, and $G(\mathcal{O})$ as above (for some local field F). Often (but not always) K or K_r or K_p^0 will denote our fixed hyperspecial maximal compact subgroup $G(\mathcal{O})$.

We have the associated spherical Hecke algebra $\mathcal{H}_K := C_c(K \backslash G(F) / K)$, a convolution algebra of \mathbb{C} -valued (or $\overline{\mathbb{Q}_\ell}$ -valued) functions on $G(F)$ where convolution is defined using the measure giving K volume 1. Similarly, $\mathcal{H}_I := C_c(I \backslash G(F) / I)$ is a convolution algebra using the measure giving I volume 1. For a compact open subset $U \subset G(F)$, $\mathbb{1}_U$ denotes the characteristic function of the set U .

The *extended affine Weyl group* of $G(F)$ is defined as the group $\widetilde{W} = N_{G(F)}T/T(\mathcal{O})$. The map $X_*(T) \rightarrow T(F)/T(\mathcal{O})$ given by $\lambda \mapsto \lambda(\pi)$ is an isomorphism of abelian groups. The finite Weyl group $W_0 := N_{G(F)}T/T(F)$ can be identified with $N_{G(\mathcal{O})}T/T(\mathcal{O})$, by choosing representatives of $N_{G(F)}T/T(F)$ in $G(\mathcal{O})$. Thus we have a canonical isomorphism

$$\widetilde{W} = X_*(T) \rtimes W_0.$$

We will denote elements in this group typically by the notation $t_\nu w$ (for $\nu \in X_*(T)$ and $w \in W_0$).

Our choice of $B \supset T$ determines a unique *opposite* Borel subgroup \bar{B} such that $B \cap \bar{B} = T$. We have a notion of B -positive (resp. \bar{B} -positive) root α and coroot α^\vee . Also, the group W_0 is a Coxeter group generated by the simple reflections s_α in the vector space $X_*(T) \otimes \mathbb{R}$ through the walls fixed by the B -positive (or \bar{B} -positive) simple roots α . Let w_0 denote the unique element of W_0 having greatest length with respect to this Coxeter system.

We will often need to consider \widetilde{W} as a subset of $G(F)$. We choose the following conventions. For each $w \in W_0$, we fix once and for all a lift in the group $N_{G(\mathcal{O})}T$. We identify each $\nu \in X_*(T)$ with the element $\nu(\pi) \in T(F) \subset G(F)$.

Let \mathbf{a} denote the alcove in the building of $G(F)$ which is fixed by the Iwahori I , or equivalently, the unique alcove in the apartment corresponding to T which is contained in the \bar{B} -positive (i.e., the B -negative) Weyl chamber, whose closure contains the origin (the vertex fixed by the maximal compact subgroup $G(\mathcal{O})$)¹.

The group \widetilde{W} permutes the set of affine roots $\alpha + k$ (α a root, $k \in \mathbb{Z}$) (viewed as affine linear functions on $X_*(T) \otimes \mathbb{R}$), and hence permutes (transitively) the set of alcoves. Let Ω denote the subgroup which stabilizes the base alcove \mathbf{a} . Then we have a semi-direct product

$$\widetilde{W} = W_{\text{aff}} \rtimes \Omega,$$

where W_{aff} (the *affine Weyl group*) is the Coxeter group generated by the reflections S_{aff} through the walls of \mathbf{a} . In the case where G is an almost simple group of rank l , with simple B -positive roots $\alpha_1, \dots, \alpha_l$, then S_{aff} consists of the l simple reflections $s_i = s_{\alpha_i}$ generating W_0 , along with one more simple affine reflection $s_0 = t_{-\tilde{\alpha}} s_{\tilde{\alpha}}$, where $\tilde{\alpha}$ is the highest B -positive root.

The Coxeter system $(W_{\text{aff}}, S_{\text{aff}})$ determines a length function ℓ and a Bruhat order \leq on W_{aff} , which extend naturally to \widetilde{W} : for $x_i \in W_{\text{aff}}$ and $\sigma_i \in \Omega$ ($i = 1, 2$), we define $x_1\sigma_1 \leq x_2\sigma_2$ in \widetilde{W} if and only if $\sigma_1 = \sigma_2$ and $x_1 \leq x_2$ in W_{aff} . Similarly, we set $\ell(x_1\sigma_1) = \ell(x_1)$.

By the Bruhat-Tits decomposition, the inclusion $\widetilde{W} \hookrightarrow G(F)$ induces a bijection

$$\widetilde{W} = I \backslash G(F) / I.$$

In the function-field case (e.g., $F = \mathbb{F}_p((t))$), the affine flag variety $\mathcal{Fl} = G(F)/I$ is naturally an ind-scheme, and the closures of the I -orbits $\mathcal{Fl}_w := IwI/I$ are determined by the Bruhat order on \widetilde{W} :

$$\mathcal{Fl}_x \subset \overline{\mathcal{Fl}_y} \iff x \leq y.$$

Similar statements hold for the affine Grassmannian, $\text{Grass} = G(F)/G(\mathcal{O})$. Now the $G(\mathcal{O})$ -orbits $\mathcal{Q}_\lambda := G(\mathcal{O})\lambda G(\mathcal{O})/G(\mathcal{O})$ are given (using the Cartan decomposition) by the B -dominant coweights $X_+(T)$:

$$X_+(T) \leftrightarrow G(\mathcal{O}) \backslash G(F) / G(\mathcal{O}).$$

By definition, λ is B -dominant if $\langle \alpha, \lambda \rangle \geq 0$ for all B -positive roots α . Here $\langle \cdot, \cdot \rangle : X^*(T) \times X_*(T) \rightarrow \mathbb{Z}$ is the canonical duality pairing.

The closure relations in Grass are given by the partial order on B -dominant coweights λ and μ :

$$\mathcal{Q}_\lambda \subset \overline{\mathcal{Q}_\mu} \iff \lambda \preceq \mu,$$

where $\lambda \prec \mu$ means that $\mu - \lambda$ is a sum of B -positive coroots.

Unless otherwise stated, a *dominant* coweight $\lambda \in X_*(T)$ will always mean one that is B -dominant.

For the group $G = \text{GL}_n$ or GSp_{2n} , there is a \mathbb{Z}_p -ind-scheme M which is a deformation of the affine Grassmannian $\text{Grass}_{\mathbb{Q}_p}$ to the affine flag variety $\mathcal{Fl}_{\mathbb{F}_p}$ for the underlying p -adic group G (see [HN1], and Remark 4.1). A very similar

¹This choice of base alcove results from our convention of embedding $X_*(T) \hookrightarrow G(F)$ by the rule $\lambda \mapsto \lambda(\pi)$; to see this, consider how vectors spanning the standard periodic lattice chain in §3 are identified with vectors in $X_*(T) \otimes \mathbb{R}$.

deformation Fl_X over a smooth curve X (due to Beilinson) exists for any group G in the function field setting, and has been extensively studied by Gaitsgory [Ga]. For any dominant coweight $\lambda \in X_+(T)$, the symbol M_λ will always denote the \mathbb{Z}_p -scheme which is the scheme-theoretic closure in M of $\mathcal{Q}_\lambda \subset \mathrm{Grass}_{\mathbb{Q}_p}$.

2.3. Duality notation. If A is any abelian scheme over a scheme S , we denote the dual abelian scheme by \hat{A} . The existence of \hat{A} over an arbitrary base is a delicate matter; see [CF], §I.1.

If M is a module over a ring R , we denote the dual module by $M^\vee = \mathrm{Hom}_R(M, R)$. Similar notation applies to quasi-coherent \mathcal{O}_S -modules over a scheme S .

If G is a connected reductive group over a local field F , then the Langlands dual (over \mathbb{C} or $\overline{\mathbb{Q}_\ell}$) will be denoted \hat{G} . The Langlands L -group is the semi-direct product ${}^L G = \hat{G} \rtimes W_F$, where W_F is the Weil-group of F .

2.4. Miscellaneous notation. We will use the following abbreviation for elements of R^n (here R can be any set): let a_1, \dots, a_r be a sequence of positive integers whose sum is n . A vector of the form $(x_1, \dots, x_1, x_2, \dots, x_2, \dots, x_r, \dots, x_r)$, where for $i = 1, \dots, r$, the element x_i is repeated a_i times, will be denoted by

$$(x_1^{a_1}, x_2^{a_2}, \dots, x_r^{a_r}).$$

We will denote by \mathbb{A} the adèles of \mathbb{Q} , by \mathbb{A}_f the finite adèles, and by \mathbb{A}_f^p the finite adèles away from p (with the exception of two instances, where \mathbb{A} denotes affine space!).

3. Iwahori and parahoric subgroups

3.1. Stabilizers of periodic lattice chains. We discuss the definitions for the groups GL_n and GSp_{2n} .

3.1.1. *The linear case.* For each $i \in \{1, \dots, n\}$, let e_i denote the i -th standard vector $(0^{i-1}, 1, 0^{n-i})$ in F^n , and let $\Lambda_i \subset F^n$ denote the free \mathcal{O} -module with basis $\pi^{-1}e_1, \dots, \pi^{-1}e_i, e_{i+1}, \dots, e_n$. We consider the diagram

$$\Lambda_\bullet : \Lambda_0 \longrightarrow \Lambda_1 \longrightarrow \dots \longrightarrow \Lambda_{n-1} \longrightarrow \pi^{-1}\Lambda_0,$$

where the morphisms are inclusions. The lattice chains $\pi^n \Lambda_\bullet$ ($n \in \mathbb{Z}$) fit together to form an infinite complete lattice chain Λ_i , ($i \in \mathbb{Z}$). If we identify each Λ_i with \mathcal{O}^n , then the diagram above becomes

$$\mathcal{O}^n \xrightarrow{m_1} \mathcal{O}^n \xrightarrow{m_2} \dots \xrightarrow{m_{n-1}} \mathcal{O}^n \xrightarrow{m_n} \mathcal{O}^n,$$

where m_i is the morphism given by the diagonal matrix

$$m_i = \mathrm{diag}(1, \dots, \pi, \dots, 1),$$

the element π appearing in the i th place. We define the (*standard*) *Iwahori subgroup* I of $\mathrm{GL}_n(F)$ by

$$I = \bigcap_i \mathrm{Stab}_{\mathrm{GL}_n(F)}(\Lambda_i).$$

Similarly, for any non-empty subset $J \subset \{0, 1, \dots, n-1\}$, we define the *parahoric subgroup* of $\mathrm{GL}_n(F)$ corresponding to the subset J by

$$\mathcal{P}_J = \bigcap_{i \in J} \mathrm{Stab}_{\mathrm{GL}_n(F)}(\Lambda_i).$$

Note that \mathcal{P}_J is a compact open subgroup of $\mathrm{GL}_n(F)$, and that $\mathcal{P}_{\{0\}} = \mathrm{GL}_n(\mathcal{O})$ is a hyperspecial maximal compact subgroup, in the terminology of Bruhat-Tits, cf. [T].

3.1.2. *The symplectic case.* The definitions for the group of symplectic similitudes GSp_{2n} are similar. We define this group using the alternating matrix

$$\tilde{I} = \begin{bmatrix} 0 & \tilde{I}_n \\ -\tilde{I}_n & 0 \end{bmatrix},$$

where \tilde{I}_n denotes the $n \times n$ matrix with 1 along the anti-diagonal and 0 elsewhere. Let $(x, y) := x^t \tilde{I} y$ denote the corresponding alternating pairing on F^{2n} . For an \mathcal{O} -lattice $\Lambda \subset F^{2n}$, we define $\Lambda^\perp := \{x \in F^{2n} \mid (x, y) \in \mathcal{O} \text{ for all } y \in \Lambda\}$. The lattice Λ_0 is self-dual (i.e., $\Lambda_0^\perp = \Lambda_0$). Consider the infinite lattice chain in F^{2n}

$$\cdots \longrightarrow \Lambda_{-2n} = \pi \Lambda_0 \longrightarrow \cdots \longrightarrow \Lambda_{-1} \longrightarrow \Lambda_0 \longrightarrow \cdots \longrightarrow \Lambda_{2n} = \pi^{-1} \Lambda_0 \longrightarrow \cdots$$

We have $\Lambda_i^\perp = \Lambda_{-i}$ for all $i \in \mathbb{Z}$. Now we define the (*standard*) Iwahori subgroup I of $\mathrm{GSp}_{2n}(F)$ by

$$I = \bigcap_i \mathrm{Stab}_{\mathrm{GSp}_{2n}(F)}(\Lambda_i).$$

For any non-empty subset $J \subset \{-n, \dots, -1, 0, 1, \dots, n\}$ such that $i \in J \Leftrightarrow -i \in J$, we define the *parahoric subgroup* corresponding to J by

$$\mathcal{P}_J = \bigcap_{i \in J} \mathrm{Stab}_{\mathrm{GSp}_{2n}(F)}(\Lambda_i).$$

3.2. Bruhat-Tits group schemes. In Bruhat-Tits theory, parahoric groups are defined as the groups $\mathfrak{G}_{\Delta_J}^0(\mathcal{O})$, where $\mathfrak{G}_{\Delta_J}^0$ is the neutral component of a group scheme \mathfrak{G}_{Δ_J} , defined and smooth over \mathcal{O} , which has generic fiber the F -group G , and whose \mathcal{O} -points are the subgroup of $G(F)$ fixing the facet Δ_J of the Bruhat-Tits building corresponding to the set J ; see [BT2], p. 356. By [T], 3.4.1 (see also [BT2], 1.7) we can characterize the group scheme \mathfrak{G}_{Δ_J} as follows: it is the unique \mathcal{O} -group scheme \mathcal{P} satisfying the following three properties:

- (1) \mathcal{P} is smooth and affine over \mathcal{O} ;
- (2) The generic fiber \mathcal{P}_F is G_F ;
- (3) For any unramified extension F' of F , letting $\mathcal{O}_{F'} \subset F'$ denote ring of integers, the group $\mathcal{P}(\mathcal{O}_{F'}) \subset G(F')$ is the subgroup of elements which fix the facet Δ_J in the Bruhat-Tits building of $G_{F'}$.

Let us show how automorphism groups of periodic lattice chains Λ_\bullet give a concrete realization of the Bruhat-Tits parahoric group schemes, in the GL_n and GSp_{2n} cases. We will discuss the Iwahori subgroups of GL_n in some detail, leaving for the reader the obvious generalizations to parahoric subgroups of GL_n (and GSp_{2n}).

For any \mathcal{O} -algebra R , we may consider the diagram $\Lambda_{\bullet,R} = \Lambda_\bullet \otimes_{\mathcal{O}} R$, and we may define the \mathcal{O} -group scheme Aut whose R -points are the isomorphisms of the diagram $\Lambda_{\bullet,R}$. More precisely, an element of $\mathrm{Aut}(R)$ is an n -tuple of R -linear automorphisms

$$(g_0, \dots, g_{n-1}) \in \mathrm{Aut}(\Lambda_{0,R}) \times \cdots \times \mathrm{Aut}(\Lambda_{n-1,R})$$

such that the following diagram commutes

$$\begin{array}{ccccccc}
 \Lambda_{0,R} & \longrightarrow & \cdots & \longrightarrow & \Lambda_{n-1,R} & \longrightarrow & \Lambda_{n,R} \\
 \downarrow g_0 & & & & \downarrow g_{n-1} & & \downarrow g_0 \\
 \Lambda_{0,R} & \longrightarrow & \cdots & \longrightarrow & \Lambda_{n-1,R} & \longrightarrow & \Lambda_{n,R}.
 \end{array}$$

The group functor Aut is obviously represented by an affine group scheme, also denoted Aut . Further, it is not hard to see that Aut is formally smooth, hence smooth, over \mathcal{O} . To show this, one has to check the lifting criterion for formal smoothness: if R is an \mathcal{O} -algebra containing a nilpotent ideal $\mathcal{J} \subset R$, then any automorphism of $\Lambda_{\bullet} \otimes_{\mathcal{O}} R/\mathcal{J}$ can be lifted to an automorphism of $\Lambda_{\bullet} \otimes_{\mathcal{O}} R$. This is proved on page 135 of [RZ]. Thus Aut satisfies condition (1) above.

Alcoves in the Bruhat-Tits building for GL_n over F can be described as complete periodic \mathcal{O} -lattice chains in F^n

$$\cdots \longrightarrow \mathcal{L}_0 \longrightarrow \mathcal{L}_1 \longrightarrow \cdots \longrightarrow \mathcal{L}_{n-1} \longrightarrow \mathcal{L}_n = \pi^{-1}\mathcal{L}_0 \longrightarrow \cdots$$

where the arrows are inclusions. We can regard Λ_{\bullet} as the base alcove in this building. It is clear that since π is invertible over the generic fiber F , the automorphism g_0 determines the other g_i 's over F and so $\text{Aut}_F = \text{GL}_n$. By construction we have $\text{Aut}(\mathcal{O}) = \text{Stab}_{\text{GL}_n(F)}(\Lambda_{\bullet})$. This is unchanged if we replace F by an unramified extension F' . It follows that Aut satisfies conditions (2) and (3) above. Thus, by uniqueness, $\text{Aut} = \mathfrak{G}_{\Delta_J}$ for $J = \{0, \dots, n-1\}$.

Further, one can check that the special fiber Aut_k is an extension of the Borel subgroup B_k by a connected unipotent group over k ; hence the special fiber is connected. It follows that Aut is a connected group scheme (cf. [BT2], 1.2.12). So in this case $\text{Aut} = \mathfrak{G}_{\Delta_J} = \mathfrak{G}_{\Delta_J}^0$. We conclude that $\text{Aut}(\mathcal{O})$ is the Bruhat-Tits Iwahori subgroup fixing the base alcove Λ_{\bullet} .

- EXERCISE 3.1. 1) Check the lifting criterion which shows Aut is formally smooth directly for the case $n = 2$, by explicit calculations with 2×2 matrices.
 2) By identifying $\text{Aut}(\mathcal{O})$ with its image in $\text{GL}_n(F)$ under the inclusion $g_{\bullet} \mapsto g_0$, show that the Iwahori subgroup is the preimage of $B(k)$ under the canonical surjection $\text{GL}_n(\mathcal{O}) \rightarrow \text{GL}_n(k)$.
 3) Prove that $\text{Aut}_k \rightarrow \text{Aut}(\Lambda_{0,k})$, $g_{\bullet} \mapsto g_0$, has image B_k , and kernel a connected unipotent group.

4. Local models

Given a certain triple (G, μ, K_p) consisting of a \mathbb{Z}_p -group G , a minuscule coweight μ for G , and a parahoric subgroup $K_p \subset G(\mathbb{Q}_p)$, one may construct a projective \mathbb{Z}_p -scheme \mathbf{M}^{loc} which (étale locally) models the singularities found in the special fiber of a certain Shimura variety with parahoric-level structure at p . The advantage of \mathbf{M}^{loc} is that it is defined in terms of linear algebra and is therefore easier to study than the Shimura variety itself. These schemes are called “local models”, or sometimes “Rapoport-Zink local models”; the most general treatment is given in [RZ], but in special cases they were also investigated in [DP] and [deJ].

In this section we recall the definitions of local models associated to GL_n and GSp_{2n} . For simplicity, we limit the discussion to models for Iwahori-level structure. In each case, the local model is naturally associated to a dominant minuscule coweight μ , which we shall always mention. In fact, it turns out that if the Shimura

data give rise to (G, μ) , then the Rapoport-Zink local model \mathbf{M}^{loc} (for Iwahori-level structure) can be identified with the space $M_{-w_0\mu}$ mentioned in §2. See Remark 4.1 below.

4.1. Linear case. We use the notation Λ_\bullet from section 3 to denote the “standard” lattice chain over $\mathcal{O} = \mathbb{Z}_p$ here.

Fix an integer d with $1 \leq d \leq n - 1$. We define the scheme \mathbf{M}^{loc} by defining its R -points for any \mathbb{Z}_p -algebra R as the set of isomorphism classes of commutative diagrams

$$\begin{array}{ccccccc} \Lambda_{0,R} & \longrightarrow & \cdots & \longrightarrow & \Lambda_{n-1,R} & \longrightarrow & \Lambda_{n,R} \xrightarrow{p} \Lambda_{0,R} \\ \uparrow & & & & \uparrow & & \uparrow & & \uparrow \\ \mathcal{F}_0 & \longrightarrow & \cdots & \longrightarrow & \mathcal{F}_{n-1} & \longrightarrow & \mathcal{F}_n \xrightarrow{p} \mathcal{F}_0 \end{array}$$

where the vertical arrows are inclusions, and each \mathcal{F}_i is an R -submodule of $\Lambda_{i,R}$ which is Zariski-locally on R a direct factor of corank d . It turns out that this is identical to the space $M_{-w_0\mu}$ of §2, where $\mu = (0^{n-d}, (-1)^d)$. It is clear that \mathbf{M}^{loc} is represented by a closed subscheme of a product of Grassmannians over \mathbb{Z}_p , hence it is a projective \mathbb{Z}_p -scheme. One can also formulate the moduli problem using quotients of rank d instead of submodules of corank d .

Another way to formulate the same moduli problem which is sometimes useful (see [HN1]) is given by adding an indeterminate t to the story (following a suggestion of G. Laumon). We replace the “standard” lattice chain term $\Lambda_{i,R}$ with $\mathcal{V}_{i,R} := \alpha^{-i}R[t]^n$, where α is the $n \times n$ matrix

$$\begin{pmatrix} 0 & 1 & & & \\ & & \ddots & \ddots & \\ & & & 0 & 1 \\ t+p & & & & 0 \end{pmatrix} \in \text{GL}_n(R[t, t^{-1}, (t+p)^{-1}]).$$

One can identify $\mathbf{M}^{\text{loc}}(R)$ with the set of chains

$$\mathcal{L}_\bullet = (\mathcal{L}_0 \subset \mathcal{L}_1 \subset \cdots \subset \mathcal{L}_n = (t+p)^{-1}\mathcal{L}_0)$$

of $R[t]$ -submodules of $R[t, t^{-1}, (t+p)^{-1}]^n$ satisfying the following properties

- (1) for all $i = 0, \dots, n - 1$, we have $t\mathcal{V}_{i,R} \subset \mathcal{L}_i \subset \mathcal{V}_{i,R}$;
- (2) as an R -module, $\mathcal{L}_i/t\mathcal{V}_{i,R}$ is locally a direct factor of $\mathcal{V}_{i,R}/t\mathcal{V}_{i,R}$ of corank d .

REMARK 4.1. With the second definition, it is easy to see that the geometric generic fiber $\mathbf{M}_{\mathbb{Q}_p}^{\text{loc}}$ is contained in the affine Grassmannian $\text{GL}_n(\overline{\mathbb{Q}_p}((t)))/\text{GL}_n(\overline{\mathbb{Q}_p}[[t]])$, and the geometric special fiber $\mathbf{M}_{\mathbb{F}_p}^{\text{loc}}$ is contained in the affine flag variety $\text{GL}_n(\overline{\mathbb{F}_p}((t)))/I_{\overline{\mathbb{F}_p}}$, where $I_{\overline{\mathbb{F}_p}} := \text{Aut}(\overline{\mathbb{F}_p}[[t]])$ is the Iwahori subgroup of $\text{GL}_n(\overline{\mathbb{F}_p}[[t]])$ corresponding to the upper triangular Borel subgroup $B \subset \text{GL}_n$. Moreover, it is possible to view \mathbf{M}^{loc} as a piece of a \mathbb{Z}_p -ind-scheme M which forms a deformation of the affine Grassmannian to the affine flag variety over the base $\text{Spec}(\mathbb{Z}_p)$, in analogy with Beilinson’s deformation Fl_X over a smooth \mathbb{F}_p -curve X in the function field case (cf. [Ga], [HN1]). In fact, letting e_0 denote the base point in the affine Grassmannian $\text{GL}_n(\mathbb{Q}_p((t)))/\text{GL}_n(\mathbb{Q}_p[[t]])$, and for $\lambda \in X_+(T)$, letting \mathcal{Q}_λ denote the $\text{GL}_n(\mathbb{Q}_p[[t]])$ -orbit of the point $\lambda(t)e_0$, it turns out that \mathbf{M}^{loc} coincides with

the scheme-theoretic closure $M_{-w_0\mu}$ of $\mathcal{Q}_{-w_0\mu}$ taken in the model M . A similar statement holds in the symplectic case.

The identification $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$ is explained in §8. It is a consequence of the determinant condition and the “homology” definition of our local models; also the flatness of \mathbf{M}^{loc} (see §7) plays a role.

4.2. The symplectic case. Recall that for our group $\text{GSp}_{2n} = \text{GSp}(V, (\cdot, \cdot))$, we have an identification of $X_*(T)$ with the lattice $\{(a_1, \dots, a_n, b_n, \dots, b_1) \in \mathbb{Z}^{2n} \mid \exists c \in \mathbb{Z}, a_i + b_i = c, \forall i\}$. The Shimura coweight that arises here has the form $\mu = (0^n, (-1)^n)$.

For the group GSp_{2n} , the symbol Λ_\bullet now denotes the self-dual \mathbb{Z}_p -lattice chain in \mathbb{Q}_p^{2n} , discussed in section 3 in the context of GSp_{2n} . Let (\cdot, \cdot) denote the alternating pairing on \mathbb{Z}_p^{2n} discussed in that section, and let the dual Λ^\perp of a lattice Λ be defined using (\cdot, \cdot) . As above, there are (at least) two equivalent ways to define $\mathbf{M}^{\text{loc}}(R)$ for a \mathbb{Z}_p -algebra R . We define $\mathbf{M}^{\text{loc}}(R)$ to be the set of isomorphism classes of diagrams

$$\begin{array}{ccccccc} \Lambda_{0,R} & \longrightarrow & \cdots & \longrightarrow & \Lambda_{n-1,R} & \longrightarrow & \Lambda_{n,R} \\ \uparrow & & & & \uparrow & & \uparrow \\ \mathcal{F}_0 & \longrightarrow & \cdots & \longrightarrow & \mathcal{F}_{n-1} & \longrightarrow & \mathcal{F}_n \end{array}$$

where the vertical arrows are inclusions of R -submodules with the following properties:

- (1) for $i = 0, \dots, n$, Zariski-locally on R the submodule \mathcal{F}_i is a direct factor of $\Lambda_{i,R}$ of corank n ;
- (2) \mathcal{F}_0 is isotropic with respect to (\cdot, \cdot) and \mathcal{F}_n is isotropic with respect to $p(\cdot, \cdot)$.

As in the linear case, this can be described in a way more transparently connected to affine flag varieties. In this case, $\mathcal{V}_{i,R}$ has the same meaning as in the linear case, except that the ambient space is now $R[t, t^{-1}, (t+p)^{-1}]^{2n}$. We may describe $\mathbf{M}^{\text{loc}}(R)$ as the set of chains

$$\mathcal{L}_\bullet = (\mathcal{L}_0 \subset \mathcal{L}_1 \subset \cdots \subset \mathcal{L}_n)$$

of $R[t]$ -submodules of $R[t, t^{-1}, (t+p)^{-1}]^{2n}$ satisfying the following properties

- (1) for $i = 0, 1, \dots, n$, $t\mathcal{V}_{i,R} \subset \mathcal{L}_i \subset \mathcal{V}_{i,R}$;
- (2) as R -modules, $\mathcal{L}_i/t\mathcal{V}_{i,R}$ is locally a direct factor of $\mathcal{V}_{i,R}/t\mathcal{V}_{i,R}$ of corank n ;
- (3) \mathcal{L}_0 is self-dual with respect to $t^{-1}(\cdot, \cdot)$, and \mathcal{L}_n is self-dual with respect to $t^{-1}(t+p)(\cdot, \cdot)$.

4.3. Generic and special fibers. In the linear case with $\mu = (0^{n-d}, (-1)^d)$, the generic fiber of $M_{-w_0\mu}$ is the Grassmannian $\text{Gr}(d, n)$ of d -planes in \mathbb{Q}_p^n . In the symplectic case with $\mu = (0^n, (-1)^n)$, the generic fiber of $M_{-w_0\mu}$ is the Grassmannian of isotropic n -planes in \mathbb{Q}_p^{2n} with respect to the alternating pairing (\cdot, \cdot) .

In each case, the special fiber is a union of finitely many Iwahori-orbits IwI/I in the affine flag variety, indexed by elements w in the extended affine Weyl group \widetilde{W} for GL_n (resp. GSp_{2n}) ranging over the so-called $-w_0\mu$ -admissible subset $\text{Adm}(-w_0\mu)$, [KR]. Let $\lambda \in X_+(T)$. Then by definition

$$\text{Adm}(\lambda) = \{w \in \widetilde{W} \mid \exists \nu \in W\lambda, \text{ such that } w \leq t_\nu\}.$$

Here, $W\lambda$ is the set of conjugates of λ under the action of the finite Weyl group W_0 , and $t_\nu \in \widetilde{W}$ is the translation element corresponding to ν , and \leq denotes the Bruhat order on \widetilde{W} . Actually (see §8.1), the set that arises naturally from the moduli problem is the $-w_0\mu$ -permissible subset $\text{Perm}(-w_0\mu) \subset \widetilde{W}$ from [KR]. Let us recall the definition of this set, following loc. cit. Let $\lambda \in X_+(T)$ and suppose $t_\lambda \in W_{\text{aff}}\tau$, for $\tau \in \Omega$. Then $\text{Perm}(\lambda)$ consists of the elements $x \in W_{\text{aff}}\tau$ such that $x(a) - a \in \text{Conv}(\lambda)$ for every vertex $a \in \mathbf{a}$. Here $\text{Conv}(\lambda)$ denotes the convex hull of $W\lambda$ in $X_*(T) \otimes \mathbb{R}$.

The strata in the special fiber of $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$ are naturally indexed by the set $\text{Perm}(-w_0\mu)$, which agrees with $\text{Adm}(-w_0\mu)$ by the following non-trivial combinatorial theorem due to Kottwitz and Rapoport.

THEOREM 4.2 ([KR]; see also [HN2]). *For every minuscule coweight λ of either GL_n or GSp_{2n} , we have the equality*

$$\text{Perm}(\lambda) = \text{Adm}(\lambda).$$

Using the well-known correspondence between elements in the affine Weyl group and the set of alcoves in the standard apartment of the Bruhat-Tits building, one can “draw” pictures of $\text{Adm}(\mu)$ for low-rank groups. Figures 1 and 2 depict this set for $G = \text{GL}_3$, $\mu = (-1, 0, 0)$, and $G = \text{GSp}_4$, $\mu = (-1, -1, 0, 0)$ ². Actually, we draw the image of $\text{Adm}(\mu)$ in the apartment for PGL_3 (resp. PGSp_4); the base alcove is labeled by τ .

4.4. Computing the singularities in the special fiber of \mathbf{M}^{loc} . In certain cases, the singularities in $\mathbf{M}_{\mathbb{F}_p}^{\text{loc}}$ can be analyzed directly by writing down equations. As the simplest example of how this is done, we analyze the local model for GL_2 , $\mu = (0, -1)$. For a \mathbb{Z}_p -algebra R , we are looking at the set of pairs $(\mathcal{F}_0, \mathcal{F}_1)$ of locally free rank 1 R -submodules of R^2 such that the following diagram commutes

$$\begin{array}{ccccc} R \oplus R & \xrightarrow{\begin{bmatrix} p & 0 \\ 0 & 1 \end{bmatrix}} & R \oplus R & \xrightarrow{\begin{bmatrix} 1 & 0 \\ 0 & p \end{bmatrix}} & R \oplus R \\ \uparrow & & \uparrow & & \uparrow \\ \mathcal{F}_0 & \longrightarrow & \mathcal{F}_1 & \longrightarrow & \mathcal{F}_0 \end{array}$$

Obviously this functor is represented by a certain closed subscheme of $\mathbb{P}_{\mathbb{Z}_p}^1 \times \mathbb{P}_{\mathbb{Z}_p}^1$. Locally around a fixed point $(\mathcal{F}_0, \mathcal{F}_1) \in \mathbb{P}^1(R) \times \mathbb{P}^1(R)$ we choose coordinates such that \mathcal{F}_0 is represented by the homogeneous column vector $[1 : x]^t$ and \mathcal{F}_1 by the vector $[y : 1]^t$, for $x, y \in R$. We see that $(\mathcal{F}_0, \mathcal{F}_1) \in \mathbf{M}^{\text{loc}}(R)$ if and only if

$$xy = p,$$

so \mathbf{M}^{loc} is locally the same as $\text{Spec}(\mathbb{Z}_p[X, Y]/(XY - p))$, the usual deformation of $\mathbb{A}_{\mathbb{Q}_p}^1$ to a union of two $\mathbb{A}_{\mathbb{F}_p}^1$'s which intersect transversally at a point. Indeed, \mathbf{M}^{loc} is globally this kind of deformation:

- In the generic fiber, the matrices are invertible and so \mathcal{F}_0 uniquely determines \mathcal{F}_1 ; thus $\mathbf{M}_{\mathbb{Q}_p}^{\text{loc}} \cong \mathbb{P}_{\mathbb{Q}_p}^1$;

²Note that in Figure 2 there is an alcove of length one contained in the Bruhat-closure of all four distinct Siegel translations. This already tells us something about the singularities: the special fiber of the Siegel variety for GSp_4 is not a union of divisors with normal crossings; see §8.




FIGURE 1. The admissible alcoves $\text{Adm}(\mu)$ for GL_3 , $\mu = (-1, 0, 0)$. The base alcove is labeled by τ .

- In the special fiber, $p = 0$ and one can check that $\mathbf{M}_{\mathbb{F}_p}^{\text{loc}}$ is the union of the closures of two Iwahori-orbits in the affine flag variety $\text{GL}_2(\mathbb{F}_p((t)))/I_{\mathbb{F}_p}$, each of dimension 1, which meet in a point. Thus $\mathbf{M}_{\mathbb{F}_p}^{\text{loc}}$ is the union of two $\mathbb{P}_{\mathbb{F}_p}^1$'s meeting in a point.

We refer to the work of U. Görtz for many more complicated calculations of this kind: [Go1], [Go2], [Go3], [Go4].

5. Some PEL Shimura varieties with parahoric level structure at p

5.1. PEL-type data. Given a Shimura datum $(\mathbf{G}, \{h\}, \mathbf{K})$ one can construct a Shimura variety $Sh(\mathbf{G}, h)_{\mathbf{K}}$ which has a canonical model over the reflex field \mathbf{E} , a number field determined by the datum. We write G for the p -adic group $\mathbf{G}_{\mathbb{Q}_p}$. Let us assume that the compact open subgroup $\mathbf{K} \subset \mathbf{G}(\mathbb{A}_f)$ is of the form $\mathbf{K} = K^p K_p$, where $K^p \subset \mathbf{G}(\mathbb{A}_f^p)$ is a sufficiently small compact open subgroup, and $K_p \subset G(\mathbb{Q}_p)$ is a parahoric subgroup.

Let us fix once and for all embeddings $\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$, and $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}_p}$. We denote by \mathfrak{p} the corresponding place of \mathbf{E} over p and by $E = \mathbf{E}_{\mathfrak{p}}$ the completion of \mathbf{E} at \mathfrak{p} .

If the Shimura datum comes from PEL-type data, then it is possible to define a moduli problem (in terms of chains of abelian varieties with additional structure)




FIGURE 2. The admissible alcoves $\text{Adm}(\mu)$ for GSp_4 , $\mu = (-1, -1, 0, 0)$.

over the ring \mathcal{O}_E . This moduli problem is representable by a quasi-projective \mathcal{O}_E -scheme whose generic fiber is the base-change to E of the initial Shimura variety $Sh(\mathbf{G}, h)_{\mathbf{K}}$ (or at least a finite union of Shimura varieties, one of which is the canonical model $Sh(\mathbf{G}, h)_{\mathbf{K}}$). This is done in great generality in Chapter 6 of [RZ]. Our aim in this section is only to make somewhat more explicit the definitions in loc. cit., in two very special cases attached to the linear and symplectic groups.

First, let us recall briefly PEL-type data. Let B denote a finite-dimensional semi-simple \mathbb{Q} -algebra with positive involution ι . Let $V \neq 0$ be a finitely-generated left B -module, and let (\cdot, \cdot) be a non-degenerate alternating form $V \times V \rightarrow \mathbb{Q}$ on the underlying \mathbb{Q} -vector space, such that $(bv, w) = (v, b^t w)$, for $b \in B$, $v, w \in V$. The form (\cdot, \cdot) determines a “transpose” involution on $\text{End}(V)$, denoted by $*$ (so viewing the left-action of b as an element of $\text{End}(V)$, we have $b^t = b^*$). We denote by \mathbf{G} the \mathbb{Q} -group whose points in a \mathbb{Q} -algebra R are

$$\{g \in \text{GL}_{B \otimes R}(V \otimes R) \mid g^* g = c(g) \in R^\times\}.$$

We assume \mathbf{G} is a *connected* reductive group; this means we are excluding the orthogonal case. Consider the \mathbb{R} -algebra $C := \text{End}_B(V) \otimes \mathbb{R}$. We let $h_0 : \mathbb{C} \rightarrow C$ denote an \mathbb{R} -algebra homomorphism satisfying $h_0(\bar{z}) = h_0(z)^*$, for $z \in \mathbb{C}$. We fix a choice of $i = \sqrt{-1}$ in \mathbb{C} once and for all, and we assume the symmetric bilinear form $(\cdot, h_0(i)\cdot) : V_{\mathbb{R}} \times V_{\mathbb{R}} \rightarrow \mathbb{R}$ is positive definite. Let h denote the inverse of the restriction of h_0 to \mathbb{C}^\times . Then h induces an algebraic homomorphism

$$h : \mathbb{C}^\times \rightarrow \mathbf{G}(\mathbb{R})$$

of real groups which defines on $V_{\mathbb{R}}$ a Hodge structure of type $(1, 0) + (0, 1)$ (in the terminology of [Del2], section 1) and which satisfies the usual Riemann conditions with respect to (\cdot, \cdot) (see [Ko92], Lemma 4.1). For any choice of (sufficiently small) compact open subgroup \mathbf{K} , the data $(\mathbf{G}, h, \mathbf{K})$ determine a (smooth) Shimura variety over a reflex field \mathbf{E} ; cf. [Del].

We recall that h gives rise to a minuscule coweight

$$\mu := \mu_h : \mathbb{G}_{m, \mathbb{C}} \rightarrow \mathbf{G}_{\mathbb{C}}$$

as follows: the complexification of the real group \mathbb{C}^{\times} is the torus $\mathbb{C}^{\times} \times \mathbb{C}^{\times}$, the factors being indexed by the two \mathbb{R} -algebra automorphisms of \mathbb{C} ; we assume the first factor corresponds to the identity and the second to complex conjugation. Then we define

$$\mu(z) := h_{\mathbb{C}}(z, 1).$$

By definition of Shimura data, the homomorphism $h : \mathbb{C}^{\times} \rightarrow \mathbf{G}_{\mathbb{R}}$ is only specified up to $\mathbf{G}(\mathbb{R})$ -conjugation, and therefore μ is only well-defined up to $\mathbf{G}(\mathbb{C})$ -conjugation. However, this conjugacy class is at least defined over the reflex field \mathbf{E} (in fact we define \mathbf{E} as the field of definition of the conjugacy class of μ). Via our choice of field embeddings $\overline{\mathbb{C}} \hookrightarrow \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$, we get a well-defined conjugacy class of minuscule coweights

$$\mu : \mathbb{G}_{m, \overline{\mathbb{Q}}_p} \rightarrow G_{\overline{\mathbb{Q}}_p},$$

which is defined over E .

The argument of [Ko84], Lemma (1.1.3) shows that E is contained in any subfield of $\overline{\mathbb{Q}}_p$ which splits G . Therefore, when G is split over \mathbb{Q}_p (the case of interest in this report), it follows that $E = \mathbb{Q}_p$ and the conjugacy class of μ contains a \mathbb{Q}_p -rational and B -dominant element, usually denoted also by the symbol μ . It is this same μ which was mentioned in the definitions of local models in section 4.

For use in the definition to follow, we decompose the $B_{\mathbb{C}}$ -module $V_{\mathbb{C}}$ as $V_{\mathbb{C}} = V_1 \oplus V_2$, where $h_0(z)$ acts by z on V_1 and by \bar{z} on V_2 , for $z \in \mathbb{C}$. Our conventions imply that $\mu(z)$ acts by z^{-1} on V_1 and by 1 on V_2 ($z \in \mathbb{C}^{\times}$). We choose $E' \subset \overline{\mathbb{Q}}_p$ a finite extension field $E' \supset E$ over which this decomposition is defined:

$$V_{E'} = V_1 \oplus V_2.$$

(We are implicitly using the diagram $\mathbb{C} \hookrightarrow \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$ to make sense of this.)

Recall that we are interested in defining an \mathcal{O}_E -integral model for $Sh(\mathbf{G}, h)_{\mathbf{K}}$ in the case where $K_p \subset \mathbf{G}(\mathbb{Q}_p)$ is a parahoric (more specifically, an Iwahori) subgroup.

To define an integral model over \mathcal{O}_E , we need to specify certain additional data. We suppose \mathcal{O}_B is a $\mathbb{Z}_{(p)}$ -order in B whose p -adic completion $\mathcal{O}_B \otimes \mathbb{Z}_p$ is a maximal order in $B_{\mathbb{Q}_p}$, stable under the involution ι . Using the terminology of [RZ], 6.2, we assume we are given a self-dual multichain \mathcal{L} of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices in $V_{\mathbb{Q}_p}$ (the notion of multichain \mathcal{L} is a generalization of the lattice chain Λ_{\bullet} appearing in section 4; specifying \mathcal{L} is equivalent to specifying a parahoric subgroup, namely $K_p := \text{Aut}(\mathcal{L})$, of $\mathbf{G}(\mathbb{Q}_p)$). We can then give the definition of a model Sh_{K^p} that depends on the above data and the choice of a small compact open subgroup K^p ³.

³In the sequel, we sometimes drop the subscript K^p on Sh_{K^p} , or replace it with the subscript K_p , depending on whether K^p , or K_p (or both) is understood.

DEFINITION 5.1. A point of the functor Sh_{K^p} with values in the \mathcal{O}_E -scheme S is given by the following set of data up to isomorphism ⁴.

- (1) An \mathcal{L} -set of abelian S -schemes $A = \{A_\Lambda\}$, $\Lambda \in \mathcal{L}$, compatibly endowed with an action of \mathcal{O}_B :

$$i : \mathcal{O}_B \otimes \mathbb{Z}_{(p)} \rightarrow \text{End}(A) \otimes \mathbb{Z}_{(p)};$$

- (2) A \mathbb{Q} -homogeneous principal polarization $\bar{\lambda}$ of the \mathcal{L} -set A ;
- (3) A K^p -level structure

$$\bar{\eta} : V \otimes \mathbb{A}_f^p \cong H_1(A, \mathbb{A}_f^p) \text{ mod } K^p$$

that respects the bilinear forms on both sides up to a scalar in $(\mathbb{A}_f^p)^\times$, and commutes with the $B = \mathcal{O}_B \otimes \mathbb{Q}$ -actions.

We impose the condition that under

$$i : \mathcal{O}_B \otimes \mathbb{Z}_{(p)} \rightarrow \text{End}(A) \otimes \mathbb{Z}_{(p)},$$

we have $i(b') = \lambda^{-1} \circ (i(b))^\vee \circ \lambda$; in other words, i intertwines ι and the Rosati involution on $\text{End}(A) \otimes \mathbb{Z}_{(p)}$ determined by $\bar{\lambda}$. In addition, we impose the following determinant condition: for each $b \in \mathcal{O}_B$ and $\Lambda \in \mathcal{L}$:

$$\det_{\mathcal{O}_S}(b, \text{Lie}(A_\Lambda)) = \det_{E'}(b, V_1).$$

We will not explain all the notions entering this definition; we refer to loc. cit., Chapter 6 as well as [Ko92], section 5, for complete details. However, in the simple examples we make explicit below, these notions will be made concrete and their importance will be highlighted. For example, an \mathcal{L} -set of abelian varieties $\{A_\Lambda\}$ comes with a family of “periodicity isomorphisms”

$$\theta_a : A_\Lambda^a \rightarrow A_{a\Lambda},$$

see [RZ], Def. 6.5, and we will describe these explicitly in the examples to follow.

Note that one can see from this definition why some of the conditions on PEL data are imposed. For example, since the Rosati involution is always positive (see [Mu], section 21), we see that the involution ι on B must be positive for the moduli problem to be non-empty.

5.2. Some “fake” unitary Shimura varieties. This section concerns the so-called “simple” or “fake unitary” Shimura varieties investigated by Kottwitz in [Ko92b]. They are indeed “simple” in the sense that they are compact Shimura varieties for which there are no problems due to endoscopy (see loc. cit.).

Kottwitz made assumptions ensuring that the local group $\mathbf{G}_{\mathbb{Q}_p}$ be unramified, and that the level structure at p be given by a hyperspecial maximal compact subgroup. Here we will work in a situation where $\mathbf{G}_{\mathbb{Q}_p}$ is *split*, but we only impose parahoric-level structure at p . For simplicity, we explain only the case $F_0 = \mathbb{Q}$ (notation of loc. cit.).

⁴We say $\{A_\Lambda\}$ is isomorphic to $\{A'_\Lambda\}$ if there is a compatible family of prime-to- p isogenies $A_\Lambda \rightarrow A'_\Lambda$ which preserve all the structures.

5.2.1. *The group-theoretic set-up.* Let F be an imaginary quadratic extension of \mathbb{Q} , and let $(D, *)$ be a division algebra with center F , of dimension n^2 over F , together with an involution $*$ which induces on F the non-trivial element of $\text{Gal}(F/\mathbb{Q})$. Let \mathbf{G} be the \mathbb{Q} -group whose points in a commutative \mathbb{Q} -algebra R are

$$\{x \in D \otimes_{\mathbb{Q}} R \mid x^*x \in R^\times\}.$$

The map $x \mapsto x^*x$ is a homomorphism of \mathbb{Q} -groups $\mathbf{G} \rightarrow \mathbb{G}_m$ whose kernel \mathbf{G}_0 is an inner form of a unitary group over \mathbb{Q} associated to F/\mathbb{Q} . Let us suppose we are given an \mathbb{R} -algebra homomorphism

$$h_0 : \mathbb{C} \rightarrow D \otimes_{\mathbb{Q}} \mathbb{R}$$

such that $h_0(z)^* = h_0(\bar{z})$ and the involution $x \mapsto h_0(i)^{-1}x^*h_0(i)$ is positive.

Given the data $(D, *, h_0)$ above, we want to explain how to find the PEL-data $(B, \iota, V, (\cdot, \cdot), h_0)$ used in the definition of the scheme Sh_{K^p} .

Let $B = D^{opp}$ and let $V = D$ be viewed as a left B -module, free of rank 1, using right multiplications. Thus we can identify $C := \text{End}_B(V)$ with D (left multiplications). For $h_0 : \mathbb{C} \rightarrow C_{\mathbb{R}}$ we use the homomorphism $h_0 : \mathbb{C} \rightarrow D \otimes_{\mathbb{Q}} \mathbb{R}$ we are given.

Next, one can show that there exist elements $\xi \in D^\times$ such that $\xi^* = -\xi$ and the involution $x \mapsto \xi x^* \xi^{-1}$ is positive. To see this, note that the Skolem-Noether theorem implies that the involutions of the second type on D are precisely the maps of the form $x \mapsto bx^*b^{-1}$, for $b \in D^\times$ such that $b(b^*)^{-1}$ lies in the center F . Since positive involutions of the second kind exist (see [Mu], p. 201-2), for some such b the involution $x \mapsto bx^*b^{-1}$ is positive. We have $N_{F/\mathbb{Q}}(b(b^*)^{-1}) = 1$, so by Hilbert's Theorem 90, we may alter any such b by an element in F^\times so that $b^* = b$. There exists $\epsilon \in F^\times$ such that $\epsilon^* = -\epsilon$. We then put $\xi = \epsilon b$.

We define the positive involution ι by $x^\iota := \xi x^* \xi^{-1}$, for $x \in B = D^{opp}$.

Now we define the non-degenerate alternating pairing $(\cdot, \cdot) : D \times D \rightarrow \mathbb{Q}$ by

$$(x, y) = \text{tr}_{D/\mathbb{Q}}(x\xi y^*).$$

It is clear that $(bx, y) = (x, b^\iota y)$ for any $b \in B = D^{opp}$, remembering that the left action of b is right multiplication by b . We also have $(h_0(z)x, y) = (x, h_0(\bar{z})y)$, since $h_0(z) \in D$ acts by left multiplication on D .

Finally, we claim that $(\cdot, h_0(i)\cdot)$ is always positive or negative definite; thus we can always arrange for it to be positive definite by replacing ξ with $-\xi$ if necessary. To prove the definiteness, choose an isomorphism

$$D \otimes_{\mathbb{Q}} \mathbb{R} \xrightarrow{\sim} M_n(\mathbb{C})$$

such that the positive involution $x \mapsto x^\iota$ goes over to the standard positive involution $X \mapsto \bar{X}^t$ on $M_n(\mathbb{C})$. Let $H \in M_n(\mathbb{C})$ be the image of $\xi h_0(i)^{-1}$ under this isomorphism, so that the symmetric pairing $\langle x, y \rangle = (x, h_0(i)y)$ goes over to the pairing

$$\langle X, Y \rangle = \text{tr}_{M_n(\mathbb{C})/\mathbb{R}}(X \bar{Y}^t H).$$

We conclude by invoking the following exercise for the reader.

EXERCISE 5.2. The matrix H is Hermitian and either positive or negative definite. If positive definite, we then have $\text{tr}(X \bar{X}^t H) > 0$ whenever $X \neq 0$.

(Hint: use the argument of [Mu], p. 200.)

5.2.2. *The minuscule coweight μ .* How is the minuscule coweight μ described in terms of the above data? Recall our decomposition

$$D_{\mathbb{C}} = V_1 \oplus V_2.$$

The homomorphism h_0 makes $D_{\mathbb{C}}$ into a $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$ -module. Of course

$$\begin{aligned} \mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} &\xrightarrow{\sim} \mathbb{C} \times \mathbb{C} \\ z_1 \otimes z_2 &\mapsto (z_1 z_2, \bar{z}_1 z_2), \end{aligned}$$

which induces the above decomposition of $D_{\mathbb{C}}$ ($h_0(z_1 \otimes 1)$ acts by z_1 on V_1 and by \bar{z}_1 on V_2).

The factors V_1, V_2 are stable under right multiplications of

$$D_{\mathbb{C}} = D \otimes_{F, \nu} \mathbb{C} \times D \otimes_{F, \nu^*} \mathbb{C},$$

where $\nu, \nu^* : F \hookrightarrow \mathbb{C}$ are the two embeddings. (We may assume our fixed choice $\bar{\mathbb{Q}} \hookrightarrow \mathbb{C}$ extends ν .) Also $h_0(z \otimes 1)$ is the endomorphism given by left multiplication by a certain element of $D_{\mathbb{C}}$. We can choose an isomorphism

$$D \otimes_{F, \nu} \mathbb{C} \times D \otimes_{F, \nu^*} \mathbb{C} \cong M_n(\mathbb{C}) \times M_n(\mathbb{C})$$

such that $h_0(z \otimes 1)$ can be written explicitly as

$$h_0(z \otimes 1) = \text{diag}(\bar{z}^{n-d}, z^d) \times \text{diag}(z^{n-d}, \bar{z}^d),$$

for some integer d , $0 \leq d \leq n$. (One can then identify V_1 resp. V_2 as the span of certain columns of the two matrices.) We know that $\mu(z) = h_{\mathbb{C}}(z, 1)$ acts by z^{-1} on V_1 and by 1 on V_2 . Hence we can identify $\mu(z)$ as

$$\mu(z) = \text{diag}(1^{n-d}, (z^{-1})^d) \times \text{diag}((z^{-1})^{n-d}, 1^d).$$

We may label this by $(0^{n-d}, (-1)^d) \in \mathbb{Z}^n$, via the usual identification applied to the first factor.

Here is another way to interpret the number d . Let W (resp. W^*) be the (unique up to isomorphism, n -dimensional) simple right-module for $D \otimes_{F, \nu} \mathbb{C}$ (resp. $D \otimes_{F, \nu^*} \mathbb{C}$). Then as right $D_{\mathbb{C}}$ -modules we have

$$V_1 = W^d \oplus (W^*)^{n-d}, \quad \text{resp. } V_2 = W^{n-d} \oplus (W^*)^d.$$

Finally, let us remark that if we choose the identification of $D \otimes_{\mathbb{Q}} \mathbb{R} = D \otimes_{F, \nu} \mathbb{C}$ with $M_n(\mathbb{C})$ in such a way that the positive involution $x \mapsto h_0(i)^{-1} x^* h_0(i)$ goes over to $X \mapsto \bar{X}^t$, then we get an isomorphism

$$G(\mathbb{R}) \cong \text{GU}(d, n-d).$$

(See also [Ko92b], section 1.)

In applications, it is sometimes necessary to prescribe the value of d ahead of time (with the additional constraint that $1 \leq d \leq n-1$). However, it can be a delicate matter to arrange things so that a prescribed value of d is achieved. To see how this is done for the case of $d=1$, see [HT], Lemma I.7.1.

5.2.3. *Assumptions on p and integral data.* We first make some assumptions on the prime p ⁵, and then we specify the integral data at p .

First assumption on p : The prime p splits in F as a product of distinct prime ideals

$$(p) = \mathfrak{p}\mathfrak{p}^*,$$

where \mathfrak{p} is the prime determined by our fixed choice of embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$, and \mathfrak{p}^* is its image under the non-trivial element of $\text{Gal}(F/\mathbb{Q})$.

Under this assumption $F_{\mathfrak{p}} = F_{\mathfrak{p}^*} = \mathbb{Q}_p$. Further the algebra $D_{\mathbb{Q}_p}$ is a product

$$D \otimes \mathbb{Q}_p = D_{\mathfrak{p}} \times D_{\mathfrak{p}^*}$$

where each factor is a central simple \mathbb{Q}_p -algebra. We have $D_{\mathfrak{p}} \xrightarrow{\sim} D_{\mathfrak{p}^*}^{opp}$ via $*$. Therefore for any \mathbb{Q}_p -algebra R we can identify the group $G(R)$ with the group

$$\{(x_1, x_2) \in (D_{\mathfrak{p}} \otimes R)^{\times} \times (D_{\mathfrak{p}^*} \otimes R)^{\times} \mid x_1 = c(x_2^*)^{-1}, \text{ for some } c \in R^{\times}\}.$$

Therefore there is an isomorphism of \mathbb{Q}_p -groups $G \cong D_{\mathfrak{p}}^{\times} \times \mathbb{G}_m$ given by $(x_1, x_2) \mapsto (x_1, c)$.

Second assumption on p : The algebra $D_{\mathbb{Q}_p}$ splits: $D_{\mathfrak{p}} \cong M_n(\mathbb{Q}_p)$.

In this case the involution $*$ becomes isomorphic to the involution on $M_n(\mathbb{Q}_p) \times M_n(\mathbb{Q}_p)$ given by

$$* : (X, Y) \mapsto (Y^t, X^t).$$

Our assumptions imply that $G = \text{GL}_n \times \mathbb{G}_m$, a split p -adic group (and thus $E := \mathbf{E}_{\mathfrak{p}} = \mathbb{Q}_p$). Why is this helpful? As we shall see, this allows us to use the local models for GL_n described in section 4 to describe the reduction modulo p of the Shimura variety Sh_{K^p} , see §6.3.3. Also, we can use the description in [HN1] of nearby cycles on such models to compute the semi-simple local zeta function at p of Sh_{K^p} , see [HN3] and Theorem 11.7. One expects that this is still possible in the general case (where $D_{\mathfrak{p}}^{\times}$ is not a split group), but there the crucial facts about nearby cycles on the corresponding local models are not yet established.

Integral data. We need to specify a $\mathbb{Z}_{(p)}$ -order $\mathcal{O}_B \subset B$ and a self-dual multichain $\mathcal{L} = \{\Lambda\}$ of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices. To give a multichain we need to specify first a (partial) \mathbb{Z}_p -lattice chain in $V_{\mathbb{Q}_p} = D_{\mathfrak{p}} \times D_{\mathfrak{p}^*}$. We do this one factor at a time. First, we may fix an isomorphism

$$(5.2.1) \quad D_{\mathbb{Q}_p} = D_{\mathfrak{p}} \times D_{\mathfrak{p}^*} \cong M_n(\mathbb{Q}_p) \times M_n(\mathbb{Q}_p)$$

such that the involution $x \mapsto x^t = \xi x^* \xi^{-1}$ goes over to $(X, Y) \mapsto (Y^t, X^t)$. So ξ gets identified with an element of the form $(\chi^t, -\chi)$, for $\chi \in \text{GL}_n(\mathbb{Q}_p)$ ⁶, and our pairing $\langle x, y \rangle = \text{tr}_{D/\mathbb{Q}}(x\xi y^*) = \text{tr}_{D/\mathbb{Q}}(xy^t \xi)$ goes over to

$$\langle (X_1, X_2), (Y_1, Y_2) \rangle = \text{tr}_{D_{\mathbb{Q}_p}/\mathbb{Q}_p}(X_1 Y_2^t \chi^t, -X_2 Y_1^t \chi).$$

Next we define a (partial) \mathbb{Z}_p -lattice chain $\Lambda_{-n}^* \subset \dots \subset \Lambda_0^* = p^{-1} \Lambda_{-n}^*$ in $D_{\mathfrak{p}^*}$ by setting

$$\Lambda_{-i}^* = \chi^{-1} \text{diag}(p^i, 1^{n-i}) M_n(\mathbb{Z}_p),$$

⁵It would make sense to include in our discussion another case, where p remains inert in F , and where the group $\mathbf{G}_{\mathbb{Q}_p}$ is a quasi-split unitary group associated to the extension F_p/\mathbb{Q}_p . However, we shall postpone discussion of this case to a future occasion.

⁶For use in §11, note that if we multiply ξ by any integral power of p , we change neither its properties nor the isomorphism class of the symplectic space $V, (\cdot, \cdot)$. Hence we may assume $\chi^{-1} \in \text{GL}_n(\mathbb{Q}_p) \cap M_n(\mathbb{Z}_p)$.

for $i = 0, 1, \dots, n$. We can then extend this by periodicity to define Λ_i^* for all $i \in \mathbb{Z}$. Similarly, we define the \mathbb{Z}_p -lattice chain $\Lambda_0 \subset \dots \subset \Lambda_n = p^{-1}\Lambda_0$ in $D_{\mathfrak{p}}$ by setting

$$\Lambda_i = \text{diag}((p^{-1})^i, 1^{n-i}) M_n(\mathbb{Z}_p),$$

for $i = 0, 1, \dots, n$ (and then extending by periodicity to define for all i). We note that

$$(\Lambda_i \oplus \Lambda_i^*)^\perp = \Lambda_{-i} \oplus \Lambda_{-i}^*,$$

where \perp is defined in the usual way using the pairing (\cdot, \cdot) . Setting $\mathcal{O}_B \subset B$ to be the unique maximal $\mathbb{Z}_{(p)}$ -order such that under our fixed identification $D_{\mathbb{Q}_p} \cong M_n(\mathbb{Q}_p) \times M_n(\mathbb{Q}_p)$, we have

$$\mathcal{O}_B \otimes \mathbb{Z}_p \xrightarrow{\sim} M_n^{opp}(\mathbb{Z}_p) \times M_n^{opp}(\mathbb{Z}_p),$$

one can now check that $\mathcal{L} := \{\Lambda \oplus \Lambda^*\}$ is a self-dual multichain of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices. It is clear that $(\mathcal{O}_B \otimes \mathbb{Z}_p)^\vee = \mathcal{O}_B \otimes \mathbb{Z}_p$.

5.2.4. *The moduli problem.* We have now constructed all the data that enters into the definition of Sh_{K^p} . By the determinant condition, the abelian varieties have (relative) dimension $\dim(V_1) = n^2$. An S -point in our moduli space is a chain of abelian schemes over S of relative dimension n^2 , equipped with $\mathcal{O}_B \otimes \mathbb{Z}_{(p)}$ -actions, indexed by \mathcal{L} (we set $A_i = A_{\Lambda_i \oplus \Lambda_i^*}$ for all $i \in \mathbb{Z}$)

$$\dots \xrightarrow{\alpha} A_0 \xrightarrow{\alpha} A_1 \xrightarrow{\alpha} \dots \xrightarrow{\alpha} A_n \xrightarrow{\alpha} \dots$$

such that

- each α is an isogeny of height $2n$ (i.e., of degree p^{2n});
- there is a “periodicity isomorphism” $\theta_p : A_{i+n} \rightarrow A_i$ such that for each i the composition

$$A_i \xrightarrow{\alpha} A_{i+1} \xrightarrow{\alpha} \dots \xrightarrow{\alpha} A_{i+n} \xrightarrow{\theta_p} A_i$$

is multiplication by $p : A_i \rightarrow A_i$;

- the morphisms α commute with the $\mathcal{O}_B \otimes \mathbb{Z}_{(p)}$ -actions;
- the determinant condition holds: for every i and $b \in \mathcal{O}_B$,

$$\det_{\mathcal{O}_S}(b, \text{Lie}(A_i)) = \det_{E'}(b, V_1).$$

(See [RZ], Def. 6.5.)

In addition, we have a principal polarization and a K^p -level structure (see [RZ], Def. 6.9). Giving a polarization is equivalent to giving a commutative diagram whose vertical arrows are isogenies

$$\begin{array}{ccccccccccc} \dots & \xrightarrow{\alpha} & A_{-1} & \xrightarrow{\alpha} & A_0 & \xrightarrow{\alpha} & A_1 & \xrightarrow{\alpha} & \dots & \xrightarrow{\alpha} & A_n & \xrightarrow{\alpha} & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \\ \dots & \xrightarrow{\alpha^\vee} & \widehat{A}_1 & \xrightarrow{\alpha^\vee} & \widehat{A}_0 & \xrightarrow{\alpha^\vee} & \widehat{A}_{-1} & \xrightarrow{\alpha^\vee} & \dots & \xrightarrow{\alpha^\vee} & \widehat{A}_{-n} & \xrightarrow{\alpha^\vee} & \dots \end{array}$$

such that for each i the quasi-isogeny

$$A_i \rightarrow \widehat{A}_{-i} \rightarrow \widehat{A}_i$$

is a rational multiple of a polarization of A_i . If up to a \mathbb{Q} -multiple the vertical arrows are all isomorphisms, we say the polarization is principal.

The fact that $\text{End}_B(V)$ is a division algebra implies that the moduli space Sh_{K^p} is proper over \mathcal{O}_E (Kottwitz verified the valuative criterion of properness in

the case of maximal hyperspecial level structure in [Ko92] p. 392, using the theory of Néron models; the same proof applies here.)

5.3. Siegel modular varieties with $\Gamma_0(p)$ -level structure. The set-up is much simpler here. The group \mathbf{G} is $\mathrm{GSp}(V)$ where V is the standard symplectic space \mathbb{Q}^{2n} with the alternating pairing (\cdot, \cdot) given by the matrix \tilde{I} in 3.1.2. We have $B = \mathbb{Q}$ with involution $\iota = \mathrm{id}$, and $h_0 : \mathbb{C} \rightarrow \mathrm{End}(V_{\mathbb{R}})$ is defined as the unique \mathbb{R} -algebra homomorphism such that

$$h_0(i) = \tilde{I}.$$

For the multichain \mathcal{L} we use the standard complete self-dual lattice chain Λ_{\bullet} in \mathbb{Q}_p^{2n} that appeared in section 4. We take $\mathcal{O}_B = \mathbb{Z}_{(p)}$.

The group $G = \mathrm{GSp}_{2n, \mathbb{Q}_p}$ is split, so again we have $E = \mathbb{Q}_p$, so $\mathcal{O}_E = \mathbb{Z}_p$. It turns out that the minuscule coweight μ is

$$\mu = (0^n, (-1)^n),$$

in other words, the same that appeared in the definition of local models in the symplectic case in section 4.

The moduli problem over \mathbb{Z}_p can be expressed as follows. For a \mathbb{Z}_p -scheme S , an S -point is an element of the set of 4-tuples (taken up to isomorphism)

$$\mathcal{A}_{K^p}(S) = \{(A_{\bullet}, \lambda_0, \lambda_n, \bar{\eta})\}$$

consisting of

- a chain A_{\bullet} of (relative) n -dimensional abelian varieties $A_0 \xrightarrow{\alpha} A_1 \xrightarrow{\alpha} \dots \xrightarrow{\alpha} A_n$ such that each morphism $\alpha : A_i \rightarrow A_{i+1}$ is an isogeny of degree p over S ;
- principal polarizations $\lambda_0 : A_0 \xrightarrow{\sim} \widehat{A}_0$ and $\lambda_n : A_n \xrightarrow{\sim} \widehat{A}_n$ such that the composition of

$$\begin{array}{ccccc} A_0 & \xrightarrow{\alpha} & \dots & \xrightarrow{\alpha} & A_n \\ \lambda_0^{-1} \uparrow & & & & \downarrow \lambda_n \\ \widehat{A}_0 & \xleftarrow{\alpha^\vee} & \dots & \xleftarrow{\alpha^\vee} & \widehat{A}_n \end{array}$$

- starting and ending at any A_i or \widehat{A}_i is multiplication by p ;
- a level K^p -structure $\bar{\eta}$ on A_0 .

EXERCISE 5.3. Show that the above description of \mathcal{A}_{K^p} is equivalent to the definition given in Definition 6.9 of [RZ] (cf. our Def. 5.1) for the group-theoretic data (B, ι, V, \dots) we described above.

Note that the only information imparted by the determinant condition in this case is that $\dim(A_i) = n$ for every i .

There is another convenient description of the same moduli problem, used by de Jong [deJ]. We define another moduli problem \mathcal{A}'_{K^p} whose S -points is the set of 4-tuples

$$\mathcal{A}'_{K^p}(S) = \{(A_0, \lambda_0, \bar{\eta}, H_{\bullet})\}$$

consisting of

- an n -dimensional abelian variety A_0 with principal polarization λ_0 and K^p -level structure $\bar{\eta}$;

- a chain H_\bullet of finite flat group subschemes of $A_0[p] := \ker(p : A_0 \rightarrow A_0)$ over S

$$(0) = H_0 \subset H_1 \subset \dots \subset H_n \subset A_0[p]$$

such that H_i has rank p^i over S and H_n is totally isotropic with respect to the Riemann form e_{λ_0} , defined by the diagram

$$\begin{array}{ccc} A_0[p] \times A_0[p] & \xrightarrow{e_{\lambda_0}} & \mu_p \\ \downarrow \text{id} \times \lambda_0 & & \uparrow \text{can} \\ A_0[p] \times \widehat{A_0}[p] & \xrightarrow{\cong} & A_0[p] \times \widehat{A_0}[p]. \end{array}$$

Here $\widehat{A_0}[p] = \text{Hom}(A_0[p], \mathbb{G}_m)$ denotes the Cartier dual of the finite group scheme $A_0[p]$ and can denotes the canonical pairing (which takes values in the p -th roots of unity group subscheme $\mu_p \subset \mathbb{G}_m$). (See [Mu], section 20, or [Mi], section 16.)

The isomorphism $\mathcal{A} \xrightarrow{\sim} \mathcal{A}'$ is given by

$$(A_\bullet, \lambda_0, \lambda_n, \bar{\eta}) \mapsto (A_0, \lambda_0, \bar{\eta}, H_\bullet); \quad H_i := \ker[\alpha^i : A_0 \rightarrow A_i].$$

The inverse map is given by setting $A_i = A_0/H_i$ (the condition on H_n allows us to define a principal polarization $\lambda_n : A_0/H_n \xrightarrow{\sim} (\widehat{A_0/H_n})$ using λ_0).

In [deJ], de Jong analyzed the singularities of \mathcal{A} in the case $n = 2$, and deduced that the model \mathcal{A} is flat in that case (by passing from \mathcal{A} to a local model \mathbf{M}^{loc} according to the procedure of section 6 and then by writing down equations for \mathbf{M}^{loc}).

In the sequel, we will denote the model \mathcal{A} (and \mathcal{A}') by the symbol Sh , sometimes adding the subscript K_p when the level-structure at p is not already understood.

6. Relating Shimura varieties and their local models

6.1. Local model diagrams. Here we describe the desiderata for local models of Shimura varieties. Quite generally, consider a diagram of finite-type \mathcal{O}_E -schemes

$$\mathcal{M} \xleftarrow{\varphi} \widetilde{\mathcal{M}} \xrightarrow{\psi} \mathcal{M}^{\text{loc}}.$$

DEFINITION 6.1. We call such a diagram a *local model diagram* provided the following conditions are satisfied:

- (1) the morphisms φ and ψ are smooth and φ is surjective;
- (2) étale locally $\mathcal{M} \cong \mathcal{M}^{\text{loc}}$: there exists an étale covering $V \rightarrow \mathcal{M}$ and a section $s : V \rightarrow \widetilde{\mathcal{M}}$ of φ over V such that $\psi \circ s : V \rightarrow \mathcal{M}^{\text{loc}}$ is étale.

In practice \mathcal{M} is the scheme we are interested in, and \mathcal{M}^{loc} is somehow simpler to study; $\widetilde{\mathcal{M}}$ is just some intermediate scheme used to link the other two. Every property that is local for the étale topology is shared by \mathcal{M} and \mathcal{M}^{loc} . For example, if \mathcal{M}^{loc} is flat over $\text{Spec}(\mathcal{O}_E)$, then so is \mathcal{M} . The singularities in \mathcal{M} and \mathcal{M}^{loc} are the same.

6.2. The general definition of local models. We briefly recall the general definition of local models, following [RZ], Def. 3.27. We suppose we have data G, μ, V, V_1, \dots coming from a PEL-type data as in section 5.1. We assume μ and V_1 are defined over a finite extension $E' \supset E$. We suppose we are given a self-dual multichain of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices $\mathcal{L} = \{\Lambda\}$.

DEFINITION 6.2 ([RZ], 3.27). A point of \mathbf{M}^{loc} with values in an \mathcal{O}_E -scheme S is given by the following data.

- (1) A functor from the category \mathcal{L} to the category of $\mathcal{O}_B \otimes_{\mathbb{Z}_p} \mathcal{O}_S$ -modules on S

$$\Lambda \mapsto t_\Lambda, \quad \Lambda \in \mathcal{L};$$

- (2) A morphism of functors $\psi_\Lambda : \Lambda \otimes_{\mathbb{Z}_p} \mathcal{O}_S \rightarrow t_\Lambda$.

We require the following conditions are satisfied:

- (i) t_Λ is a locally free \mathcal{O}_S -module of finite rank. For the action of \mathcal{O}_B on t_Λ we have the determinant condition

$$\det_{\mathcal{O}_S}(a; t_\Lambda) = \det_{E'}(a; V_1), \quad a \in \mathcal{O}_B;$$

- (ii) the morphisms ψ_Λ are surjective;
- (iii) for each Λ the composition of the following map is zero:

$$t_\Lambda^\vee \xrightarrow{\psi_\Lambda^\vee} (\Lambda \otimes \mathcal{O}_S)^\vee \xrightarrow{\cong} \Lambda^\perp \otimes \mathcal{O}_S \xrightarrow{\psi_{\Lambda^\perp}} t_{\Lambda^\perp}.$$

It is clear that one can associate to any PEL-type Shimura variety $Sh = Sh_{K_p}$ a scheme \mathbf{M}^{loc} (just use the same PEL-type data and multichain \mathcal{L} used to define Sh_{K_p}). It is less clear why the resulting scheme \mathbf{M}^{loc} really is a local model for Sh_{K_p} , in the sense described above. We shall see this below, thus justifying the terminology “local model”. Then we will show that in our two examples – the “fake” unitary and the Siegel cases – this definition agrees with the concrete ones defined for GL_n and GSp_{2n} in section 4.

6.3. Constructing local model diagrams for Shimura varieties.

6.3.1. *The abstract construction.* For one of our models $Sh = Sh_{K_p}$ from §5, we want to construct a local model diagram

$$Sh \xleftarrow{\varphi} \widetilde{Sh} \xrightarrow{\psi} \mathbf{M}^{\text{loc}}.$$

In the following we use freely the notation of the appendix, §14. For an abelian scheme $a : A \rightarrow S$, let $M(A)$ be the locally free \mathcal{O}_S -module dual to the de Rham cohomology

$$M^\vee(A) = H_{DR}^1(A/S) := R^1 a_* (\Omega_{A/S}^\bullet).$$

This is a locally free \mathcal{O}_S -module of rank $2 \dim(A/S)$. We have the Hodge filtration

$$0 \rightarrow \text{Lie}(\widehat{A})^\vee \rightarrow M(A) \rightarrow \text{Lie}(A) \rightarrow 0.$$

This is dual to the usual Hodge filtration on de Rham cohomology

$$0 \subset \omega_{A/S} := a_* \Omega_{A/S}^1 \subset H_{DR}^1(A/S).$$

We shall call $M(A)$ the *crystal associated to A/S* (this is perhaps non-standard terminology). If A carries an action of \mathcal{O}_B , then by functoriality so does $M(A)$. Note also that $M(A)$ is covariant as a functor of A . So if \mathcal{L} denotes a self-dual multichain of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices, and $\{A_\Lambda\}$ denotes an \mathcal{L} -set of abelian schemes over S with \mathcal{O}_B -action and polarization (as in Definition 5.1), then applying the functor $M(\cdot)$ gives us a polarized multichain $\{M(A_\Lambda)\}$ of $\mathcal{O}_B \otimes \mathcal{O}_S$ -modules of type (\mathcal{L}) , in the sense of [RZ], Def. 3.6, 3.10, 3.14.

One key consequence of the conditions imposed in loc. cit., Def. 3.6, is that locally on S there is an isomorphism of polarized multichains of $\mathcal{O}_B \otimes \mathcal{O}_S$ -modules

$$\gamma_\Lambda : M(A_\Lambda) \xrightarrow{\sim} \Lambda \otimes_{\mathbb{Z}_p} \mathcal{O}_S.$$

In fact we have the following result which guarantees this.

THEOREM 6.3 ([**RZ**], Theorems 3.11, 3.16). *Let $\mathcal{L} = \{\Lambda\}$ be a (self-dual) multichain of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices in V . Let S be any \mathbb{Z}_p -scheme where p is locally nilpotent. Then any (polarized) multichain $\{M_\Lambda\}$ of $\mathcal{O}_B \otimes_{\mathbb{Z}_p} \mathcal{O}_S$ -modules of type (\mathcal{L}) is locally (for the étale topology on S) isomorphic to the (polarized) multichain $\{\Lambda \otimes_{\mathbb{Z}_p} \mathcal{O}_S\}$. Moreover, the functor Isom*

$$T \mapsto \text{Isom}(\{M_\Lambda \otimes \mathcal{O}_T\}, \{\Lambda \otimes \mathcal{O}_T\}),$$

is represented by a smooth affine scheme over S .

The analogous statements hold for any \mathbb{Z}_p -scheme S , see [**P**]. In particular for such S we have a smooth affine group scheme \mathcal{G} over S given by

$$\mathcal{G}(T) = \text{Aut}(\{\Lambda \otimes \mathcal{O}_T\}),$$

and the functor Isom is obviously a left-torsor under \mathcal{G} . This generalizes the smoothness of the groups Aut in section 3.2. Moreover, by the same arguments as in section 3.2, for $S = \text{Spec}(\mathbb{Z}_p)$ the group $\mathcal{G}_{\mathbb{Z}_p}$ is a Bruhat-Tits parahoric group scheme corresponding to the parahoric subgroup of $G(\mathbb{Q}_p) = \mathcal{G}(\mathbb{Q}_p)$ which stabilizes the multichain \mathcal{L} ⁷.

In the special case of lattice chains for GSp_{2n} , the theorem was proved by de Jong [**deJ**] (he calls what are “polarized (multi)chains” here by the name “systems of \mathcal{O}_S -modules of type II”).

Now we define the local model diagram for Sh . We assume $\mathcal{O}_E = \mathbb{Z}_p$ for simplicity. Let us define \widetilde{Sh} to be the \mathbb{Z}_p -scheme representing the functor whose points in a \mathbb{Z}_p -scheme S is the set of pairs

$$(\{A_\Lambda\}, \bar{\lambda}, \bar{\eta}) \in Sh(S); \quad \gamma_\Lambda : M(A_\Lambda) \xrightarrow{\sim} \Lambda \otimes_{\mathbb{Z}_p} \mathcal{O}_S,$$

where γ_Λ is an isomorphism of polarized multichains of $\mathcal{O}_B \otimes \mathcal{O}_S$ -modules. The morphism

$$\varphi : \widetilde{Sh} \rightarrow Sh$$

is the obvious morphism which forgets γ_Λ . By Theorem 6.3, φ is smooth (being a torsor for a smooth group scheme) and surjective. Now we want to define

$$\psi : \widetilde{Sh}(S) \rightarrow \mathbf{M}^{\text{loc}}(S).$$

We define it to send an S -point $(\{A_\Lambda\}, \bar{\lambda}, \bar{\eta}, \gamma_\Lambda)$ to the morphism of functors

$$\Lambda \otimes_{\mathbb{Z}_p} \mathcal{O}_S \rightarrow \text{Lie}(A_\Lambda)$$

induced by the composition $\gamma_\Lambda^{-1} : \Lambda \otimes_{\mathbb{Z}_p} \mathcal{O}_S \cong M(A_\Lambda)$ with the canonical surjective morphism

$$M(A_\Lambda) \rightarrow \text{Lie}(A_\Lambda).$$

It is not completely obvious that the morphisms $\Lambda \otimes \mathcal{O}_S \rightarrow \text{Lie}(A_\Lambda)$ satisfy the condition (iii) of Definition 6.2. We will explain it in the Siegel case below, as a

⁷More precisely, the *connected component* of \mathcal{G} is the Bruhat-Tits group scheme. As G. Pappas points out, in some cases (e.g. the unitary group for ramified quadratic extensions), the stabilizer group \mathcal{G} is not connected.

consequence of Proposition 5.1.10 of [BBM] (our Prop. 14.1). We omit discussion of this point in other cases.

The theory of Grothendieck-Messing ([Me]) shows that the morphism ψ is formally smooth. Since both schemes are of finite type over \mathbb{Z}_p , it is smooth. In summary:

THEOREM 6.4 ([RZ], §3). *The diagram*

$$Sh \xleftarrow{\varphi} \widetilde{Sh} \xrightarrow{\psi} \mathbf{M}^{\text{loc}}$$

is a local model diagram. The morphism φ is a torsor for the smooth affine group scheme \mathcal{G} .

PROOF. We have indicated why condition (1) of Definition 6.1 is satisfied. Condition (2) is proved in [RZ], 3.30-3.35; see also [deJ], Cor. 4.6. \square

We will describe the local model diagrams more explicitly for each of our two main examples next. Our goal is to show that their local models are none other than the ones defined in section 4.

6.3.2. *Symplectic case.* Following [deJ] and [GN] we change conventions slightly and replace the de Rham homology functor with the cohomology functor

$$A/S \mapsto H_{DR}^1(A/S).$$

What kind of data do we get by applying the de Rham cohomology functor to a point in our moduli problem Sh from section 5.3? For notational convenience, let us now number the chains of abelian varieties in the opposite order:

$$\{A_{\Lambda_{\bullet}}\} = A_n \rightarrow A_{n-1} \rightarrow \cdots \rightarrow A_0.$$

LEMMA 6.5. *The result of applying H_{DR}^1 to a point $(\{A_{\Lambda_{\bullet}}\}, \lambda_0, \lambda_n)$ in $Sh(S)$ is a datum of form $(M_0 \xrightarrow{\alpha} M_1 \xrightarrow{\alpha} \cdots \xrightarrow{\alpha} M_n, q_0, q_n)$ satisfying*

- M_i is a locally free \mathcal{O}_S -module of rank $2n$;
- $\text{Coker}(M_{i-1} \rightarrow M_i)$ is a locally free $\mathcal{O}_S/p\mathcal{O}_S$ -module of rank 1;
- for $i = 0, n$, $q_i : M_i \otimes M_i \rightarrow \mathcal{O}_S$ is a non-degenerate symplectic pairing;
- for any i , the composition of

$$\begin{array}{ccc} M_0 & \xrightarrow{\alpha} & \cdots \xrightarrow{\alpha} & M_n \\ q_0 \uparrow & & & \downarrow q_n \\ M_0^\vee & \xleftarrow{\alpha^\vee} & \cdots \xleftarrow{\alpha^\vee} & M_n^\vee \end{array}$$

starting and ending at M_i or $M_i^\vee := \text{Hom}(M_i, \mathcal{O}_S)$ is multiplication by p .

PROOF. The pairings q_0, q_n come from the polarizations λ_0, λ_n . The various properties are easy to check, using the canonical natural isomorphism $H_{DR}^1(\widehat{A}/S) = (H_{DR}^1(A/S))^\vee$; cf. Prop. 14.1. \square

Our next goal is to rephrase Definition 6.2 in terms of data similar to that in Lemma 6.5, which will take us closer to the definition of \mathbf{M}^{loc} in §4.

Let $\mathcal{L} = \Lambda_{\bullet}$ be the standard self-dual lattice chain in $V = \mathbb{Q}_p^{2n}$, with respect to the usual pairing $(x, y) = x^t \tilde{I}y$. Clearly we may rephrase Definition 6.2 using the sub-objects $\omega'_\Lambda := \ker(\psi_\Lambda)$ of $\Lambda \otimes \mathcal{O}_S$ rather than the quotients t_Λ . Then condition (iii) becomes

(iii') $(\omega'_\Lambda)^{perp} \subset \omega'_{\Lambda^\perp}$, which is equivalent to $(\omega'_\Lambda)^{perp} = \omega'_{\Lambda^\perp}$,

in other words, under the canonical pairing $(\cdot, \cdot) : \Lambda \otimes \mathcal{O}_S \times \Lambda^\perp \otimes \mathcal{O}_S \rightarrow \mathcal{O}_S$, the submodules ω'_Λ and ω'_{Λ^\perp} are perpendicular. If $\Lambda = \Lambda^\perp$, this means

$$(\cdot, \cdot)|_{\omega'_\Lambda \times \omega'_\Lambda} \equiv 0,$$

and if $\Lambda^\perp = p\Lambda$, this means

$$p(\cdot, \cdot)|_{\omega'_\Lambda \times \omega'_\Lambda} \equiv 0,$$

since the pairing on $\Lambda \otimes \mathcal{O}_S \times \Lambda \otimes \mathcal{O}_S$ is defined by composing the standard pairing on $\Lambda \otimes \mathcal{O}_S \times \Lambda^\perp \otimes \mathcal{O}_S$ with the periodicity isomorphism

$$p : \Lambda \xrightarrow{\sim} \Lambda^\perp$$

in the second variable. For the “standard system” $(\Lambda_0 \rightarrow \Lambda_1 \rightarrow \dots \rightarrow \Lambda_n, q_0, q_n)$ as in Lemma 6.5, the (perfect) pairings are given by

$$q_0 = (\cdot, \cdot) : \Lambda_0 \times \Lambda_0 \rightarrow \mathbb{Z}_p$$

$$q_n = p(\cdot, \cdot) : \Lambda_n \times \Lambda_n \rightarrow \mathbb{Z}_p.$$

Note that if $\omega'_i := \omega'_{\Lambda_i}$, the identity $(\omega'_i)^{perp} = \omega'_{-i}$ (iii) of Def. 6.2) means that ω'_\bullet is uniquely determined by the elements $\omega'_0, \dots, \omega'_n$. Conversely, suppose we are given $\omega'_0, \dots, \omega'_n$ such that $(\omega'_0)^{perp} = \omega'_0$ and $(\omega'_n)^{perp} = p\omega'_n =: \omega'_{-n}$. Then we can define $\omega'_{-i} = (\omega'_i)^{perp}$ for $i = 0, \dots, n$, and then extend by periodicity to get an infinite chain ω'_\bullet as in Definition 6.2 (condition (iii) being satisfied by fiat).

We thus have the following reformulation of Definition 6.2, which shows that that definition agrees with the one in section 4 for GSp_{2n} .

LEMMA 6.6. *In the Siegel case, an S -point of $\mathbf{M}^{\mathrm{loc}}$ (in the sense of Definition 6.2) is a commutative diagram*

$$\begin{array}{ccccccc} \Lambda_0 \otimes \mathcal{O}_S & \longrightarrow & \Lambda_1 \otimes \mathcal{O}_S & \longrightarrow & \dots & \longrightarrow & \Lambda_n \otimes \mathcal{O}_S \\ \uparrow & & \uparrow & & & & \uparrow \\ \omega'_0 & \longrightarrow & \omega'_1 & \longrightarrow & \dots & \longrightarrow & \omega'_n, \end{array}$$

such that

- for each i , ω'_i is a locally free \mathcal{O}_S -submodule of $\Lambda_i \otimes \mathcal{O}_S$ of rank n ;
- ω'_0 is totally isotropic for (\cdot, \cdot) and ω'_n is totally isotropic for $p(\cdot, \cdot)$.

Finally, we promised to explain why the morphism $\psi : \widetilde{Sh} \rightarrow \mathbf{M}^{\mathrm{loc}}$ really takes values in $\mathbf{M}^{\mathrm{loc}}$. We must also redefine it in terms of cohomology. Recall we now have the Hodge filtration

$$\omega_{A_\Lambda/S} \subset H^1_{DR}(A_\Lambda/S).$$

We define ψ to send $(\{A_0 \leftarrow \dots \leftarrow A_n\}, \lambda_0, \lambda_n, \bar{\eta}, \gamma_\Lambda)$ to the locally free, rank n , \mathcal{O}_S -submodules

$$\gamma_\Lambda(\omega_{A_\Lambda}) \subset \Lambda \otimes \mathcal{O}_S,$$

where now γ_Λ is an isomorphism of polarized multichains of \mathcal{O}_S -modules

$$\gamma_\Lambda : H^1_{DR}(A_\Lambda/S) \xrightarrow{\sim} \Lambda \otimes \mathcal{O}_S.$$

The following result ensures that this map really takes values in $\mathbf{M}^{\mathrm{loc}}$.

LEMMA 6.7. *The morphism ψ takes values in $\mathbf{M}^{\mathrm{loc}}$, i.e., condition (iii') holds.*

PROOF. Setting $\omega_{A_{\lambda_i}} = \omega_i$, we need to see that the Hodge filtration ω_0 resp. ω_n is totally isotropic with respect to the pairing q_0 resp. q_n induced by the polarization λ_0 resp. λ_n . But this is Proposition 5.1.10 of [BBM] (our Prop. 14.1). See also [deJ], Cor. 2.2. \square

Comparison of homology and cohomology local models. One further remark is in order. Let us consider a point

$$A = (A_0 \rightarrow \cdots \rightarrow A_n, \lambda_0, \lambda_n, \bar{\eta})$$

in our moduli problem Sh . Note that this data gives us another point in Sh , namely

$$\widehat{A} = (\widehat{A}_n \rightarrow \cdots \rightarrow \widehat{A}_0, \lambda_n^{-1}, \lambda_0^{-1}, \bar{\eta}).$$

(We need to use the assumption that the polarizations λ_i are required to be *symmetric* isogenies $A_i \rightarrow \widehat{A}_i$, in the sense that $\widehat{\lambda}_i = \lambda_i$.)

The moduli problem Sh is thus equipped with an automorphism of order 2, given by $A \mapsto \widehat{A}$.

This comes in handy in comparing the “homology” and “cohomology” constructions of the local model diagram. Namely, since $M(A_i) = H_{DR}^1(\widehat{A}_i)$ (Prop. 14.1), an isomorphism $\gamma_\bullet : M(A_\bullet) \xrightarrow{\sim} \Lambda_\bullet \otimes \mathcal{O}_S$ is simultaneously an isomorphism $\gamma_\bullet : H_{DR}^1(\widehat{A}_\bullet) \xrightarrow{\sim} \Lambda_\bullet \otimes \mathcal{O}_S$. In the “homology” version, ψ sends (A, γ_\bullet) to the quotient chain

$$\Lambda_\bullet \otimes \mathcal{O}_S \rightarrow \text{Lie}(A_\bullet),$$

defined using γ_\bullet^{-1} . On the other hand, in the “cohomology” version, ψ sends $(\widehat{A}, \gamma_\bullet)$ to the sub-object chain

$$\omega_{\widehat{A}_\bullet} \subset \Lambda_\bullet \otimes \mathcal{O}_S$$

(identifying ω_\bullet with $\gamma_\bullet(\omega_\bullet)$). But the exact sequence

$$0 \rightarrow \omega_{\widehat{A}} \rightarrow M(A) \rightarrow \text{Lie}(A) \rightarrow 0$$

(Prop. 14.1) means that the two chains correspond: they give exactly the same element of \mathbf{M}^{loc} . In summary, we have the following result relating the “homology” and “cohomology” constructions of the local model diagram.

PROPOSITION 6.8. *There is a commutative diagram*

$$\begin{array}{ccc} \widetilde{Sh}^{\text{hom}} & \xrightarrow{\psi^{\text{hom}}} & \mathbf{M}^{\text{loc}} \\ \downarrow & & \downarrow = \\ \widetilde{Sh}^{\text{coh}} & \xrightarrow{\psi^{\text{coh}}} & \mathbf{M}^{\text{loc}}, \end{array}$$

where the left vertical arrow is the automorphism $(A, \gamma_\bullet) \mapsto (\widehat{A}, \gamma_\bullet)$.

6.3.3. “Fake” unitary case. Here the “standard” polarized multichain of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices is given by $\{\Lambda_i \oplus \Lambda_i^*\}$, in the notation of section 5.2.3. Recall that

$$\mathcal{O}_B \otimes \mathbb{Z}_p \cong M_n^{\text{opp}}(\mathbb{Z}_p) \times M_n^{\text{opp}}(\mathbb{Z}_p),$$

according to the decomposition of $B_{\mathbb{Q}_p}^{\text{opp}} = D_{\mathbb{Q}_p}$

$$D_{\mathbb{Q}_p} = D_{\mathfrak{p}} \times D_{\mathfrak{p}^*} \cong M_n(\mathbb{Q}_p) \times M_n(\mathbb{Q}_p).$$

Let W (resp. W^*) be \mathbb{Z}_p^n viewed as a left $\mathcal{O}_B \otimes \mathbb{Z}_p$ -module, via right multiplications by elements of the first (resp. second) factor of $M_n(\mathbb{Z}_p) \times M_n(\mathbb{Z}_p)$. The ring $B_{\mathbb{Q}_p}$ has two simple left modules: $W_{\mathbb{Q}_p}$ and $W_{\mathbb{Q}_p}^*$. We may write

$$V_{E'} = V_1 \oplus V_2$$

as before. The determinant condition now implies (at least over E') that

$$V_1 = W_{E'}^d \oplus (W_{E'}^*)^{n-d};$$

(comp. section 5.2.2). Using the “sub-object” variant of Definition 6.2, it follows that an S -point of \mathbf{M}^{loc} is a commutative diagram (here Λ_i being understood as $\Lambda_i \otimes \mathcal{O}_S$)

$$\begin{array}{ccccccc} \Lambda_0 \oplus \Lambda_0^* & \longrightarrow & \Lambda_1 \oplus \Lambda_1^* & \longrightarrow & \cdots & \longrightarrow & \Lambda_n \oplus \Lambda_n^* \\ \uparrow & & \uparrow & & & & \uparrow \\ \mathcal{F}_0 \oplus \mathcal{F}_0^* & \longrightarrow & \mathcal{F}_1 \oplus \mathcal{F}_1^* & \longrightarrow & \cdots & \longrightarrow & \mathcal{F}_n \oplus \mathcal{F}_n^* \end{array}$$

where $\mathcal{F}_i \oplus \mathcal{F}_i^*$ is an $\mathcal{O}_B \otimes \mathcal{O}_S$ -submodule of $\Lambda_i \oplus \Lambda_i^*$ which, locally on S , is a direct factor isomorphic to $W_{\mathcal{O}_S}^{n-d} \oplus (W_{\mathcal{O}_S}^*)^d$.

The analogue of condition (iii'), which is imposed in Definition 6.2, is

$$(\mathcal{F}_i \oplus \mathcal{F}_i^*)^{\text{perp}} = \mathcal{F}_{-i} \oplus \mathcal{F}_{-i}^*.$$

On the other hand, from the definition of $\langle \cdot, \cdot \rangle$ in section 5.2.3 it is immediate that

$$(\mathcal{F}_i \oplus \mathcal{F}_i^*)^{\text{perp}} = \mathcal{F}_i^{*,\text{perp}} \oplus \mathcal{F}_i^{\text{perp}}.$$

We see thus that the first factor \mathcal{F}_\bullet uniquely determines the second factor \mathcal{F}_\bullet^* (and vice-versa). Thus \mathbf{M}^{loc} is given by chains of right $M_n(\mathcal{O}_S) = M_n(\mathbb{Z}_p) \otimes \mathcal{O}_S$ -modules

$$\mathcal{F}_0 \rightarrow \mathcal{F}_1 \rightarrow \cdots \rightarrow \mathcal{F}_n$$

which are locally direct factors in

$$M_n(\mathcal{O}_S) \rightarrow \text{diag}(p^{-1}, 1^{n-1})M_n(\mathcal{O}_S) \rightarrow \cdots \rightarrow p^{-1}M_n(\mathcal{O}_S),$$

each term locally isomorphic to $(\mathcal{O}_S^n)^{n-d}$. By Morita equivalence, \mathbf{M}^{loc} is just given by the definition in section 4 (for the integer d).

7. Flatness

Because of the local model diagram, the flatness of the moduli problem Sh can be investigated by considering its local model. The following fundamental result is due to U. Görtz. It applies to all parahoric subgroups.

THEOREM 7.1 ([Go1], [Go2]). *Suppose \mathbf{M}^{loc} is a local model attached to a group $\text{Res}_{F/\mathbb{Q}_p}(\text{GL}_n)$ or $\text{Res}_{F/\mathbb{Q}_p}(\text{GSp}_{2n})$, where F/\mathbb{Q}_p is an unramified extension. Then \mathbf{M}^{loc} is flat over \mathcal{O}_E . Moreover, its special fiber is reduced, and has rational singularities.*

We give the idea for the proof. One reduces to the case where $F = \mathbb{Q}_p$. In order to prove flatness over $\mathcal{O}_E = \mathbb{Z}_p$ it is enough to prove the following facts (comp. [Ha], III.9.8):

- 1) The special fiber is reduced, as a scheme over \mathbb{F}_p ;
- 2) The model is *topologically flat*: every closed point in the special fiber is contained in the the scheme-theoretic closure of the generic fiber.

The innovation behind the proof of 1) is to embed the special fiber into the affine flag variety and then to make systematic use of the theory of Frobenius-splitting to prove affine Schubert varieties are compatibly Frobenius-split. See [Go1].

To prove 2), suppose μ is such that $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$. It is enough by a result of Kottwitz-Rapoport (Theorem 4.2) to prove that the generic element in a stratum of the special fiber indexed by a translation element in $\text{Adm}(-w_0\mu)$ can be lifted to characteristic zero. This statement is checked by hand in [Go1].

We will provide an alternative, calculation-free, proof by making use of nearby cycles⁸. We freely make use of material on nearby cycles from §10, 11.

We fix an element $\lambda \in W(-w_0\mu)$ and consider the stratum of \mathbf{M}^{loc} indexed by t_λ . We want to show this stratum is in the closure of the generic fiber.

The nearby cycles sheaf $R\Psi^{\mathbf{M}^{\text{loc}}}(\overline{\mathbb{Q}}_\ell)$ is supported only on this closure (Theorem 10.1), and so it is enough to show that

$$\text{Tr}^{ss}(\Phi_p^r, R\Psi_{t_\lambda}^{\mathbf{M}^{\text{loc}}}(\overline{\mathbb{Q}}_\ell)) \neq 0.$$

But it is clear that

$$z_{-w_0\mu, r}(t_\lambda) \neq 0$$

from the definition of Bernstein functions (see [Lu] or [HKP]), and we are done by Theorem 11.3 (which also holds for the group GSp_{2n} , see [HN1]).

As G. Pappas has observed [P], the local models attached (by [RZ]) to the groups above can fail to be flat if F/\mathbb{Q}_p is *ramified*. In their joint works [PR1], [PR2], Pappas and Rapoport provide alternative definitions of local models in that case (in fact they treat nearly all the groups considered in [RZ]), and these new models *are* flat. However, these new models cannot always be described as the scheme representing a “concrete” moduli problem.

8. The Kottwitz-Rapoport stratification

Let us assume Sh is the model over \mathcal{O}_E for one of the Shimura varieties $Sh(\mathbf{G}, h)_{\mathbf{K}}$ discussed in §5, i.e. a “fake” unitary or a Siegel modular variety. We assume (for simplicity of statements) that K_p is an Iwahori subgroup of $G(\mathbb{Q}_p)$. Let us summarize what we know so far.

The group $G_{\mathbb{Q}_p}$ is either isomorphic to $\text{GL}_{n, \mathbb{Q}_p} \times \mathbb{G}_{m, \mathbb{Q}_p}$ or $\text{GSp}_{2n, \mathbb{Q}_p}$. These groups being split over \mathbb{Q}_p , we have $E = \mathbb{Q}_p$ and $\mathcal{O}_E = \mathbb{Z}_p$.

The Shimura datum h gives rise to a dominant minuscule cocharacter μ of $\text{GL}_{n, \mathbb{Q}_p}$ or $\text{GSp}_{2n, \mathbb{Q}_p}$, respectively. The functorial description of the local model \mathbf{M}^{loc} shows that it can be embedded into the deformation M from the affine Grassmannian $\text{Grass}_{\mathbb{Q}_p}$ to the affine flag variety $\mathcal{Fl}_{\mathbb{F}_p}$ associated to G , and has generic fibre $\mathcal{Q}_{-w_0\mu}$. Since the local model is flat with reduced special fiber [Go1], [Go2] (see §7) and is closed in M , it coincides with the scheme-theoretic closure $M_{-w_0\mu}$ of $\mathcal{Q}_{-w_0\mu}$ in this deformation. We thus identify $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$. (For all this, keep in mind we use the “homology” definition of the local model diagram.)

The relation between the model of the Shimura variety over \mathbb{Z}_p and its local model is given by a diagram

$$Sh \xleftarrow{\varphi} \widetilde{Sh} \xrightarrow{\psi} \mathbf{M}^{\text{loc}}$$

⁸We emphasize that this proof is much less elementary than the original proof of Görtz [Go1], relying as it does on the full strength of [HN1].

of \mathbb{Z}_p -schemes, where φ is a torsor under the smooth affine group scheme \mathcal{G} of §6.3 (also termed Aut in §4), and ψ is smooth. The fibres of φ are geometrically connected (more precisely, this holds for the restriction of φ to any geometric connected component of \widetilde{Sh}). One can show that étale-locally around each point of the special fiber of Sh , the schemes Sh and \mathbf{M}^{loc} are isomorphic.

The stratification of the special fibre of \mathbf{M}^{loc} (by Iwahori-orbits) induces stratifications of the special fibers of \widetilde{Sh} and Sh (see below). The resulting stratification of $Sh_{\mathbb{F}_p}$ is called the *Kottwitz-Rapoport (or KR-) stratification*.

8.1. Construction of the KR-stratification. Essentially following [GN], we will recall the construction and basic properties of the KR-stratification. The difference between their treatment and ours is that they construct local models in terms of de Rham *cohomology*, whereas here they are constructed in terms of de Rham *homology*. This is done for compatibility with the computations in §11.

For later use in §11, we give a detailed treatment here for the case of “fake” unitary Shimura varieties.

Let k denote the algebraic closure of the residue field of \mathbb{Z}_p , and let $\widetilde{\Lambda}_\bullet = \Lambda_\bullet \oplus \Lambda_\bullet^*$ denote the self-dual multichain of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices from §5.2.3. Recall that a point in $\mathbf{M}^{\text{loc}}(k)$ is a “quotient chain” of k -vector spaces

$$\widetilde{\Lambda}_\bullet \otimes k \rightarrow t_{\widetilde{\Lambda}_\bullet},$$

self-dual in the sense of Definition 6.2 (iii), and such that each $t_{\widetilde{\Lambda}_i}$ satisfies the determinant condition, that is,

$$t_{\widetilde{\Lambda}_i} = W_k^d \oplus (W^*)_k^{n-d}$$

as $\mathcal{O}_B \otimes k$ -modules. We can identify this object with a lattice chain in the affine flag variety for $\text{GL}_n(k((t)))$ as follows. Let $\mathcal{V}_{\bullet,k}$ denote the “standard” complete lattice chain from §4.1. Using duality and Morita equivalence (see §6.3.3), the quotient $t_{\widetilde{\Lambda}_\bullet}$ can be identified with a quotient t_{Λ_\bullet} of the standard lattice chain $\mathcal{V}_{\bullet,k} \subset k((t))^n$. Then we may write

$$t_{\Lambda_i} = \mathcal{V}_{i,k} / \mathcal{L}_i$$

for a unique lattice chain $\mathcal{L}_\bullet = (\mathcal{L}_0 \subset \cdots \subset \mathcal{L}_n = t^{-1}\mathcal{L}_0)$ consisting of $k[[t]]$ -submodules of $k((t))^n$ which satisfy for each $i = 0, \dots, n$,

- $t\mathcal{V}_{i,k} \subset \mathcal{L}_i \subset \mathcal{V}_{i,k}$;
- the k -vector space $\mathcal{V}_{i,k} / \mathcal{L}_i$ has dimension d (determinant condition).

The set of such lattice chains \mathcal{L}_\bullet is the special fiber of the model $M_{-w_0\mu}$ attached to the dominant coweight $-w_0\mu = (1^d, 0^{n-d})$ of GL_n . Indeed, the two conditions above mean that for each i ,

$$\text{inv}_K(\mathcal{L}_i, \mathcal{V}_{i,k}) = \mu$$

and thus

$$\text{inv}_K(\mathcal{V}_{i,k}, \mathcal{L}_i) = -w_0\mu.$$

Here inv_K is the standard notion of relative position of $k[[t]]$ -lattices in $k((t))^n$, relative to the base point $\mathcal{V}_{0,k} = k[[t]]^n$: we say $\text{inv}_K(g\mathcal{V}_{0,k}, g'\mathcal{V}_{0,k}) = \lambda \in X_+(T)$ if $g^{-1}g' \in K\lambda K$, where $K = \text{GL}_n(k[[t]])$. Recall we have identified μ with $(0^{n-d}, (-1)^d)$ and have embedded coweights into the loop group by the rule $\lambda \mapsto \lambda(t)$.

If $\mathcal{L}_\bullet = x(\mathcal{V}_{\bullet,k})$ for $x \in \widetilde{W}(\text{GL}_n)$, this means that $x \in \text{Perm}(-w_0\mu)$, which is also the set $\text{Adm}(-w_0\mu)$, see §4.

Recall that the Iwahori subgroup $I = I_{k((t))}$ fixing $\mathcal{V}_{\bullet,k}$ preserves the subset $M_{-w_0\mu,k} \subset \mathcal{F}l_k$ and so via the identification $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$, it also acts on the local model. The Iwahori-orbits give a cellular decomposition

$$\mathbf{M}_k^{\text{loc}} = \coprod_{w \in \text{Adm}(\mu)} \mathbf{M}_w^{\text{loc}}.$$

Here we define $\mathbf{M}_w^{\text{loc}}$ to be the set of \mathcal{L}_{\bullet} above such that

$$\text{inv}_I(\mathcal{V}_{\bullet,k}, \mathcal{L}_{\bullet}) = w^{-1},$$

or equivalently

$$\text{inv}_I(\mathcal{L}_{\bullet}, \mathcal{V}_{\bullet,k}) = w,$$

for $w^{-1} \in \text{Adm}(-w_0\mu)$ (which happens if and only if $w \in \text{Adm}(\mu)$). Here we define $\text{inv}_I(g\mathcal{V}_{\bullet,k}, g'\mathcal{V}_{\bullet,k}) = w$ if $g^{-1}g' \in IwI$.

Each stratum is smooth (in fact $\mathbf{M}_w^{\text{loc}} = \mathbb{A}^{\ell(w)}$), and the closure relations are determined by the Bruhat order on \widetilde{W} ; that is, $\mathbf{M}_w^{\text{loc}} \subset \overline{\mathbf{M}_{w'}^{\text{loc}}}$ if and only if $w \leq w'$.

There is a surjective homomorphism $I_{k((t))} \rightarrow \text{Aut}_k$, where Aut is the group scheme \mathcal{G} of Theorem 6.3 which acts on the whole local model diagram. The action of $I_{k((t))}$ on $\mathbf{M}_k^{\text{loc}}$ factors through Aut_k , so that the strata above can also be thought of as Aut_k -orbits. The morphism ψ is clearly equivariant for Aut_k , hence we have a stratification of \widetilde{Sh}_k

$$\widetilde{Sh}_k = \coprod_{w \in \text{Adm}(\mu)} \psi^{-1}(\mathbf{M}_w^{\text{loc}}),$$

whose strata are non-empty (Lemma 13.1), smooth, and stable under the action of Aut_k . Since φ_k is a torsor for the smooth group scheme Aut_k , the stratification descends to Sh_k :

$$Sh_k = \coprod_{w \in \text{Adm}(\mu)} Sh_w$$

such that $\varphi^{-1}(Sh_w) = \psi^{-1}(\mathbf{M}_w^{\text{loc}})$. These strata are still smooth, non-empty, and satisfy closure relations determined by the Bruhat order.

All statements above remain true over the base field \mathbb{F}_p instead of its algebraic closure k .

8.2. Relating nearby cycles on local models and Shimura varieties.

We will need in §11 the following result relating the nearby cycles on Sh , \widetilde{Sh} , and \mathbf{M}^{loc} , which follows immediately from the above remarks and Theorem 10.1 below (cf. [GN]):

LEMMA 8.1. *There are canonical isomorphisms*

$$\varphi^* R\Psi^{Sh}(\overline{\mathbb{Q}}_{\ell}) = R\Psi^{\widetilde{Sh}}(\overline{\mathbb{Q}}_{\ell}) = \psi^* R\Psi^{\mathbf{M}^{\text{loc}}}(\overline{\mathbb{Q}}_{\ell}).$$

Moreover, $R\Psi^{Sh}(\overline{\mathbb{Q}}_{\ell})$ is constant on each stratum Sh_w , and if $\Phi_{\mathfrak{p}} \in \text{Gal}(\overline{\mathbb{Q}}_p/E)$ is a geometric Frobenius element, then for any elements $x \in Sh_w(k_r)$ and $x_0 \in \mathbf{M}_w^{\text{loc}}(k_r)$, we have

$$\text{Tr}^{ss}(\Phi_{\mathfrak{p}}^r, R\Psi_x^{Sh}(\overline{\mathbb{Q}}_{\ell})) = \text{Tr}^{ss}(\Phi_{\mathfrak{p}}^r, R\Psi_{x_0}^{\mathbf{M}^{\text{loc}}}(\overline{\mathbb{Q}}_{\ell})).$$

Here we use the notion of semi-simple trace, which is explained below in §9.3. The above lemma plays a key role in the determination of the semi-simple local zeta function in §11.

8.3. The Genestier-Ngô comparison with p -rank. We assume in this section that Sh is the Siegel modular variety with Iwahori-level structure from §5.3.

Recall that any n -dimensional abelian variety A over an algebraically closed field k of characteristic p has

$$\#A[p](k) = p^j,$$

for some integer $0 \leq j \leq n$. The integer j is called the p -rank of A . Ordinary abelian varieties are those whose p -rank is n , the largest possible. The p -rank is constant on isogeny classes, and therefore it determines a well-defined function on the set of geometric points $Sh_{\mathbb{F}_p}$. The level sets determine the *stratification by p -rank*.

It is natural to ask how this stratification relates to the KR-stratification. In [GN], Genestier and Ngô have very elegantly derived the relationship using local models and work of de Jong [deJ]. As they point out, their theorem yields interesting results even in the case of Siegel modular varieties having *good reduction* at p : they derive a short and beautiful proof that the ordinary locus in such Shimura varieties is open and dense in the special fiber (comp. [W1]).

To state their result, we define for $w \in \widetilde{W}(\mathrm{GSp}_{2n})$ an integer $r(w)$ as follows. Its image \bar{w} in the finite Weyl group $W(\mathrm{GSp}_{2n})$ is a permutation of the set $\{1, \dots, 2n\}$ commuting with the involution $i \mapsto 2n+1-i$. The set of fixed points of \bar{w} is stable under the involution, and therefore has even cardinality (the involution is without fixed-points). Define

$$2r(w) = \#\{\text{fixed points of } \bar{w}\}.$$

THEOREM 8.2 (Genestier-Ngô [GN]). *The p -rank is constant on each KR-stratum Sh_w . More precisely, the p -rank of a point in Sh_w is the integer $r(w)$.*

COROLLARY 8.3 ([GN]). *The ordinary locus in $Sh_{\mathbb{F}_p}$ is precisely the union of the KR-strata indexed by the translation elements in $\mathrm{Adm}(\mu)$, that is, the elements t_λ , for $\lambda \in W\mu$. Moreover, the ordinary locus is dense and open in $Sh_{\mathbb{F}_p}$.*

PROOF. By the theorem, the p -rank is n on Sh_w if and only if $r(w) = n$; writing $w = t_\lambda \bar{w}$, this is equivalent to $\bar{w} = 1$. We conclude the first statement by noting that $w \in \mathrm{Adm}(\mu) \Rightarrow \lambda \in W\mu$.

Finally, the union of the strata $\mathbf{M}_{t_\lambda}^{\mathrm{loc}}$ for $\lambda \in W\mu$ is clearly dense and open in $\mathbf{M}_{\mathbb{F}_p}^{\mathrm{loc}}$. \square

REMARK 8.4. It should be noted that in [GN] the local model, and thus the KR-stratification, is defined in terms of the “cohomology” local model diagram, whereas here everything is stated using the “homology” version. Furthermore, in [GN] the “standard” lattice chain is “opposite” from ours, so that an element $w \in \widetilde{W}(\mathrm{GSp}_{2n})$ is used to index a double coset $\bar{I}w\bar{I}/\bar{I}$, where \bar{I} is an “opposite” Iwahori subgroup. Nevertheless, our conventions and those of [GN] yield *the same answer*, that is, the p -rank on Sh_w is given by $r(w)$ in both cases. This may be seen by using the comparison between “homology” and “cohomology” local model diagrams in Prop. 6.8, and by imitating the proof of [GN] with our conventions in force.

8.4. The smooth locus of $Sh_{\mathbb{F}_p}$. Also, in [GN] one finds the proof of the following related fact.

PROPOSITION 8.5 (Genestier-Ngô [GN]). *The smooth locus of $Sh_{\mathbb{F}_p}$ is the union of the KR-strata indexed by t_λ , for $\lambda \in W\mu$ (in particular the smooth locus agrees with the ordinary locus).*

8.4.1. *The geometric proof of [GN].* The crucial observation is that any stratum Sh_w , where w is not a translation element, is contained in the singular locus. Genestier and Ngô deduce this by showing that for such w ,

$$(8.4.1) \quad \text{Tr}^{ss}(\Phi_p^r, R\Psi_w^{Sh}(\overline{\mathbb{Q}}_\ell)) \neq 1,$$

which by the general geometric principle explained below, shows that w is singular. Now (8.4.1) itself is proved by combining the main theorems of [HN1] and [H2], and by taking into account Lemma 8.1.

Here is the geometric principle implicit in [GN] and a sketch of the proof from [GN]. We will use freely the material from sections 9.3 and 10 below.

LEMMA 8.6. *Let $S = (S, s, \eta)$ be a trait, with $k(s) = \mathbb{F}_q$ a finite field. Suppose $M \rightarrow S$ is a finite type flat model with M_η smooth. Then $x \in M(\mathbb{F}_{q^r})$ is a smooth point of $M_{\bar{s}}$ only if $\text{Tr}^{ss}(\Phi_q^r, R\Psi_x^M(\overline{\mathbb{Q}}_\ell)) = 1$.*

PROOF. Let $M' \subset M$ be the open subscheme obtained by removing the singular locus of the special fiber of M . We see that $M' \rightarrow S$ is smooth (since $M' \rightarrow S$ is flat of finite-type, it suffices to check the smoothness fiber by fiber, and by construction $M'_\eta = M_\eta$ and M'_s are both smooth). Now we invoke the general fact (Theorem 10.1) that for smooth models M' , $R\Psi^{M'}(\overline{\mathbb{Q}}_\ell) \cong \overline{\mathbb{Q}}_\ell$, the constant sheaf on the special fiber. This implies that the semi-simple trace of nearby cycles at $x \in M'_s$ is 1. \square

Proof of Proposition 8.5. We consider the stratum Sh_w , or equivalently, $\mathbf{M}_w^{\text{loc}}$, for $w \in \text{Adm}(\mu)$. We recall that $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$ and the stratum $\mathbf{M}_w^{\text{loc}}$ is the Iwahori-orbit indexed by $x := w^{-1}$, contained in $M_{-w_0\mu}$. For such an x , we have from [HN1], [H2], [HP] an explicit formula for the semi-simple trace of Frobenius on nearby cycles at x

$$\text{Tr}^{ss}(\Phi_q, R\Psi_x^{\mathbf{M}^{\text{loc}}}(\overline{\mathbb{Q}}_\ell)) = (-1)^{\ell(t_\mu) + \ell(x)} R_{x, t_{\lambda(x)}}(q),$$

where $\lambda(x)$ is the translation part of x ($x = t_{\lambda(x)}w$, for $\lambda(x) \in X_*$, and $w \in W_0$), and $R_{x,y}(q)$ is the Kazhdan-Lusztig R -polynomial. This polynomial can be computed explicitly, but we need only the fact that it is always a polynomial in q of degree $\ell(y) - \ell(x)$. It follows from this and the above lemma that whenever x corresponds to a stratum of codimension ≥ 1 , every point of that stratum is singular.

8.4.2. *A combinatorial proof.* There is however a more elementary way to proceed: we prove below that every $\text{codim} \geq 1$ stratum in $\mathbf{M}_{\mathbb{F}_p}^{\text{loc}}$ is contained at least two irreducible components. The same goes for $Sh_{\mathbb{F}_p}$, proving the proposition. (There is even a third proof of the proposition, given in [GH].)

PROPOSITION 8.7. *Let μ be minuscule. For any $x \in \text{Adm}(\mu)$ of codimension 1, there exist exactly two distinct translation elements λ_1, λ_2 in $W\mu$ such that $x \leq t_{\lambda_i}$ (for $i = 1, 2$). Thus, any codimension 1 KR-stratum in the special fiber of a Shimura variety Sh with Iwahori-level structure at p is contained in exactly two irreducible components.*

PROOF. We give a purely combinatorial proof. Suppose $x \in \text{Adm}(\mu)$ has codimension 1. We have $x < t_\nu$, for some $\nu \in W\mu$. By properties of the Bruhat

order, there exists an affine reflection $s_{\beta+k}$, where β is a B -positive root, such that $x = t_\nu s_{\beta+k}$. Since $s_{\beta+k} = t_{-k\beta^\vee} s_\beta$, this means $x = t_{\nu-k\beta^\vee} s_\beta$. The translation part must lie in $\text{Perm}(\mu) \cap X_* = W\mu$, hence we have $x = t_\lambda s_\beta$, for $\lambda \in W\mu$. By comparing lengths, we have $t_\lambda s_\beta < t_\lambda$ and $t_\lambda s_\beta < s_\beta t_\lambda s_\beta = t_{s_\beta \lambda}$. We claim that $\langle \beta, \lambda \rangle < 0$. Indeed, writing $\epsilon_\beta \in [-1, 0)$ for the infimum of the set $\beta(\mathbf{a})$ (recalling that our base alcove \mathbf{a} is contained in the \bar{B} -positive chamber), we have

$$\begin{aligned} t_\lambda s_\beta < t_\lambda &\Leftrightarrow \mathbf{a} \text{ and } t_{-\lambda} \mathbf{a} \text{ are on opposite sides of the hyperplane } \beta = 0 \\ &\Leftrightarrow \beta(-\lambda + \mathbf{a}) \in [0, \infty) \\ &\Leftrightarrow -\langle \beta, \lambda \rangle + (\epsilon_\beta, 0) \in [0, \infty) \\ &\Leftrightarrow \langle \beta, \lambda \rangle < 0. \end{aligned}$$

We see that $s_\beta \lambda \neq \lambda$, and so x precedes at least the two distinct translation elements t_λ and $t_{s_\beta \lambda}$ in $\text{Adm}(\mu)$. It remains to prove that these are the only such translation elements. So suppose now that $t_\lambda s_\beta < t_{\lambda'}$, where $\lambda' \in W\mu$; we will show that $\lambda' \in \{\lambda, s_\beta \lambda\}$. As above, there is an affine reflection $s_{\alpha+n}$ such that $t_\lambda s_\beta = t_{\lambda' s_{\alpha+n}} = t_{\lambda' - n\alpha^\vee} s_\alpha$, where α is B -positive. We see that $\alpha = \beta$, and $\lambda' - n\beta^\vee = \lambda$. Thus, λ' , λ , and $s_\beta \lambda$ all lie on the line $\lambda + \mathbb{R}\beta^\vee$. Since all elements in $W\mu$ are vectors with the same Euclidean length, this can only occur if $\lambda' \in \{\lambda, s_\beta \lambda\}$. \square

9. Langlands' strategy for computing local L -factors

The well-known general strategy for computing the local L -factor at p of a Shimura variety in terms of automorphic L -functions is due to the efforts of many people, beginning with Eichler, Shimura, Kuga, Sato, and Ihara, and reaching its final conjectural form with Langlands, Rapoport, and Kottwitz.

Let us fix a rational prime p , and a compact open subgroup $K_p \subset G(\mathbb{Q}_p)$ at p ; we consider the Shimura variety $Sh(\mathbf{G}, h)_{\mathbf{K}}$ as in §5.1.

Roughly, the method of Langlands is to start with a cohomological definition of the local factor of the Hasse-Weil zeta function for $Sh(\mathbf{G}, h)_{\mathbf{K}}$, and express its logarithm via the Grothendieck-Lefschetz trace formula as a certain sum of orbital integrals for the group $\mathbf{G}(\mathbb{A})$. This involves both a process of counting points (with ‘‘multiplicity’’ – the trace of the correspondence on the stalk of an appropriate sheaf), and then a ‘‘pseudo-stabilization’’ like that done to stabilize the geometric side of the Arthur-Selberg trace formula (we are ignoring the appearance of endoscopic groups other than \mathbf{G} itself in this stage). At this point, we can apply the Arthur-Selberg stable trace formula and express the sum as a trace of a function on automorphic representations appearing in the discrete part of $L^2(\mathbf{G}(\mathbb{Q}) \backslash \mathbf{G}(\mathbb{A}))$. This equality of traces implies a relation like that in Theorem 11.7 below.

More details on the general strategy as well as the precise conjectural description of $IH^i(Sh \times_E \overline{\mathbb{Q}}_p, \overline{\mathbb{Q}}_\ell)$ in terms of automorphic representations of \mathbf{G} can be found in [Ko90], [Ko92b], and [BR]. These sources discuss the case of good reduction at \mathfrak{p} .

Some details of the analogous strategy in case of bad reduction will be given below in §11. Let us explain more carefully the relevant definitions.

9.1. Definition of local factors of the Hasse-Weil Zeta function. Let \mathfrak{p} denote a prime of a number field \mathbf{E} , lying over p . Let X be a smooth d -dimensional

variety over \mathbf{E} . We define the Hasse-Weil zeta function (of a complex variable s) as an Euler product

$$Z(s, X) = \prod_{\mathfrak{p}} Z_{\mathfrak{p}}(s, X),$$

where the local factors are defined as follows.

DEFINITION 9.1. The factor $Z_{\mathfrak{p}}(s, X)$ is defined to be

$$\prod_{i=0}^{2d} \det(1 - N\mathfrak{p}^{-s}\Phi_{\mathfrak{p}}; H_c^i(X \times_E \overline{\mathbb{Q}}_{\mathfrak{p}}, \overline{\mathbb{Q}}_{\ell})^{\Gamma_{\mathfrak{p}}^0})^{(-1)^{i+1}},$$

where

- ℓ is an auxiliary prime, $\ell \neq p$;
- $\Phi_{\mathfrak{p}}$ is the inverse of an arithmetic Frobenius element for the extension $\mathbf{E}_{\mathfrak{p}}/\mathbb{Q}_p$;
- $N\mathfrak{p} = \text{Norm}_{\mathbf{E}_{\mathfrak{p}}/\mathbb{Q}_p} \mathfrak{p}$;
- $\Gamma_{\mathfrak{p}}^0 \subset \Gamma_{\mathfrak{p}} := \text{Gal}(\overline{\mathbb{Q}}_{\mathfrak{p}}/\mathbf{E}_{\mathfrak{p}})$ is the inertia subgroup.

It is believed that $Z(s, X)$ has good analytical properties (it should satisfy a functional equation and have a meromorphic analytic continuation on \mathbb{C}) and that analytical invariants (e.g. residues, special values, orders of zeros and poles) carry important arithmetic information about X . Moreover, it is believed that the local factor is indeed independent of the choice of the auxiliary prime ℓ . At present these remain only guiding principles, as very little has been actually proved. As Taniyama originally proposed, a promising strategy for establishing the functional equation is to express the zeta function as a product of automorphic L -functions, whose analytic properties are easier to approach. The hope that this can be done is at the heart of the Langlands program.

9.2. Definition of local factors of automorphic L -functions. Let π_p be an irreducible admissible representation of $G(\mathbb{Q}_p)$. Let us assume that the local Langlands conjecture holds for the group $G(\mathbb{Q}_p)$. Then associated to π_p is a local Langlands parameter, that is, a homomorphism

$$\varphi_{\pi_p} : W_{\mathbb{Q}_p} \times \text{SL}_2(\mathbb{C}) \rightarrow {}^L G,$$

where $W_{\mathbb{Q}_p}$ is the Weil group for \mathbb{Q}_p and, letting \widehat{G} denote the Langlands dual group over \mathbb{C} associated to $G_{\mathbb{Q}_p}$, the L -group is defined to be ${}^L G = W_{\mathbb{Q}_p} \rtimes \widehat{G}$. Let $r = (r, V)$ be a rational representation of ${}^L G$. The local L -function attached to π_p is defined using φ_{π_p} and r as follows.

DEFINITION 9.2. We define $L(s, \pi_p, r)$ to be

$$\det(1 - p^{-s} r \varphi_{\pi_p}(\Phi_p \times \begin{bmatrix} p^{-1/2} & 0 \\ 0 & p^{1/2} \end{bmatrix}); (\ker N)^{\Gamma_p^0})^{-1},$$

where

- $\Phi_p \in W_{\mathbb{Q}_p}$ is the inverse of the arithmetic Frobenius for \mathbb{Q}_p ;
- N is a nilpotent endomorphism on V coming from the action of \mathfrak{sl}_2 on the representation $r\varphi_{\pi_p}$, namely $N := d(r\varphi_{\pi_p})(1 \times \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix})$; equivalently, N is determined by $r\varphi_{\pi_p}(1 \times \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}) = \exp(N)$;

- the action of inertia $\Gamma_p^0 \subset W_{\mathbb{Q}_p}$ on $\ker(N)$ is the restriction of $r\varphi_{\pi_p}$ to $\Gamma_p^0 \times \text{id} \subset W_{\mathbb{Q}_p} \times \text{SL}_2(\mathbb{C})$.

REMARK 9.3. If π_p is a spherical representation, then $(\ker N)^{\Gamma_p^0} = V$ and therefore in that case the local factor takes the more familiar form

$$\det(1 - p^{-s}r(\text{Sat}(\pi_p)); V)^{-1},$$

where $\text{Sat}(\pi_p)$ is the Satake parameter of π_p ; see [Bo], [Ca].

Any irreducible admissible representation π of $\mathbf{G}(\mathbb{A})$ has a tensor factorization $\pi = \otimes_v \pi_v$ (v ranges over all places of \mathbb{Q}) where π_v is an admissible representation of the local group $\mathbf{G}(\mathbb{Q}_v)$. If $v = \infty$, there is a suitable definition of $L(s, \pi_\infty, r)$ (see [Ta]). We then define the automorphic L -function

$$L(s, \pi, r) = \prod_v L(s, \pi_v, r).$$

9.3. Problems in case of bad reduction, and definition of semi-simple local factors.

9.3.1. *Semi-simple zeta function.* Let us fix \mathfrak{p} and set $E = \mathbf{E}_{\mathfrak{p}}$. In the case where X possesses an integral model over \mathcal{O}_E , one can study the local factor by reduction modulo \mathfrak{p} . In the case of good reduction (meaning this model is smooth over \mathcal{O}_E), the inertia group acts trivially and the cohomology of $X \times_E \overline{\mathbb{Q}}_p$ can be identified with that of the special fiber $X \times_{k_E} \overline{\mathbb{F}}_p$ (k_E being the residue field of E). By the Grothendieck-Lefschetz trace formula, the local zeta function then satisfies the familiar identity

$$\log(Z_{\mathfrak{p}}(s, X)) = \sum_{r=1}^{\infty} \#X(k_{E,r}) \frac{N\mathfrak{p}^{-rs}}{r},$$

where $k_{E,r}$ is the degree r extension of k_E in its algebraic closure k . In the case of bad reduction, inertia can act non-trivially and the smooth base-change theorems of ℓ -adic cohomology no longer apply in such a simple way. Following Rapoport [R1], we bypass the first difficulty by working with the *semi-simple* local zeta function, defined below. The second difficulty forces us to work with nearby cycles (see §10): if X is a proper scheme over \mathcal{O}_E , then there is a $\Gamma_{\mathfrak{p}}$ -equivariant isomorphism

$$H^i(X \times_E \overline{\mathbb{Q}}_p, \overline{\mathbb{Q}}_{\ell}) = H^i(X \times_{k_E} \overline{k}_E, R\Psi(\overline{\mathbb{Q}}_{\ell})),$$

so that the Grothendieck-Lefschetz trace formula gives rise to the problem of “counting counts $x \in X(k_{E,r})$ with multiplicity”, i.e., to computing the semi-simple trace on the stalks of the complex of nearby cycles:

$$\text{Tr}^{ss}(\Phi_{\mathfrak{p}}^r, R\Psi(\overline{\mathbb{Q}}_{\ell})_x),$$

in order to understand the semi-simple zeta function.

How do we define semi-simple trace and semi-simple zeta functions? We recall the discussion from [HN1]. Suppose V is a finite-dimensional continuous ℓ -adic representation of the Galois group $\Gamma_{\mathfrak{p}}$. Grothendieck’s local monodromy theorem states that this representation is necessarily *quasi-unipotent*: there exists a finite index subgroup $\Gamma_1 \subset \Gamma_{\mathfrak{p}}$ such that Γ_1 acts unipotently on V . Hence there is a finite increasing $\Gamma_{\mathfrak{p}}$ -invariant filtration $V_{\bullet} = (0 \subset V_1 \subset \dots \subset V_m = V)$ such that $\Gamma_{\mathfrak{p}}$ acts

on $\oplus_k gr_k V_\bullet$ through a finite quotient. Such a filtration is called *admissible*. Then we define

$$\mathrm{Tr}^{ss}(\Phi_p; V) = \sum_k \mathrm{Tr}(\Phi_p; gr_k(V_\bullet)^{\Gamma_p^0}).$$

This is independent of the choice of admissible filtration. Moreover, the function $V \mapsto \mathrm{Tr}^{ss}(\Phi_p, V)$ factors through the Grothendieck group of the category of ℓ -adic representations V , and using this one proves that it extends naturally to give a “sheaf-function dictionary” à la Grothendieck: a complex \mathcal{F} in the “derived” category $D_c^b(X \times_\eta s, \overline{\mathbb{Q}}_\ell)$,⁹ gives rise to the $\overline{\mathbb{Q}}_\ell$ -valued function

$$x \mapsto \mathrm{Tr}^{ss}(\Phi_p^r, \mathcal{F}_x)$$

on $X(k_{E,r})$. Furthermore, the formation of this function is compatible with the pull-back and proper-push-forward operations on the derived category, and a Grothendieck-Leftschetz trace formula holds. (For details, see [HN1].) We can then *define* the semi-simple local zeta function $Z_p^{ss}(s, X)$ by the identity

DEFINITION 9.4.

$$\log(Z_p^{ss}(s, X)) = \sum_{r=1}^{\infty} \left(\sum_i (-1)^i \mathrm{Tr}^{ss}(\Phi_p^r; H_c^i(X \times_E \overline{\mathbb{Q}}_p, \overline{\mathbb{Q}}_\ell)) \right) \frac{Np^{-rs}}{r}.$$

REMARK 9.5. Note that in the case where Γ_p^0 acts unipotently (not just quasi-unipotently) on the cohomology of the generic fiber, then we have

$$\mathrm{Tr}^{ss}(\Phi_p^r; H_c^i(X \times_E \overline{\mathbb{Q}}_p, \overline{\mathbb{Q}}_\ell)) = \mathrm{Tr}(\Phi_p^r; H_c^i(X \times_E \overline{\mathbb{Q}}_p, \overline{\mathbb{Q}}_\ell)).$$

As we shall see below in §10.2, Γ_p^0 does indeed act unipotently on the cohomology of a proper Shimura variety with Iwahori-level structure at p . The definition of Z^{ss} thus simplifies in that case.

As before, the global semi-simple zeta function is defined to be an Euler product over all finite places of the local functions: $Z^{ss}(s, X) = \prod_p Z_p^{ss}(s, X)$.

9.3.2. *Semi-simple local L-functions.* Retain the notation of §9.2. The Langlands parameters $\varphi = \varphi_{\pi_p}$ being used here have the property that the representation $r\varphi$ on V arises from a representation (ρ, N) of the Weil-Deligne group $W'_{\mathbb{Q}_p}$ on V , see [Ta]. In that case we have for $w \in W_{\mathbb{Q}_p}$

$$\begin{aligned} \rho(w) &= r\varphi(w \times \begin{bmatrix} \|w\|^{1/2} & 0 \\ 0 & \|w\|^{-1/2} \end{bmatrix}) \\ \exp(N) &= r\varphi(1 \times \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}), \end{aligned}$$

where $\|w\|$ is the power to which w raises elements in the residue field of \mathbb{Q}_p .

Now suppose that via some choice of isomorphism $\overline{\mathbb{Q}}_\ell \cong \mathbb{C}$, the pair (ρ, N) comes from an ℓ -adic representation $(\rho_\lambda, V_\lambda)$ of $W_{\mathbb{Q}_p}$, by the rule

$$\rho_\lambda(\Phi^n \sigma) = \rho(\Phi^n \sigma) \exp(N t_\ell(\sigma)),$$

where $n \in \mathbb{Z}$, $\sigma \in \Gamma_p^0$, and $t_\ell : \Gamma_p^0 \rightarrow \mathbb{Q}_\ell$ is a nonzero homomorphism (cf. [Ta], Thm 4.2.1).

⁹This is the “derived category of $\overline{\mathbb{Q}}_\ell$ -sheaves” – although this is somewhat misleading terminology (see [KW] for a detailed account) – on $X \times_{k_E} \overline{k_E}$ equipped with a continuous action of Γ_p compatible with the action of its quotient $\mathrm{Gal}(\overline{k_E}/k_E)$ on $X \times \overline{k_E}$. See §10.

For each $\sigma \in \Gamma_p^0$, $\rho_\lambda(\sigma) = \rho(\sigma)\exp(Nt_\ell(\sigma))$ is the multiplicative Jordan decomposition of $\rho_\lambda(\sigma)$ into its semi-simple and unipotent parts, respectively. Therefore a vector $v \in V$ is fixed by $\rho_\lambda(\sigma)$ if and only if it is fixed by both $\rho(\sigma)$ and $\exp(Nt_\ell(\sigma))$. Thus we have the following result.

LEMMA 9.6. *If (ρ, N, V) and $(\rho_\lambda, V_\lambda)$ are the two avatars above of a representation of the Weil-Deligne group $W_{\mathbb{Q}_p}^1$, then*

$$(\ker N)^{\Gamma_p^0} = V_\lambda^{\Gamma_p^0}.$$

Furthermore ρ is trivial on Γ_p^0 if and only if $\rho_\lambda(\Gamma_p^0)$ acts unipotently on V_λ , and in that case

$$(\ker N)^{\Gamma_p^0} = \ker N = V_\lambda^{\Gamma_p^0}.$$

COROLLARY 9.7. *The local L -function can also be expressed as*

$$L(s, \pi_p, r) = \det(1 - p^{-s}\rho_\lambda(\Phi_p); V_\lambda^{\Gamma_p^0})^{-1}.$$

Note the similarity with Definition 9.1. The representation $\rho_\lambda(\Gamma_p^0)$ being quasi-unipotent, we can define the semi-simple L -function as in Definition 9.4.

DEFINITION 9.8.

$$\log(L^{ss}(s, \pi_p, r)) = \sum_{r=1}^{\infty} \text{Tr}^{ss}(\rho_\lambda(\Phi_p^r); V_\lambda) \frac{p^{-rs}}{r}.$$

(The symbol r occurs with two different meanings here, but this should not cause confusion.)

REMARK 9.9. In analogy with Remark 9.5, in the case that (ρ, N, V) has $\rho(\Gamma_p^0) = 1$, in view of Lemma 9.6, we have

$$\text{Tr}^{ss}(\rho_\lambda(\Phi_p^r); V) = \text{Tr}(\rho_\lambda(\Phi_p^r); V),$$

and the definition of L^{ss} simplifies. The dictionary set-up by the local Langlands correspondence asserts that if π_p has an Iwahori-fixed vector, then $\rho(\Gamma_p^0)$ is trivial (cf. [W2]). Moreover, in the situation of parahoric level structure at p , the only representations π_p which will arise necessarily have Iwahori-fixed vectors. Hence the simplified definition of L^{ss} will apply in our situation.

Let $\pi = \otimes_v \pi_v$ be an irreducible admissible representation of $\mathbf{G}(\mathbb{A})$. It is to be hoped that a reasonable definition of $L^{ss}(s, \pi_v, r)$ exists for Archimedean places v , and if so we can then define

$$L^{ss}(s, \pi, r) = \prod_v L^{ss}(s, \pi_v, r).$$

10. Nearby cycles

10.1. Definitions and general facts. Let X be a scheme of finite type over a finite (or algebraically closed) field k . (The following also works if we assume that k is the fraction field of a discrete valuation ring R with finite residue field, and that X is finite-type over R , cf. [Ma].) Denote by \bar{k} an algebraic closure of k , and by $X_{\bar{k}}$ the base change $X \times_k \bar{k}$.

We denote by $D_c^b(X, \overline{\mathbb{Q}}_\ell)$ the 'derived' category of $\overline{\mathbb{Q}}_\ell$ -sheaves on X . Note that this is not actually the derived category of the category of $\overline{\mathbb{Q}}_\ell$ -sheaves, but is

defined via a limit process. See [BBD] 2.2.14 or [Weil2] 1.1.2, or [KW] for more details. Nevertheless, $D_c^b(X, \overline{\mathbb{Q}}_\ell)$ is a triangulated category which admits the usual functorial formalism, and which can be equipped with a 'natural' t -structure having as its core the category of $\overline{\mathbb{Q}}_\ell$ -sheaves. If $f : X \rightarrow Y$ is a morphism of schemes of finite type over k , we have the derived functors f_* , $f^! : D_c^b(X, \overline{\mathbb{Q}}_\ell) \rightarrow D_c^b(Y, \overline{\mathbb{Q}}_\ell)$ and f^* , $f^! : D_c^b(Y, \overline{\mathbb{Q}}_\ell) \rightarrow D_c^b(X, \overline{\mathbb{Q}}_\ell)$. Occasionally we denote these same functors using the symbols Rf_* , etc.

Let (S, s, η) denote an Henselian trait: S is the spectrum of a complete discrete valuation ring, with special point s and generic point η . The key examples for us are $S = \text{Spec}(\mathbb{Z}_p)$ (the p -adic setting) and $S = \text{Spec}(\mathbb{F}_p[[t]])$ (the function-field setting). Let $k(s)$ resp. $k(\eta)$ denote the residue fields of s resp. η .

We choose a separable closure $\bar{\eta}$ of η and define the Galois group $\Gamma = \text{Gal}(\bar{\eta}/\eta)$ and the inertia subgroup $\Gamma_0 = \ker[\text{Gal}(\bar{\eta}/\eta) \rightarrow \text{Gal}(\bar{s}/s)]$, where \bar{s} is the residue field of the normalization \bar{S} of S in $\bar{\eta}$.

Now let X denote a finite-type scheme over S . The category $D_c^b(X \times_s \eta, \overline{\mathbb{Q}}_\ell)$ is the category of sheaves $\mathcal{F} \in D_c^b(X_{\bar{s}}, \overline{\mathbb{Q}}_\ell)$ together with a continuous action of $\text{Gal}(\bar{\eta}/\eta)$ which is compatible with the action on $X_{\bar{s}}$. (Continuity is tested on cohomology sheaves.)

For $\mathcal{F} \in D_c^b(X_\eta, \overline{\mathbb{Q}}_\ell)$, we define the *nearby cycles sheaf* to be the object in $D_c^b(X \times_s \eta, \overline{\mathbb{Q}}_\ell)$ given by

$$R\Psi^X(\mathcal{F}) = \bar{i}^* R\bar{j}_*(\mathcal{F}_{\bar{\eta}}),$$

where $\bar{i} : X_{\bar{s}} \hookrightarrow X_{\bar{S}}$ and $\bar{j} : X_{\bar{\eta}} \hookrightarrow X_{\bar{S}}$ are the closed and open immersions of the geometric special and generic fibers of X/S , and $\mathcal{F}_{\bar{\eta}}$ is the pull-back of \mathcal{F} to $X_{\bar{\eta}}$.

Here we list the basic properties of $R\Psi^X$, extracted from the standard references: [BBD], [II], [SGA7 I], [SGA7 XIII]. The final listed property has been proved (in this generality) only recently, and is due to the author and U. Görtz [GH].

THEOREM 10.1. *The following properties hold for the functors*

$$R\Psi : D_c^b(X_\eta, \overline{\mathbb{Q}}_\ell) \rightarrow D_c^b(X \times_s \eta, \overline{\mathbb{Q}}_\ell) :$$

- (a) *$R\Psi$ commutes with proper-push-forward: if $f : X \rightarrow Y$ is a proper S -morphism, then the canonical base-change morphism of functors to $D_c^b(Y \times_s \eta, \overline{\mathbb{Q}}_\ell)$ is an isomorphism:*

$$R\Psi f_* \xrightarrow{\sim} f_* R\Psi;$$

In particular, if $X \rightarrow S$ is proper there is a $\text{Gal}(\bar{\eta}/\eta)$ -equivariant isomorphism

$$H^i(X_{\bar{\eta}}, \overline{\mathbb{Q}}_\ell) = H^i(X_{\bar{s}}, R\Psi(\overline{\mathbb{Q}}_\ell)).$$

- (b) *Suppose $f : X \rightarrow S$ is finite-type but not proper. Suppose that there is a compactification $j : X \hookrightarrow \bar{X}$ over S such that the boundary $\bar{X} \setminus X$ is a relative normal crossings divisor over S . Then there is a $\text{Gal}(\bar{\eta}/\eta)$ -equivariant isomorphism*

$$H_c^i(X_{\bar{\eta}}, \overline{\mathbb{Q}}_\ell) = H_c^i(X_{\bar{s}}, R\Psi(\overline{\mathbb{Q}}_\ell)).$$

- (c) *$R\Psi$ commutes with smooth pull-back: if $p : X \rightarrow Y$ is a smooth S -morphism, then the base-change morphism is an isomorphism:*

$$p^* R\Psi \xrightarrow{\sim} R\Psi p^*.$$

- (d) If $\mathcal{F} \in D_c^b(X, \overline{\mathbb{Q}}_\ell)$, we define $R\Phi(\mathcal{F})$ (the vanishing cycles) to be the cone of the canonical morphism

$$\mathcal{F}_{\bar{s}} \rightarrow R\Psi(\mathcal{F}_\eta);$$

there is a distinguished triangle

$$\mathcal{F}_{\bar{s}} \rightarrow R\Psi(\mathcal{F}_\eta) \rightarrow R\Phi(\mathcal{F}) \rightarrow \mathcal{F}_{\bar{s}}[1].$$

If $X \rightarrow S$ is smooth, then $R\Phi(\overline{\mathbb{Q}}_\ell) = 0$; in particular, $\overline{\mathbb{Q}}_\ell \cong R\Psi(\overline{\mathbb{Q}}_\ell)$ in this case;

- (e) $R\Psi$ commutes with Verdier duality, and preserves perversity of sheaves (for the middle perversity);
- (f) For $x \in X_{\bar{s}}$, $R^i\Psi(\overline{\mathbb{Q}}_\ell)_x = H^i(X_{(\bar{x})\bar{\eta}}, \overline{\mathbb{Q}}_\ell)$, where $X_{(\bar{x})\bar{\eta}}$ is the fiber over $\bar{\eta}$ of the strict henselization of X in a geometric point \bar{x} with center x . In particular, the support of $R\Psi(\overline{\mathbb{Q}}_\ell)$ is contained in the scheme-theoretic closure of X_η in $X_{\bar{s}}$;
- (g) If the generic fiber X_η is non-singular, then the complex $R\Psi^X(\overline{\mathbb{Q}}_\ell)$ is mixed, in the sense of [Weil2].

REMARK 10.2. The “fake” unitary Shimura varieties $Sh(\mathbf{G}, h)_{\mathbf{K}}$ discussed in §5.2 are proper over \mathcal{O}_E and hence by (a) we can use nearby cycles to study their semi-simple local zeta functions. The Siegel modular schemes of §5.3 are not proper over \mathbb{Z}_p , so (a) does not apply directly; in fact it is not a priori clear that there is a Galois equivariant isomorphism on cohomology with compact supports as in (b). In order to apply the method of nearby cycles to study the semi-simple local zeta function, we need such an isomorphism.

CONJECTURE 10.3. Let Sh_{K_p} denote a model over \mathcal{O}_E for a PEL Shimura variety with parahoric level structure K_p , as in §5. Then the natural morphism

$$H_c^i(Sh_{K_p} \times_{\mathcal{O}_E} \overline{k_E}, R\Psi(\overline{\mathbb{Q}}_\ell)) \rightarrow H_c^i(Sh_{K_p} \times_{\mathcal{O}_E} \overline{\mathbb{Q}}_p, \overline{\mathbb{Q}}_\ell)$$

is an isomorphism.

In the Siegel case one should be able to prove this by finding a suitably nice compactification, perhaps by adapting the methods of [CF].

Such an isomorphism would allow us to study the semi-simple local zeta function in the Siegel case by the same approach applied to the “fake” unitary case in §11. The local geometric problems involving nearby cycles have already been resolved in [HN1], and Conjecture 10.3 encapsulates the remaining geometric difficulty (which is global in nature). There will be additional group-theoretic problems in applying the Arthur-Selberg trace formula, however, due to endoscopy.

10.2. Concerning the inertia action on certain nearby cycles. Let $Sh(\mathbf{G}, h)_{\mathbf{K}}$ denote a “fake” unitary Shimura variety as in §5.2, where K_p is an Iwahori subgroup. Suppose Sh is its integral model over $\mathcal{O}_E = \mathbb{Z}_p$ defined by the moduli problem in §5.2.4. Our goal in §11 is to explain how to identify its Z_p^{ss} with a product of semi-simple local L -functions (see Theorem 11.7). We first want to justify our earlier claim that the simplified definitions of Z^{ss} and L^{ss} apply to this case. The key ingredient is a theorem of D. Gaitsgory showing that the inertia action on certain nearby cycles is unipotent.

We first recall the theorem of Gaitsgory, which is contained in [Ga]. Let λ denote a dominant coweight of $G_{\overline{\mathbb{Q}}_p}$ and consider the corresponding $G(\mathbb{Q}_p[[t]])$ -orbit \mathcal{Q}_λ in the affine Grassmannian $\text{Grass}_{\overline{\mathbb{Q}}_p} = G(\overline{\mathbb{Q}}_p((t)))/G(\overline{\mathbb{Q}}_p[[t]])$. Let M_λ denote the scheme-theoretic closure of \mathcal{Q}_λ in the deformation M of $\text{Grass}_{\overline{\mathbb{Q}}_p}$ to $\mathcal{F}l_{\overline{\mathbb{F}}_p}$, from Remark 4.1. Let IC_λ denote the intersection complex of the closure $\overline{\mathcal{Q}}_\lambda$.

THEOREM 10.4 (Gaitsgory [Ga]). *The inertia group Γ_p^0 acts unipotently on $R\Psi^{M_\lambda}(IC_\lambda)$.*

See also [GH], §5, for a detailed proof of this theorem. We remark that Gaitsgory proves this statement for nearby cycles taken with respect to Beilinson’s deformation of the affine Grassmannian of $G(\mathbb{F}_p[[t]])$ to its affine flag variety, but the same proof applies in the present p -adic setting; see loc. cit.

We can apply this to the local model $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$. Because the morphisms in the local model diagram are smooth and surjective (Lemma 13.1), by taking [GH], Lemma 5.6 into account, we see that unipotency of nearby cycles on \mathbf{M}^{loc} implies unipotency of nearby cycles on Sh :

COROLLARY 10.5. *The inertia group Γ_p^0 acts unipotently on $R\Psi^{Sh}(\overline{\mathbb{Q}}_\ell)$. Consequently, by Theorem 10.1 (a), it also acts unipotently on $H^i(Sh \times_{\mathbb{Q}_p} \overline{\mathbb{Q}}_p, \overline{\mathbb{Q}}_\ell)$.*

This justifies the assertion made in Remark 9.5. Together with Remark 9.9, we thus see that the simplified definitions of Z^{ss} and L^{ss} apply in this case, which will be helpful in §11 below.

11. The semi-simple local zeta function for “fake” unitary Shimura varieties

We assume in this section that Sh is the model from §5.2: a “fake” unitary Shimura variety. We also assume that K_p is the “standard” Iwahori subgroup of $\mathbf{G}(\mathbb{Q}_p)$, i.e., the subgroup stabilizing the “standard” self-dual multichain of $\mathcal{O}_B \otimes \mathbb{Z}_p$ -lattices

$$\tilde{\Lambda}_\bullet = \Lambda_\bullet \oplus \Lambda_\bullet^*$$

from §5.2.3.

Following the strategy of Kottwitz [Ko92], [Ko92b], we will explain how to express $Z_p^{ss}(s, Sh)$ in terms of the functions $L^{ss}(s, \pi_p, r)$.

There are two equations to be proved. The first equation is an expression for the semi-simple Lefschetz number

$$(11.0.1) \quad \sum_{x \in Sh(k_r)} \text{Tr}^{ss}(\Phi_p^r, R\Psi_x(\overline{\mathbb{Q}}_\ell)) = \sum_{\gamma_0} \sum_{(\gamma, \delta)} c(\gamma_0; \gamma, \delta) O_\gamma(f^p) TO_{\delta\sigma}(\phi_r).$$

The left hand side is termed the semi-simple Lefschetz number $\text{Lef}^{ss}(\Phi_p^r)$. The right hand side has exactly the same form (with essentially the same notation, see below) as [Ko92], p. 442¹⁰. Recall that the twisted orbital integral is defined as

$$TO_{\delta\sigma}(\phi_r) = \int_{\mathbf{G}_{\delta\sigma} \backslash \mathbf{G}(L_r)} \phi_r(x^{-1} \delta\sigma(x)),$$

¹⁰Note that the factor $|\ker^1(\mathbb{Q}, \mathbf{G})| = |\ker^1(\mathbb{Q}, Z(\mathbf{G}))|$ appearing in loc. cit. does not appear here since our assumptions on \mathbf{G} guarantee that this number is 1; see loc. cit. §7.

with an appropriate choice of measures (and $O_\gamma(f^p)$ is similarly defined). However in contrast to loc. cit., here the Haar measure on $\mathbf{G}(L_r)$ is the one giving the standard Iwahori subgroup $I_r \subset \mathbf{G}(L_r)$ volume 1.

The second equation relates the sum of (twisted) orbital integrals on the right to the spectral side of the Arthur-Selberg trace formula for \mathbf{G} :

$$(11.0.2) \quad \sum_{\gamma_0} \sum_{(\gamma, \delta)} c(\gamma_0; \gamma, \delta) O_\gamma(f^p) TO_{\delta\sigma}(\phi_r) = \sum_{\pi} m(\pi) \operatorname{Tr} \pi(f^p f_p^{(r)} f_\infty).$$

All notation on the right hand side is as in [Ko92b] (cf. §4) where the “fake” unitary Shimura varieties were analyzed in the case that K_p is a *hyperspecial maximal compact* subgroup rather than an Iwahori. In particular, π ranges over irreducible admissible representations of $\mathbf{G}(\mathbb{A})$ which occur in the discrete part of $L^2(\mathbf{G}(\mathbb{Q})A_{\mathbf{G}}(\mathbb{R})^0 \backslash \mathbf{G}(\mathbb{A}))$, with multiplicity $m(\pi)$. The function $f^p \in C_c^\infty(K^p \backslash \mathbf{G}(\mathbb{A}_f^p)/K^p)$ is just the characteristic function $\mathbb{1}_{K^p}$ of K^p , whereas f_∞ is the much more mysterious function from [Ko92b], §1¹¹. The function $f_p^{(r)}$ is a kind of “base-change” of ϕ_r and will be further explained below.

The equality (11.0.2) comes from the “pseudo-stabilization” of its left hand side, similar to that done in [Ko92b], §4. One important ingredient in that is the “base-change fundamental lemma” between the test function ϕ_r and its “base-change” $f_p^{(r)}$; the novel feature here is that the function ϕ_r is more complicated than when K_p is maximal compact: it is not a spherical function, but rather an element in the center of an Iwahori-Hecke algebra (see below). The “base-change fundamental lemma” for such functions is proved in [HN3], and further discussion will be omitted here. Finally, after pseudo-stabilization and the fundamental lemma, we apply a simple form of the Arthur-Selberg trace formula, which produces the right hand side of (11.0.2). See [HN3] for details.

Our object here is to explain (11.0.1), following the strategy of [Ko92] which handles the case where K_p is maximal compact (the case of good reduction). The main difficulty is to identify the test function ϕ_r that appears in the right hand side.

11.1. Finding the test function ϕ_r via the Kottwitz conjecture. To understand the function ϕ_r , we will use the full strength of our description of local models in §6.3, in particular §6.3.3, and also the material in the appendix §14.

In (11.0.1), the index γ_0 roughly parametrizes polarized n^2 -dimensional B -abelian varieties over k_r , up to $\overline{\mathbb{Q}}$ -isogeny. The index (γ, δ) roughly parametrizes those polarized n^2 -dimensional B -abelian varieties, up to \mathbb{Q} -isogeny, which belong to the $\overline{\mathbb{Q}}$ -isogeny class indexed by γ_0 . (For precise statements, see [Ko92].) Therefore, the summand roughly counts (with “multiplicity”) the elements in $Sh(k_r)$ which belong to a fixed \mathbb{Q} -isogeny class. We will make this last statement precise, and also explain the crucial meaning of “multiplicity” here.

Let us fix a polarized n^2 -dimensional B -abelian variety over k_r , up to isomorphism: (A', \mathcal{X}', i') . We assume it possesses some K^p -level structure $\bar{\eta}'$. We fix once and for all an isomorphism of skew-Hermitian $\mathcal{O}_B \otimes \mathbb{A}_f^p$ -modules

$$(11.1.1) \quad V \otimes \mathbb{A}_f^p = H_1(A', \mathbb{A}_f^p),$$

¹¹Up to the sign $(-1)^{\dim(Sh_E)}$, a pseudo-coefficient of a representation π_∞^0 in the packet of discrete series $\mathbf{G}(\mathbb{R})$ -representations having trivial central and infinitesimal characters.

(in the terminology of [Ko92], §4). Since it comes from a level-structure η' , this isomorphism is Galois-equivariant (with the trivial action on $V \otimes \mathbb{A}_f^p$).

Associated to (A', λ', i') is also an L_r -isocrystal (H'_{L_r}, Φ) as in §14. In brief, $H' = H(A')$ is the the $W(k_r)$ -dual of $H^1_{\text{crys}}(A'/W(k_r))$, and Φ is the σ -linear bijection on H'_{L_r} such that $p^{-1}H' \supset \Phi H' \supset H'$ (i.e., Φ is V^{-1} , where V is the Verschiebung from Cor. 14.4). Because of (11.1.1) and the determinant condition, there is also an isomorphism of skew-Hermitian $\mathcal{O}_B \otimes L_r$ -modules

$$(11.1.2) \quad V \otimes_{\mathbb{Q}_p} L_r = H'_{L_r}$$

which we also fix once and for all. (See [Ko92], p. 430.)

From these isomorphisms we construct the elements $(\gamma_0; \gamma, \delta)$ that appear in (11.0.1). Namely, the absolute Frobenius $\pi_{A'}$ for A'/k_r acting on $H_1(A', \mathbb{A}_f^p)$ induces the automorphism $\gamma^{-1} \in \mathbf{G}(\mathbb{A}_f^p)$, and the σ -linear bijection Φ acting on H'_{L_r} induces the element $\delta\sigma$, for $\delta \in \mathbf{G}(L_r)$. The element $\gamma_0 \in \mathbf{G}(\mathbb{Q})$ is constructed from (γ, δ) as in [Ko92], §14. The existence of γ_0 is proved roughly as follows. By Cor. 14.4, we have $\Phi^r = \pi_{A'}^{-1}$ acting on H'_{L_r} . Hence the elements $\gamma_l \in \mathbf{G}(\mathbb{Q}_l)$ (for $l \neq p$) and $N\delta \in \mathbf{G}(L_r)$ come from l -adic (resp. p -adic) realizations of the endomorphism $\pi_{A'}^{-1}$ of A' . Using Honda-Tate theory (and a bit more), one can view $\pi_{A'}^{-1}$ as a semi-simple element $\gamma_0 \in \mathbf{G}(\mathbb{Q})$, which is well-defined up to stable conjugacy. By its very construction, γ_0 is stably conjugate to γ , resp. $N\delta$.

Let $(A_\bullet, \lambda, i, \bar{\eta}) \in Sh(k_r)$ be a point in the moduli problem (§5.2.4). We want to classify those such that (A_0, λ, i) is \mathbb{Q} -isogenous to (A', λ', i') . Let us consider the category

$$\{(A_\bullet, \lambda, i, \xi)\}$$

consisting of chains of polarized \mathcal{O}_B -abelian varieties over k_r (up to $\mathbb{Z}_{(p)}$ -isogeny), equipped with a \mathbb{Q} -isogeny of polarized \mathcal{O}_B -abelian varieties $\xi : A_0 \rightarrow A'$, defined over k_r . Of course the integral “isocrystal” functor $A \mapsto H(A)$ of §14 is a covariant functor from this category to the category of $W(k_r)$ -free $\mathcal{O}_B \otimes W(k_r)$ -modules in H'_{L_r} equipped with Frobenius and Verschiebung endomorphisms. In fact, $(A_\bullet, \lambda, i, \xi) \mapsto \xi(H(A_\bullet))$ gives an isomorphism

$$\{(A_\bullet, \lambda, i, \xi)\} \xrightarrow{\sim} Y_p,$$

where Y_p is the category consisting of all type $(\tilde{\Lambda}_\bullet)$ multichains of $\mathcal{O}_B \otimes W(k_r)$ -lattices H_\bullet in $H'_{L_r} = V \otimes L_r$, self-dual up to a scalar in \mathbb{Q}^\times ¹², such that for each i , $p^{-1}H_i \supset \Phi H_i \supset H_i$, and $\sigma^{-1}(\Phi H_i/H_i)$ satisfies the determinant condition.

11.1.1. *Interpreting the determinant condition.* The final condition on H_i comes from the determinant condition on $\text{Lie}(A_i)$, and Cor. 14.4. Let us see what this means more concretely. By Morita equivalence (see §6.3.3), a type $(\tilde{\Lambda}_\bullet)$ multichain of $\mathcal{O}_B \otimes W(k_r)$ -lattices H_\bullet , self-dual up to a scalar in \mathbb{Q}^\times , inside $H'_{L_r} = V \otimes_{\mathbb{Q}_p} L_r$ can be regarded as a complete $W(k_r)$ -lattice chain H_\bullet^0 in L_r^n . By working with H_i^0 instead of H_i , we can work in L_r^n instead of $V \otimes_{\mathbb{Q}_p} L_r$ (which has $\dim_{L_r} = 2n^2$). Recall our minuscule coweight $\mu = (0^{n-d}, (-1)^d)$ of GL_n (§5.2.2), and write Φ for the Morita equivalent σ -linear bijection of L_r^n . The determinant condition now reads

$$\Phi H_i^0/H_i^0 \cong \sigma(k_r)^d,$$

¹²In particular, these multichains are *polarized* in the sense of [RZ], Def. 3.14.

that is, the relative position of the $W(k_r)$ -lattices H_i^0 and ΦH_i^0 in L_r^n is given by

$$\text{inv}_K(H_i^0, \Phi H_i^0) = \sigma(\mu(p)) = \mu(p).$$

(We write $\mu(p)$ in place of μ to emphasize our convention that coweights λ are embedded in $\text{GL}_n(\mathbb{Q}_p)$ by the rule $\lambda \mapsto \lambda(p)$.) The same identity holds for H_i replacing H_i^0 , when we interpret μ as a coweight for the group $\mathbf{G}(L_r) \subset \text{Aut}_B(V \otimes_{\mathbb{Q}_p} L_r)$.

By Theorem 6.3 (and the proof of [Ko92], Lemma 7.2), we may find $x \in \mathbf{G}(L_r)$ such that

$$H_\bullet = x\tilde{\Lambda}_{\bullet, W(k_r)},$$

where $\tilde{\Lambda}_{\bullet, W(k_r)} = \tilde{\Lambda}_\bullet \otimes_{\mathbb{Z}_p} W(k_r)$ is the “standard” self-dual multichain of $\mathcal{O}_B \otimes W(k_r)$ -lattices in $V \otimes_{\mathbb{Q}_p} L_r$.

The determinant condition now reads: for every index i in the chain $\tilde{\Lambda}_\bullet$,

$$(11.1.3) \quad \text{inv}_K(\tilde{\Lambda}_{i, W(k_r)}, x^{-1}\delta\sigma(x)\tilde{\Lambda}_{i, W(k_r)}) = \mu(p).$$

Letting $I_r \subset \mathbf{G}(L_r)$ denote the stabilizer of $\tilde{\Lambda}_{\bullet, W(k_r)}$, we have the Bruhat-Tits decomposition

$$\widetilde{W}(\text{GL}_n \times \mathbb{G}_m) \cong \widetilde{W}(\mathbf{G}) = I_r \backslash \mathbf{G}(L_r) / I_r.$$

Equation (11.1.3) recalls the definition of the μ -permissible set (§4.3). The determinant condition can now be interpreted as:

$$x^{-1}\delta\sigma(x) \in I_r w I_r$$

for some $w \in \text{Perm}^{\mathbf{G}}(\mu)$. The equality $\text{Adm}^{\mathbf{G}}(\mu) = \text{Perm}^{\mathbf{G}}(\mu)$ holds (this translates under Morita equivalence to the analogous statement for GL_n), and therefore we have proved that the determinant condition can now be interpreted as:

$$x^{-1}\delta\sigma(x)\tilde{\Lambda}_{\bullet, W(k_r)} \in \mathbf{M}_\mu(k_r),$$

where by definition $\mathbf{M}_\mu(k_r)$ is the set of type $(\tilde{\Lambda}_\bullet)$ multichains of $\mathcal{O}_B \otimes W(k_r)$ -lattices in $V \otimes L_r$, self-dual up to a scalar in \mathbb{Q}^\times , of form

$$g\tilde{\Lambda}_{\bullet, W(k_r)}$$

for some $g \in \mathbf{G}(L_r)$ such that $I_r g I_r = I_r w I_r$ for an element $w \in \text{Perm}^{\mathbf{G}}(\mu) = \text{Adm}^{\mathbf{G}}(\mu)$.

Now let I denote the \mathbb{Q} -group of self- \mathbb{Q} -isogenies of (A', λ', i') . Our above remarks and the discussion in [Ko92], §16 show that there is a bijection from the set of points $(A_\bullet, \lambda, i, \bar{\eta}) \in \text{Sh}(k_r)$ such that (A_0, λ, i) is \mathbb{Q} -isogenous to (A', λ', i') , to the set $I(\mathbb{Q}) \backslash (Y^p \times Y_p)$, where

$$Y^p = \{y \in \mathbf{G}(\mathbb{A}_f^p) / K^p \mid y^{-1}\gamma y \in K^p\},$$

$$Y_p = \{x \in \mathbf{G}(L_r) / I_r \mid I_r x^{-1}\delta\sigma(x)\tilde{\Lambda}_{\bullet, W(k_r)} \subset \mathbf{M}_\mu(k_r)\}.$$

11.1.2. *Compatibility of “de Rham” and “crystalline” maps.*

LEMMA 11.1. *Fix $w \in \text{Adm}(\mu)$. Let $A = (A_\bullet, \lambda, i, \bar{\eta}) \in \text{Sh}(k_r)$. Suppose that A is \mathbb{Q} -isogenous to (A', λ', i') and that via a choice of $\xi : A_0 \rightarrow A'$ we have $\xi H(A_\bullet) = x\tilde{\Lambda}_{\bullet, W(k_r)}$ as above, for $x \in \mathbf{G}(L_r) / I_r$ ¹³.*

¹³Note that $x \in \mathbf{G}(L_r) / I_r$ depends on ξ , but its image in $I(\mathbb{Q}) \backslash \mathbf{G}(L_r) / I_r$ is independent of ξ , and hence the double coset $I_r x^{-1}\delta\sigma(x) I_r$ is well-defined.

Then A belongs to the KR-stratum Sh_w if and only if

$$I_r x^{-1} \delta \sigma(x) I_r = I_r w I_r.$$

PROOF. Recall that the local model \mathbf{M}^{loc} is naturally identified with the model $M_{-w_0\mu}$ and as such its special fiber carries an action of the standard Iwahori-subgroup of $\text{GL}_n(k_r[[t]])$; further the KR-stratum Sh_w is the set of points which give rise to a point in the Iwahori-orbit indexed by w^{-1} under the “de Rham” map $\psi : \widetilde{Sh} \rightarrow \mathbf{M}^{\text{loc}}$. Let us recall the definition of ψ . We choose any isomorphism

$$\gamma_\bullet : M(A_\bullet) \xrightarrow{\sim} \widetilde{\Lambda}_{\bullet, k_r}$$

of polarized $\mathcal{O}_B \otimes k_r$ -multichains. The quotient $M(A_\bullet) \rightarrow \text{Lie}(A_\bullet)$ then determines via Morita equivalence a quotient

$$\frac{\mathcal{V}_{\bullet, k_r}}{t\mathcal{V}_{\bullet, k_r}} \rightarrow \frac{\mathcal{V}_{\bullet, k_r}}{\mathcal{L}_\bullet}$$

for a uniquely determined $k_r[[t]]$ -lattice chain \mathcal{L}_\bullet satisfying $t\mathcal{V}_{\bullet, k_r} \subset \mathcal{L}_\bullet \subset \mathcal{V}_{\bullet, k_r}$ (see §8.1). The “de Rham” map ψ sends the point $(A, \gamma_\bullet) \in \widetilde{Sh}(k_r)$ to \mathcal{L}_\bullet . By definition $A \in Sh_w$ if and only if

$$\text{inv}_I(\mathcal{L}_\bullet, \mathcal{V}_{\bullet, k_r}) = w,$$

where the invariant measures the relative position of complete $k_r[[t]]$ -lattice chains in $k_r((t))^n$. (Note that w is independent of the choice of γ_\bullet .)

In the previous paragraph the ambient group is $\text{GL}_n(k_r((t)))$, the function-field analogue of $\text{GL}_n(L_r)$. But it is more natural to work directly with $\mathbf{G}(L_r)$. Given \mathcal{L}_\bullet as above, there exists a unique type $(\widetilde{\Lambda}_\bullet)$ polarized multichain of $\mathcal{O}_B \otimes W(k_r)$ -lattices $\widetilde{\mathcal{L}}_\bullet$ in $V \otimes_{\mathbb{Q}_p} L_r$ such that $p\widetilde{\Lambda}_{\bullet, W(k_r)} \subset \widetilde{\mathcal{L}}_\bullet \subset \widetilde{\Lambda}_{\bullet, W(k_r)}$ and the polarized $\mathcal{O}_B \otimes k_r$ -multichain $\widetilde{\Lambda}_{\bullet, W(k_r)} / \widetilde{\mathcal{L}}_\bullet$ is Morita equivalent to the k_r -lattice chain $\mathcal{V}_{\bullet, k_r} / \mathcal{L}_\bullet$. Thus we can think of ψ as the map

$$(A, \gamma_\bullet) \mapsto \widetilde{\mathcal{L}}_\bullet.$$

Thus, $A \in Sh_w$ if and only if

$$\text{inv}_{I_r}(\widetilde{\mathcal{L}}_\bullet, \widetilde{\Lambda}_{\bullet, W(k_r)}) = w.$$

We have an isomorphism of polarized multichains of $\mathcal{O}_B \otimes k_r$ -lattices

$$\frac{\widetilde{\Lambda}_{\bullet, W(k_r)}}{\widetilde{\mathcal{L}}_\bullet} = \text{Lie}(A_\bullet) = \sigma^{-1} \left(\frac{x^{-1} \delta \sigma(x) \widetilde{\Lambda}_{\bullet, w(k_r)}}{\widetilde{\Lambda}_{\bullet, W(k_r)}} \right).$$

The second equality comes from Cor. 14.4; the map sending (A_\bullet, ξ) to the multichain $x\widetilde{\Lambda}_{\bullet, W(k_r)}$ may be termed “crystalline” – it is defined using crystalline homology. This equality shows that the “de Rham” and “crystalline” maps are compatible (and ultimately rests on Theorem 14.2 of Oda).

Putting these remarks together, we see that $A \in Sh_w$ if and only if

$$\text{inv}_{I_r}(\widetilde{\Lambda}_{\bullet, W(k_r)}, x^{-1} \delta \sigma(x) \widetilde{\Lambda}_{\bullet, W(k_r)}) = w,$$

which completes the proof. □

REMARK 11.2. The crux of the above proof is the aforementioned compatibility between the “de Rham” and “crystalline” maps. This compatibility can be rephrased as the commutativity of the diagram at the end of §7 in [R2], when the morphisms there are suitably interpreted.

11.1.3. *Identifying ϕ_r .* To find ϕ_r we need to count the points in the set $I(\mathbb{Q}) \backslash (Y^p \times Y_p)$ with the correct “multiplicity”. The test function ϕ_r in the twisted orbital integral must be such that

$$(11.1.4) \quad \sum_A \text{Tr}^{ss}(\Phi_p^r; R\Psi_A^{Sh}(\overline{\mathbb{Q}}_\ell)) = \int_{I(\mathbb{Q}) \backslash (\mathbf{G}(\mathbb{A}_f^p) \times \mathbf{G}(L_r))} f^p(y^{-1}\gamma y) \phi_r(x^{-1}\delta\sigma(x)),$$

where A ranges over points $(A_\bullet, \lambda, i, \bar{\eta}) \in Sh(k_r)$ such that (A_0, λ, i) is \mathbb{Q} -isogenous to (A', λ', i') (other notation and measures as in [Ko92], §16, with the exception that here the Haar measure on $G(L_r)$ gives I_r volume 1).

Now by Lemma 11.1, equation (11.1.4) will hold if ϕ_r is a function in the Iwahori-Hecke algebra of $\overline{\mathbb{Q}}_\ell$ -valued functions

$$\mathcal{H}_{I_r} = C_c(I_r \backslash \mathbf{G}(L_r) / I_r)$$

such that

$$\phi_r(I_r w I_r) = \text{Tr}^{ss}(\Phi_p^r, R\Psi_{w^{-1}}^{M_{-w_0\mu}}(\overline{\mathbb{Q}}_\ell)),$$

for elements $w \in \text{Adm}(\mu)$, and zero elsewhere. Note that on the right hand side, w^{-1} really represents an Iwahori-orbit in the affine flag variety $\mathcal{Fl}_{\mathbb{F}_p}$ over the *function field*. The nearby cycles are equivariant for the Iwahori-action in a suitable sense, so that the semi-simple trace function is constant on these orbits. Hence the right hand side is a well-defined element of $\overline{\mathbb{Q}}_\ell$.

We can simply *define* the function ϕ_r by this equality. But such a description of ϕ_r will not be useful unless we can identify it with an explicit function in the Iwahori-Hecke algebra (we need to know its traces on representations with Iwahori-fixed vectors, at least, if we want to make the spectral side of (11.0.2) explicit). This however is possible, due to the following theorem. This result was conjectured by Kottwitz in a more general form, which inspired Beilinson to conjecture that nearby cycles can be used to give a geometric construction of the center of the affine Hecke algebra for *any* reductive group in the function-field setting. This latter conjecture was proved by Gaitsgory [Ga], whose ideas were adapted to prove the p-adic analogue in [HN1].

THEOREM 11.3 (The Kottwitz Conjecture; [Ga], [HN1]). *Let $G = \text{GL}_n$ or GSp_{2n} . Let λ be a minuscule dominant coweight of G , with corresponding \mathbb{Z}_p -model M_λ (cf. Remark 4.1). Let $\mathcal{H}_{k_r((t))}$ denote the Iwahori-Hecke algebra $C_c(I_r \backslash G(k_r((t))) / I_r)$. Let $z_{\lambda,r} \in Z(\mathcal{H}_{k_r((t))})$ denote the Bernstein function in the center of the Iwahori-Hecke algebra which is associated to λ . Then*

$$\text{Tr}(\Phi_p^r, R\Psi^{M_\lambda}(\overline{\mathbb{Q}}_\ell)) = p^{r \dim(M_\lambda)/2} z_{\lambda,r}.$$

Let K_r be the stabilizer in $\mathbf{G}(L_r)$ of $\tilde{\Lambda}_{0,W(k_r)}$; this is a hyperspecial maximal compact subgroup of $\mathbf{G}(L_r)$ containing I_r . The Bernstein function $z_{\lambda,r}$ is characterized as the unique element in the center of \mathcal{H}_{I_r} such that the image of $p^{r\langle \rho, \lambda \rangle} z_{\lambda,r}$ under the Bernstein isomorphism

$$- * \mathbb{I}_{K_r} : Z(\mathcal{H}_{I_r}) \xrightarrow{\sim} \mathcal{H}_{K_r} := C_c(K_r \backslash \mathbf{G}(L_r) / K_r)$$

is the spherical function $f_{\lambda,r} := \mathbb{I}_{K_r \lambda K_r}$. Here, we have used the fact that λ is minuscule. Recall that ρ is the half-sum of the B -positive roots of G , and that $\dim(M_\lambda)/2 = \langle \rho, \lambda \rangle$.

The above theorem for $\lambda = -w_0\mu$ implies that

$$\phi_r(w) = p^{r \dim(Sh)/2} z_{-w_0\mu}(w^{-1}).$$

Now invoking the identity $z_\mu(w) = z_{-w_0\mu}(w^{-1})$ (see [HKP], §3.2), we have proved the following result.

PROPOSITION 11.4. *Let $z_{\mu,r}$ denote the Bernstein function in the center of the Iwahori-Hecke algebra $\mathcal{H}_{L_r} = C_c(I_r \backslash \mathbf{G}(L_r)/I_r)$ corresponding to μ . Then the test function is given by*

$$\phi_r = p^{r \dim(\text{Sh})/2} z_{\mu,r}.$$

Remarks: 1) Because ϕ_r is central we can define its “base-change” function $b(\phi_r) =: f_p^{(r)}$, an element in the center of the Iwahori-Hecke algebra for $\mathbf{G}(\mathbb{Q}_p)$. We define the base-change homomorphism for centers of Iwahori-Hecke algebras as the unique homomorphism $b : Z(\mathcal{H}_{I_r}) \rightarrow Z(\mathcal{H}_I)$ which induces, via the Bernstein isomorphism, the usual base-change homomorphism for spherical Hecke algebras $b : \mathcal{H}_{K_r} \rightarrow \mathcal{H}_K$ for any special maximal compact K_r containing I_r (setting $I = I_r \cap \mathbf{G}(\mathbb{Q}_p)$ and $K = K_r \cap \mathbf{G}(\mathbb{Q}_p)$). This gives a well-defined homomorphism, independent of the choice of K_r . Moreover, the pair of functions $\phi_r, f_p^{(r)}$ have matching (twisted) orbital integrals; see [HN3].

2) We know how $z_{\mu,r}$ (and hence how its base-change $b(z_{\mu,r})$) acts on unramified principal series. This plays a key role in Theorem 11.7 below. In fact, we have the following helpful lemma.

LEMMA 11.5. *Let I denote our Iwahori subgroup $K_p^{\mathfrak{a}} \subset G(\mathbb{Q}_p)$, whose reduction modulo p is $B(\mathbb{F}_p)$. Suppose π_p is an irreducible admissible representation of $G(\mathbb{Q}_p)$ with $\pi_p^I \neq 0$. Let r_μ be the irreducible representation of ${}^L\mathbf{G}_{\mathbb{Q}_p}$ having extreme weight μ . Let $d = \dim(\text{Sh}_E)$. Then*

$$\text{Tr } \pi_p(p^{rd/2} b(z_{\mu,r})) = \dim(\pi_p^I) p^{rd/2} \text{Tr}(r_\mu \varphi_{\pi_p}(\Phi^r \times \begin{bmatrix} p^{-r/2} & 0 \\ 0 & p^{r/2} \end{bmatrix})).$$

PROOF. Write $G = G(\mathbb{Q}_p)$ and $B = B(\mathbb{Q}_p)$ and suppose that π_p is an irreducible subquotient of the normalized unramified principal series representation $i_B^G(\chi)$, for an unramified quasi-character $\chi : T(\mathbb{Q}_p) \rightarrow \mathbb{C}^\times$. Suppose $K_r \supset I_r$ (thus also $K \supset I$) is a hyperspecial maximal compact, and suppose that π_χ is the unique K -spherical subquotient of $i_B^G(\chi)$. Suppose $\varphi : W_{\mathbb{Q}_p} \rightarrow {}^L G$ is the unramified parameter associated to π_χ . Our normalization of the correspondence $\pi_p \mapsto \varphi_{\pi_p}$ is such that for all $r \geq 1$, the element $\varphi(\Phi^r) \in \widehat{G} \rtimes W_{\mathbb{Q}_p}$ can be described as

$$(11.1.5) \quad \varphi(\Phi^r) = \varphi_{\pi_p}(\Phi^r \times \begin{bmatrix} p^{-r/2} & 0 \\ 0 & p^{r/2} \end{bmatrix}) = (\chi \rtimes \Phi)^r$$

where Φ is a *geometric* Frobenius element in $W_{\mathbb{Q}_p}$, and where we identify χ with an element in the dual torus $\widehat{T}(\mathbb{C}) \subset \widehat{G}(\mathbb{C})$ and take the product on the right in the group ${}^L G$. Note that our normalization of the local Langlands correspondence is the one compatible with Deligne’s normalization of the reciprocity map in local class field theory, where a uniformizer is sent to a geometric Frobenius element (see [W2]).

Let $f_{\mu,r} = \mathbb{1}_{K_r \mu K_r}$ in the spherical Hecke algebra \mathcal{H}_{K_r} ; it is the image of $p^{rd/2} z_{\mu,r}$ under the Bernstein isomorphism

$$- * \mathbb{1}_{K_r} : Z(\mathcal{H}_{I_r}) \xrightarrow{\sim} \mathcal{H}_{K_r}.$$

Further, $b(p^{rd/2} z_{\mu,r}) * \mathbb{1}_K = b(f_{\mu,r})$.

Now by [Ko84], Thm (2.1.3), we know that under the usual left action of \mathcal{H}_K , $b(f_{\mu,r})$ acts on π_χ^K by the scalar

$$p^{rd/2} \operatorname{Tr}(r_\mu \varphi(\Phi^r)).$$

Taking (11.1.5) into account along with the well-known fact that elements in $Z(\mathcal{H}_f)$ act on the entire space $i_B^G(\chi)^I$ by a scalar (see e.g. [HKP]), we are done. \square

REMARK 11.6. For application in Theorem 11.7, we need the “ ℓ -adic” analogue of this lemma, i.e., we need to work with the dual group $\widehat{G}(\overline{\mathbb{Q}}_\ell)$ instead of $\widehat{G}(\mathbb{C})$. For a discussion of how to do this, see [Ko92b], §1.

11.2. The semi-simple local zeta function in terms of semi-simple L -functions. The foregoing discussion culminates in the following result from [HN3], to which we refer for details of the proof.

THEOREM 11.7 ([HN3]). *Suppose Sh is a simple (“fake” unitary) Shimura variety. Suppose K_p is an Iwahori subgroup. Suppose r_μ is the irreducible representation of $L(\mathbf{G}_{\mathbf{E}_p})$ with extreme weight μ , where μ is the minuscule coweight determined by the Shimura data. Then we have*

$$Z_{\mathfrak{p}}^{ss}(s, Sh) = \prod_{\pi_f} L^{ss}(s - \frac{d}{2}, \pi_p, r_\mu)^{a(\pi_f) \dim(\pi_f^K)}$$

where the product runs over all admissible representations π_f of $\mathbf{G}(\mathbb{A}_f)$, and the integer number $a(\pi_f)$ is given by

$$a(\pi_f) = \sum_{\pi_\infty \in \Pi_\infty} m(\pi_f \otimes \pi_\infty) \operatorname{Tr} \pi_\infty(f_\infty),$$

where $m(\pi_f \otimes \pi_\infty)$ is the multiplicity of $\pi_f \otimes \pi_\infty$ in $L^2(\mathbf{G}(\mathbb{Q})A_{\mathbf{G}}(\mathbb{R})^0 \backslash \mathbf{G}(\mathbb{A}))$. Here Π_∞ is the set of admissible representations of $G(\mathbb{R})$ whose central and infinitesimal characters are trivial. Also, d denotes the dimension of Sh_E .

Let us remark that it is our firm belief that this result continues to hold when K_p is a general parahoric subgroup of $\mathbf{G}(\mathbb{Q}_p)$, but some details are more difficult than the Iwahori case treated in [HN3], and remain to be worked out.

Finally, recall our assumptions on \mathfrak{p} implied that $\mathbf{E}_p = \mathbb{Q}_p$ and $N\mathfrak{p} = (p)$. In more general circumstances one or both of these statements will fail to hold, and the result will be a slightly more complicated expression for $Z_{\mathfrak{p}}^{ss}(s, Sh)$.

12. The Newton stratification on Shimura varieties over finite fields

12.1. Review of the Kottwitz and Newton maps. As usual L denotes the fraction field of the Witt vectors $W(\overline{\mathbb{F}}_p)$, with Frobenius automorphism σ . Let G denote a connected reductive group over \mathbb{Q}_p . Then we have the pointed set $B(G) = B(G)_{\mathbb{Q}_p}$ consisting of σ -conjugacy classes in $G(L)$. Let us also assume (only for simplicity) that the connected reductive \mathbb{Q}_p -group G is *unramified*. We have the Kottwitz map

$$\kappa_G : B(G) \rightarrow X^*(Z(\widehat{G})^{\Gamma_p}),$$

where $\Gamma_p = \operatorname{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ (see [Ko97], §7). We also have the Newton map

$$\overline{\nu} : B(G) \rightarrow \mathfrak{U}^+$$

where the notation is as follows. We choose a \mathbb{Q}_p -rational Borel subgroup $B \subset G$ and a maximal \mathbb{Q}_p -torus T which is contained in B . Then $\mathfrak{U} = X_*(T)_{\mathbb{R}}^{\Gamma_p}$, and \mathfrak{U}^+ denotes the intersection of \mathfrak{U} with the cone of B -dominant elements in $X_*(T)_{\mathbb{R}}$. We call $b \in B(G)$ *basic* if $\bar{\nu}_b$ is central, i.e., $\bar{\nu}_b \in X_*(Z)_{\mathbb{R}}$.

Suppose $\{\lambda\}$ is a conjugacy class of one-parameter subgroups of G , defined over \mathbb{Q}_p . We may represent the class by a unique cocharacter $\bar{\lambda} \in X_*(T) = X^*(\widehat{T})$ lying in the B -positive Weyl chamber of $X_*(T)_{\mathbb{R}}$. The Weyl-orbit of $\bar{\lambda}$ is stabilized by Γ_p . The notion of B -dominant being preserved by Γ_p (since B is \mathbb{Q}_p -rational), we see that $\bar{\lambda}$ is fixed by Γ_p , hence it belongs to \mathfrak{U}^+ . Also, restricting $\{\lambda\}$ to $Z(\widehat{G})$ determines a well-defined element $\lambda^{\natural} \in X^*(Z(\widehat{G})^{\Gamma_p})$.

We can now define the subset $B(G, \lambda) \subset B(G)$ to be the set of classes $[b] \in B(G)$ such that

$$\begin{aligned} \kappa_G(b) &= \lambda^{\natural} \\ \bar{\nu}_b &\preceq \bar{\lambda}. \end{aligned}$$

Here \preceq denotes the usual partial order on \mathfrak{U}^+ for which $\nu \preceq \nu'$ if $\nu' - \nu$ is a nonnegative linear combination of simple relative coroots.

12.2. Definition of the Newton stratification. Suppose that the Shimura variety $Sh_{K_p} = Sh(\mathbf{G}, h)_{K_p K_p}$ is given by a moduli problem of abelian varieties, and that $K_p \subset \mathbf{G}(\mathbb{Q}_p)$ is a parahoric subgroup. Also, assume for simplicity that $E := \mathbf{E}_{\mathfrak{p}} = \mathbb{Q}_p$. Let $G = \mathbf{G}_{\mathbb{Q}_p}$, which again for simplicity we assume is unramified. Let k denote as usual an algebraic closure of the residue field $\mathcal{O}_E/\mathfrak{p} = \mathbb{F}_p$, and let k_r denote the unique subfield of k having cardinality p^r . Let L_r denote the fraction field of the Witt vectors $W(k_r)$.

We denote by \mathfrak{a} the base alcove and suppose $0 \in \bar{\mathfrak{a}}$ is a hyperspecial vertex. Let $K_p^{\mathfrak{a}}$ (resp. K_p^0) denote the corresponding Iwahori (resp. hyperspecial maximal compact) subgroup of $G(\mathbb{Q}_p)$. We will let K_p denote a “standard” parahoric subgroup, i.e., one such that $K_p^{\mathfrak{a}} \subset K_p \subset K_p^0$.

We can define a map

$$\delta_{K_p} : Sh_{K_p}(k) \rightarrow B(G)$$

as follows. A point $A_{\bullet} = (A_{\bullet}, \lambda, i, \bar{\eta}) \in Sh_{K_p}(k_r)$ gives rise to a c -polarized virtual B -abelian variety over k_r up to prime-to- p isogeny (cf. [Ko92] §10), which we denote by (A, λ, i) . That in turn determines an L -isocrystal $(H(A)_L, \Phi)$ as in §14, cf. loc. cit. It is not hard to see that δ_{K_p} takes values in the subset $B(G, \mu) \subset B(G)$. In fact, if we choose an isomorphism $(H(A)_L, \Phi) = (V \otimes L, \delta\sigma)$ of isocrystals for the group $G(L)$, then $\delta \in G(L)$ satisfies

$$\kappa_G(\delta) = \mu^{\natural},$$

as follows from the determinant condition $\sigma(\text{Lie}(A_0)) = \Phi H(A_0)/H(A_0)$, for any A_0 in the chain A_{\bullet} (use the argument of [Ko92], p. 431). Furthermore, the Mazur inequality

$$\bar{\nu}_{\delta} \preceq \bar{\mu}$$

can be proved by reducing to the case where $K_p = K_p^0$ (which has been treated by Rapoport-Richartz [RR] – see also [Ko03])¹⁴.

¹⁴Note that our assumption that K_p be *standard*, i.e., $K_p \subset K_p^0$, was used in the last step, because we want to invoke [RR]. The results of loc. cit. probably hold for nonspecial maximal compact subgroups, so this assumption is probably unnecessary.

DEFINITION 12.1. We call the fibers of δ_{K_p} the *Newton strata* of Sh_{K_p} . The inverse image of the basic set

$$\delta_{K_p}^{-1}(B(G, \mu) \cap B(G)_{basic}).$$

is called the *basic locus* of Sh_{K_p} .

We denote the Newton stratum $\delta_{K_p}^{-1}([b])$ by $\mathcal{S}_{K_p, [b]}$, or if K_p is understood, simply by $\mathcal{S}_{[b]}$.

The following conjecture is fundamental to the subject. It asserts that all the Newton strata are nonempty.

CONJECTURE 12.2 (Rapoport, Conj. 7.1 [R2]). *The map*

$$\delta_{K_p} : Sh_{K_p}(k) \rightarrow B(G, \mu)$$

is surjective. In particular, the basic locus is nonempty.

REMARK 12.3. Note that $\text{Im}(\delta_{K_p^0})$ can be interpreted purely in terms of group theory: $[b] \in B(G, \mu)$ lies in the image of $\delta_{K_p^0}$ if and only if for one (equivalently, for all sufficiently divisible) $r \geq 1$, $[b]$ contains an element $\delta \in G(L_r)$ belonging to a triple $(\gamma_0; \gamma, \delta)$, which satisfies the conditions in [Ko90] §2 (except for the following correction, noted at the end of [Ko92]: under the canonical map $B(G)_{\mathbb{Q}_p} \rightarrow X^*(Z(\widehat{G})^{\Gamma_p})$, the σ -conjugacy class of δ goes to $\mu|_{X^*(Z(\widehat{G})^{\Gamma_p})}$ and not its negative), and for which the following four conditions also hold:

- (a) $\gamma_0 \gamma_0^* = c_0^{-1} p^{-r}$, where $c_0 \in \mathbb{Q}^\times$ is a p -adic unit;
- (b) the Kottwitz invariant $\alpha(\gamma_0; \gamma, \delta)$ is trivial;
- (c) there exists a lattice Λ in V_{L_r} such that $\delta \sigma \Lambda \supset \Lambda$;
- (d) $O_\gamma(\mathbb{I}_{K_p}) TO_{\delta \sigma}(\mathbb{I}_{K_r, \mu K_r}) \neq 0$.

Here $K_r \subset \mathbf{G}(L_r)$ is the hyperspecial maximal compact subgroup such that $K_r \cap \mathbf{G}(\mathbb{Q}_p) = K_p^0$.

To see this, use [Ko92], Lemmas 15.1, 18.1 to show that the first three conditions ensure the existence of a $c_0 p^r$ -polarized virtual B -abelian variety (A', λ', i') over k_r up to prime-to- p isogeny, giving rise to $(\gamma_0; \gamma, \delta)$. In the presence of the first three, the last condition shows that there exists a k_r -rational point $(A, \lambda, i, \bar{\eta}) \in Sh_{K_p^0}$ such that (A, λ, i) is \mathbb{Q} -isogenous to (A', λ', i') . Hence $\delta_{K_p^0}((A, \lambda, i, \bar{\eta})) = [\delta]$.

Note that by counting fixed points of $\Phi_p^r \circ f$ for any Hecke operator f away from p as in [Ko92], §16, we may replace condition (d) with

(d') For some $g \in \mathbf{G}(\mathbb{A}_f^p)$, we have $O_\gamma(\mathbb{I}_{K_p g^{-1} K_p}) TO_{\delta \sigma}(\mathbb{I}_{K_r, \mu K_r}) \neq 0$.

Since we may always choose $g = \gamma^{-1}$, we may also replace (d) or (d') with

(d'') $TO_{\delta \sigma}(\mathbb{I}_{K_r, \mu K_r}) \neq 0$.

In the Siegel case with K_p hyperspecial, we have the following result of Oort [Oo] (comp. [R2], Thm. 7.4) which in particular proves Conjecture 12.2 in that case.

THEOREM 12.4 (Oort, [Oo]). *Suppose $Sh_{K_p^0}$ is a Siegel modular variety with maximal hyperspecial level structure K_p^0 at p . Then the Newton strata are all nonempty and equidimensional (and the dimension is given by a simple formula in terms of the partially ordered set $B(\text{GSp}_{2n}, \mu)$).*

In Corollary 12.8 below, we shall generalize the part of Theorem 12.4 that asserts the nonemptiness of the Newton strata: in the Siegel case, all Newton strata of Sh_{K_p} are nonempty, when K_p is an arbitrary standard parahoric subgroup. In the “fake” unitary case, we shall later prove in Cor. 12.12 only that the basic locus is nonempty (again for standard parahorics K_p).

12.3. The relation between Newton strata, KR-strata, and affine Deligne-Lusztig varieties. Fix a σ -conjugacy class $[b] \in B(G)$, and fix an element $w \in \widetilde{W}$. Let $\widetilde{K}_p^{\mathbf{a}} \subset G(L)$ denote the Iwahori subgroup such that $\widetilde{K}_p^{\mathbf{a}} \cap G(\mathbb{Q}_p) = K_p^{\mathbf{a}}$.

DEFINITION 12.5. We define the *affine Deligne-Lusztig variety* ¹⁵

$$X_w(b)_{\widetilde{K}_p^{\mathbf{a}}} = \{x \in G(L)/\widetilde{K}_p^{\mathbf{a}} \mid x^{-1}b\sigma(x) \in \widetilde{K}_p^{\mathbf{a}} w \widetilde{K}_p^{\mathbf{a}}\},$$

and for any dominant coweight λ ,

$$X(\lambda, b)_{\widetilde{K}_p^{\mathbf{a}}} = \bigcup_{w \in \text{Adm}(\lambda)} X_w(b)_{\widetilde{K}_p^{\mathbf{a}}}.$$

A similar definition can be made for a parahoric subgroup replacing $K_p^{\mathbf{a}}$ (cf. [R2]).

A fundamental problem is to determine the relations between the Kottwitz-Rapoport and Newton stratifications. The following shows how this problem is related to the nonemptiness of certain affine Deligne-Lusztig varieties.

PROPOSITION 12.6. *Let μ be the minuscule coweight attached to the Shimura data for $Sh = Sh_{K_p^{\mathbf{a}}}$. Suppose $w \in \text{Adm}(\mu)$. Then for every $[b] \in \text{Im}(\delta_{K_p^{\mathbf{a}}})$, we have*

$$X_w(b)_{\widetilde{K}_p^{\mathbf{a}}} \neq \emptyset \Leftrightarrow Sh_w \cap \mathcal{S}_{[b]} \neq \emptyset.$$

PROOF. By Remark 12.3, for all sufficiently divisible $r \geq 1$, $[b]$ contains an element $\delta \in G(L_r)$ which is part of a Kottwitz triple $(\gamma_0; \gamma, \delta)$ satisfying the conditions (a-d). We consider such a triple, up to equivalence (we say $(\gamma'_0; \gamma', \delta')$ is equivalent to $(\gamma_0; \gamma, \delta)$ if γ'_0 and γ_0 are stably-conjugate, γ' and γ are conjugate, and δ' and δ are σ -conjugate). Then by the arguments in §11 together with [Ko92], §18, 19, we have the equality

$$(12.3.1) \quad \#\{A_{\bullet} \in Sh_w(k_r) \mid A_{\bullet} \rightsquigarrow (\gamma_0; \gamma, \delta)\} = (\text{vol}) O_{\gamma}(\mathbb{I}_{K_p}) \#X_w(\delta)_{I_r}.$$

Let us explain the notation. The notation $A_{\bullet} \rightsquigarrow (\gamma_0; \gamma, \delta)$ means that A_{\bullet} gives rise to the equivalence class of $(\gamma_0; \gamma, \delta)$; cf. [Ko92], §18, 19. The term vol denotes the nonzero rational number $|\ker^1(\mathbb{Q}, \mathbf{G})|c(\gamma_0; \gamma, \delta)$, where the second term is the number defined in loc. cit. Also $I_r = \widetilde{K}_p^{\mathbf{a}} \cap G(L_r)$.

This equality would imply the proposition, if we knew that $O_{\gamma}(\mathbb{I}_{K_p}) \neq 0$. But this follows from condition (d) in Remark 12.3. □

The following result of Wintenberger [Wi] proves a conjecture of Kottwitz and Rapoport in a suitably unramified case (cf. [R2], Conj. 5.2, and the notes at the end).

¹⁵Strictly speaking, this is only a set, not a variety. The sets are the affine analogues of the usual Deligne-Lusztig varieties in the theory of finite groups of Lie type.

THEOREM 12.7 (Wintenberger). *Let G be any connected reductive group, defined and quasi-split over L . Suppose $\{\lambda\}$ is a conjugacy class of 1-parameter subgroups, defined over L . Suppose $[b] \in B(G)$ and let K be any standard parahoric subgroup (that is, one contained in a special maximal parahoric subgroup). Then*

$$X(\lambda, b)_K \neq \emptyset \Leftrightarrow [b] \in B(G, \lambda).$$

COROLLARY 12.8. *Let Sh be either a “fake” unitary or a Siegel modular variety, as in §5.*

- (a) *In the “fake” unitary case, for any two standard parahoric subgroups $K'_p \subset K''_p$, we have $\text{Im}(\delta_{K'_p}) = \text{Im}(\delta_{K''_p})$.*
- (b) *In the Siegel case, we have $\text{Im}(\delta_{K_p}) = B(G, \mu)$ for every standard parahoric subgroup K_p .*

PROOF. Consider first (a). We need to prove $\text{Im}(\delta_{K'_p}) \supset \text{Im}(\delta_{K''_p})$. Clearly it is enough to consider the case $K'_p = K_p^{\mathfrak{a}}$ and $K''_p = K_p^0$. The natural morphism $Sh_{K_p^{\mathfrak{a}}} \rightarrow Sh_{K_p^0}$ is proper, surjective on generic fibers (look at \mathbb{C} -points), and the target is flat (even smooth). Therefore in the special fiber the morphism is surjective. This completes the proof of (a).

Consider now part (b). In the Siegel case the morphism $Sh_{K_p^{\mathfrak{a}}} \rightarrow Sh_{K_p^0}$ is still projective with flat image, so the same argument combined with Theorem 12.4 yields the stronger result of (b). Here is another argument using Theorem 12.4, Proposition 12.6 and Theorem 12.7. Let $G = \text{GSp}_{2n}$ and $\mu = (0^n, (-1)^n)$. By Oort’s theorem, $\text{Im}(\delta_{K_p^0}) = B(G, \mu)$, so it is enough to prove $B(G, \mu) \subset \text{Im}(\delta_{K_p^{\mathfrak{a}}})$. Let $[b] \in B(G, \mu)$. By Wintenberger’s theorem, there exists $w \in \text{Adm}(\mu)$ such that $X_w(b)_{\tilde{K}_p^{\mathfrak{a}}} \neq \emptyset$, which implies the result by Proposition 12.6.

Note that a similar argument provides an alternative proof for part (a). □

REMARK 12.9. Let $G = \text{GSp}_{2n}$, and suppose μ is minuscule. We can give a proof of Theorem 12.7 in this case using Oort’s theorem, as follows. Let $[b] \in B(G, \mu)$. By Corollary 12.8, there is a point $A = (A_{\bullet}, \lambda, i, \bar{\eta})$ such that $\delta_{\tilde{K}_p^{\mathfrak{a}}}(A) = [b]$. Now A belongs to *some* KR-stratum Sh_w , so $Sh_w \cap \mathcal{S}_{[b]} \neq \emptyset$. Now Proposition 12.6 implies that $X_w([b])_{\tilde{K}_p^{\mathfrak{a}}} \neq \emptyset$.

12.4. The basic locus is nonempty in the “fake” unitary case. ¹⁶

First consider a “fake” unitary variety Sh with $\mu = (0^{n-d}, (-1)^d)$. We shall consider both hyperspecial and Iwahori level structures.

Recall that the subset $\text{Adm}(\mu) \subset \widetilde{W}(\text{GL}_n)$ contains a unique minimal element. To find this element, we consider first the element

$$\tau_1 = t_{(-1, 0^{n-1})}(12 \cdots n) \in \widetilde{W}(\text{GL}_n),$$

where the cycle $(12 \cdots n)$ acts on a vector $(x_1, x_2, \dots, x_n) \in X_*(T) \otimes \mathbb{R}$ by sending it to $(x_n, x_1, \dots, x_{n-1})$. Note that this element preserves our base alcove

$$\mathfrak{a} = \{(x_1, \dots, x_n) \in X_*(T) \otimes \mathbb{R} \mid x_n - 1 < x_1 < \dots < x_{n-1} < x_n\}.$$

Hence its d -th power

$$\tau = t_{((-1)^d, 0^{n-d})}(12 \cdots n)^d \in \widetilde{W}(\text{GL}_n),$$

¹⁶This non-emptiness is implicit in both articles of [FM], and can be justified (indirectly) from their main theorems. Our object here is only to give a simple direct proof.

is the unique element of Ω which is congruent modulo W_{aff} to t_μ , so is the desired element. This element is identified with an element in $\text{GL}_n(L)$ using our usual convention (the vector part is sent to $\text{diag}((p^{-1})^d, 1^{n-d})$). Via Morita equivalence, we can view it as an element of $\mathbf{G}(L)$. In fact, via (5.2.1), τ becomes the element

$$\tau = (X, p^{-1}\chi^{-1}(X^t)^{-1}\chi),$$

where by definition $X = \text{diag}((p^{-1})^d, 1^{n-d})(12 \cdots n)^d$.

Next consider the Siegel case, where $\mu = (0^n, (-1)^n)$. The unique element $\tau \in \Omega$ which is congruent to t_μ modulo W_{aff} is given by

$$\tau = t_{((-1)^n, 0^n)}(12 \cdots 2n)^n \in \widetilde{W}(\text{GSp}_{2n}).$$

We will show that either case, $\tau \in \mathbf{G}(L)$ is *basic*. We will also show that $[\tau]$ belongs to the image of $\delta_{K_p^0}$. By virtue of Corollary 12.8, this will show that the basic locus of Sh_{K_p} is nonempty for every standard parahoric K_p .

Let us handle the second statement first.

LEMMA 12.10. *Let $\delta = \tau \in \mathbf{G}(L)$. Then there exists a Kottwitz triple $(\gamma_0; \gamma, \delta)$ satisfying the conditions in Remark 12.3. Hence $[\tau] \in \text{Im}(\delta_{K_p^0})$.*

PROOF. Consider the “fake” unitary case. Note that $\delta \in \mathbf{G}(L_n)$ is clearly fixed by σ , so that

$$(12.4.1) \quad N\delta = \delta^n = \text{diag}((p^{-d})^n) \times \text{diag}((p^{d-n})^n),$$

where the first factor is the diagonal matrix with the entry p^{-d} repeated n times (similarly for the second factor). This is the image of a unique element $\gamma_0 \in \mathbf{G}(\mathbb{Q})$ under the composition of the inclusion $\mathbf{G}(\mathbb{Q}) \hookrightarrow \mathbf{G}(\mathbb{Q}_p)$ and the isomorphism (5.2.1). In fact γ_0 belongs to the center of \mathbf{G} , and thus is clearly an elliptic element in $\mathbf{G}(\mathbb{R})$. Moreover, $\gamma_0\gamma_0^* = p^{-n}$. For all primes $l \neq p$, we define γ_l to be the image of γ_0 under the inclusion $\mathbf{G}(\mathbb{Q}) \hookrightarrow \mathbf{G}(\mathbb{Q}_l)$. We set $\gamma = (\gamma_l)_l \in \mathbf{G}(\mathbb{A}_f^p)$. The resulting triple $(\gamma_0; \gamma, \delta)$ is clearly a Kottwitz triple satisfying the conditions (a-c,d”) of Remark 12.3 (with $r = n$). One can check that $\alpha(\gamma_0; \gamma, \delta) = 1$ from the definitions, but in fact this is not necessary, as the group to which $\alpha(\gamma_0; \gamma, \delta)$ belongs is itself trivial in the “fake” unitary case (see [Ko92b], Lemma 2). Hence by Remark 12.3, δ arises from a point in $Sh_{K_p^0}(k_n)$, i.e., $[\tau]$ is in the image of $\delta_{K_p^0}$.

In the Siegel case, the same argument works, if we let $\delta = \tau \in \text{GL}_n(L_2)$ and note

$$(12.4.2) \quad N\delta = \delta^2 = \text{diag}((p^{-1})^{2n}).$$

□

LEMMA 12.11. *The element $\tau \in G(L)$ is basic.*

PROOF. We want to use the following special case of the characterization of \bar{v}_b , for certain $b \in G(L)$: suppose we are given an element $b \in G(L)$ such that for sufficiently divisible $s \in \mathbb{N}$, we may write in the semidirect product $G(L) \rtimes \langle \sigma \rangle$ the identity

$$(b\sigma)^s = (s\nu)(p) \sigma^s$$

for a rational B -dominant cocharacter $s\nu : \mathbf{G}_m \rightarrow Z(G)$ defined over \mathbb{Q}_p . Then in that case, b is basic and

$$\bar{v}_b = \frac{1}{s}(s\nu) \in X_*(T)_{\mathbb{Q}}^{\Gamma_p}.$$

This follows immediately from the general characterization of $\bar{\nu}_b$ given in [Ko85], §4.3.

This characterization applies to the element τ because of the identities (12.4.1) and (12.4.2). In fact we see that, in GL_n , the Newton point of τ is

$$\bar{\nu}_\tau = \left(\left(\frac{-d}{n}\right)^n\right) \in X_*(T(\mathrm{GL}_n))_{\mathbb{Q}}^{\Gamma_p},$$

which clearly factors through the center of G . In the Siegel case,

$$\bar{\nu}_\tau = \left(\left(\frac{-1}{2}\right)^{2n}\right) \in X_*(T(\mathrm{GSp}_{2n}))_{\mathbb{Q}}^{\Gamma_p}.$$

Thus τ is basic in each case. □

We get the following, which of course we already knew in the Siegel case, by Theorem 12.4.

COROLLARY 12.12. *Let Sh be a “fake” unitary or Siegel modular Shimura variety with level structure given by a standard parahoric subgroup K_p . Then the basic locus of Sh is nonempty.*

For (more detailed) information concerning the Newton stratification on some other Shimura varieties, the reader might consult the work of Andreatta-Goren [AG] and Goren-Oort [GOo].

13. The number of irreducible components in $Sh_{\mathbb{F}_p}$

Let Sh be a “fake” unitary or a Siegel modular variety with Iwahori-level structure as in §5. Recall that $\mathbf{M}_{\mathbb{F}_p}^{\mathrm{loc}}$ is a connected variety with an Iwahori-orbit stratification indexed by the finite subset $\mathrm{Adm}(\mu) \subset \widetilde{W}$, where $\mu = (0^{n-d}, (-1)^d)$, resp. $(0^n, (-1)^n)$. Its irreducible components are indexed by the maximal elements in this set, namely by the translation elements t_λ for λ belonging to the Weyl-orbit $W\mu$ of the coweight μ .

It is natural to hope that similar statements apply in some sense to Sh . The varieties $Sh_{\mathbb{F}_p}$ and $\widetilde{Sh}_{\mathbb{F}_p}$ are not geometrically connected (the number of connected components of $Sh_{\mathbb{F}_p}$ depends on the choice of subgroup $K^p \subset \mathbf{G}(\mathbb{A}_f^p)$; see below). Nevertheless the following two Lemmas (13.1 and 13.2) show that in every connected component of Sh , all the KR-strata are nonempty.

LEMMA 13.1 ([Ge], Prop. 1.3.2, in the Siegel case). *If Sh is either a “fake” unitary or a Siegel modular Shimura variety, and K_p is any standard parahoric subgroup, then the morphism $\psi : \widetilde{Sh} \rightarrow \mathbf{M}_{K_p}^{\mathrm{loc}}$ is surjective.*

PROOF. First we claim that it suffices to prove the lemma for $K_p^{\mathbf{a}}$, our standard Iwahori subgroup. Indeed, it is enough to observe that the following diagram commutes

$$\begin{array}{ccc} \widetilde{Sh}_{K_p^{\mathbf{a}}} & \xrightarrow{\psi} & \mathbf{M}^{\mathrm{loc}} \\ \downarrow & & \downarrow p \\ \widetilde{Sh}_{K_p} & \xrightarrow{\psi} & \mathbf{M}_{K_p}^{\mathrm{loc}}, \end{array}$$

and that the right vertical arrow p is surjective on the level of KR-strata: every Aut_{K_p} -orbit in $\mathbf{M}_{K_p}^{\text{loc}}$ contains an element in the image of p ; this follows from [KR], Prop. 9.3, 10.6. See also [Go3].

Next consider the Siegel case with Iwahori level structure, where this result is due to Genestier, loc. cit. We briefly recall his argument. It is easy to see that ψ is surjective on generic fibers, because it is Aut-equivariant, and the generic fiber of \mathbf{M}^{loc} is a single orbit under Aut. Because ψ is smooth, the complement of its image is an Aut-invariant, Zariski-closed subset of the special fiber of \mathbf{M}^{loc} . On the other hand, there is a unique closed (zero-dimensional) Aut-orbit (denoted here by τ^{-1}) in that fiber which belongs to the Zariski-closure of every other Aut-orbit, and one can show (by writing down an explicit chain of supersingular abelian varieties) that the point τ^{-1} belongs to the image of ψ . It follows that ψ is surjective.

In the “fake” unitary case with Iwahori-level structure, consider the element τ from §12.4 above. Then the element τ^{-1} indexes the unique zero-dimensional Aut-orbit in $\mathbf{M}^{\text{loc}} = M_{-w_0\mu}$. By Genestier’s argument, it is enough to prove that τ^{-1} belongs to the image of ψ , or equivalently, the stratum Sh_τ is nonempty. But we have proved in §12.4 that $[\delta] = [\tau]$ belongs to the image of $\delta_{K_p^0}$. Furthermore, it is obvious that $X_\tau(\tau)_{\tilde{K}_p^a} \neq \emptyset$, and hence by Proposition 12.6, we conclude that $Sh_\tau \cap \mathcal{S}_{[\tau]} \neq \emptyset$. (Note: Using the element $\tau \in \widetilde{W}(\text{GSp}_{2n})$, this argument applies just as well to the Siegel case, thus providing an alternative to Genestier’s step of finding an explicit chain of abelian varieties in the moduli problem which maps to τ^{-1} .) \square

Recall that Sh is not a connected scheme. In fact, for our two examples, the set of connected components of the geometric generic fiber $Sh_{\overline{\mathbb{Q}}_p}$ carries a simply transitive action by the finite abelian group

$$\pi_0 = \mathbb{Z}_{(p)}^+ \backslash (\mathbb{A}_f^p)^\times / c(K^p) = \mathbb{Q}^+ \backslash \mathbb{A}_f^\times / c(K^p)\mathbb{Z}_p^\times = \mathbb{Q}^\times \backslash \mathbb{A}^\times / c(\mathbf{K}K_\infty),$$

where $c : \mathbf{G} \rightarrow \mathbb{G}_m$ is the similitude homomorphism, K_∞ denotes the centralizer of h_0 in $\mathbf{G}(\mathbb{R})$, and the superscript $+$ designates the positive elements of the set; cf. [Del]. To see the groups above are actually isomorphic, use the fact that $c(K_\infty) \supset \mathbb{R}^+$ and $c(K_p) = \mathbb{Z}_p^\times$, as one can easily check for each of our two examples. Fixing isomorphisms $\mathbb{A}_f^p(1) = \mathbb{A}_f^p$ and $\overline{\mathbb{Q}}_p = \mathbb{C}$ once and for all, an element $(A_\bullet, \lambda, i, \bar{\eta}) \in Sh(\overline{\mathbb{Q}}_p)$ belongs to the connected component indexed by $a \in \pi_0$ if and only if the Weil-pairing $(\cdot, \cdot)_\lambda$ on $H_1(A, \mathbb{A}_f^p)$ pulls-back via $\bar{\eta}$ to the pairing $a(\cdot, \cdot)$ on $V \otimes \mathbb{A}_f^p$; see [H3] §2.

Let $\overline{\mathbb{Z}}_p \subset \overline{\mathbb{Q}}_p$ denote the subring of elements integral over \mathbb{Z}_p , and fix $a \in \pi_0$. Let Sh^0 denote the moduli space (over e.g. $\overline{\mathbb{Z}}_p$) of points $(A_\bullet, \lambda, i, \bar{\eta}) \in Sh$ such that

$$\bar{\eta}^*(\cdot, \cdot)_\lambda = a(\cdot, \cdot).$$

LEMMA 13.2. *Suppose K_p is any standard parahoric subgroup. Then the fibers of $Sh^0 \rightarrow \text{Spec}(\overline{\mathbb{Z}}_p)$ are connected. Furthermore, the morphism $\psi : \widetilde{Sh}^0 \rightarrow \mathbf{M}_{K_p}^{\text{loc}}$ is surjective.*

PROOF. By [Del], the generic fiber $Sh_{\overline{\mathbb{Q}}_p}^0$ is connected. In the “fake” unitary case, $Sh^0 \rightarrow \text{Spec}(\overline{\mathbb{Z}}_p)$ is proper and flat, and hence by the Zariski connectedness principle (comp. [Ha], Ex. III.11.4), the special fiber is also connected. In the

Siegel case $Sh^0 \rightarrow \text{Spec}(\overline{\mathbb{Z}}_p)$ is flat but not proper, so the same argument does not apply (but see [CF] IV.5.10 when K_p is maximal hyperspecial). However, the connectedness of the special fiber still holds and can be proved in an indirect way from the p -adic monodromy theorem of [CF]; for details see [Yu].

The statement regarding surjectivity follows from the proof of Lemma 13.1: the local model diagram does not “see” $\bar{\eta}$, and so we can arrange matters so that all constructions occur within Sh^0 and \widehat{Sh}^0 . \square

From now on we assume $K_p = K_p^{\mathbf{a}}$. The above lemma proves that in any connected component Sh^0 , all KR-strata Sh_w^0 are nonempty.

In fact, because the KR-stratum Sh_τ is zero-dimensional, the nonemptiness of $Sh_\tau^0 \cap \mathcal{S}_{[\tau]}$ proves slightly more. The following statement is in some sense the opposite extreme of the result of Genestier-Ngô in Corollary 8.3.

COROLLARY 13.3. *In the “fake” unitary or the Siegel case with $K_p = K_p^{\mathbf{a}}$, let Sh_τ^0 denote the the zero-dimensional KR-stratum in a connected component Sh^0 . Then Sh_τ^0 is nonempty and is contained in the basic locus of $Sh_{K_p^{\mathbf{a}}}$.*

How can we describe the irreducible components in $Sh_{\mathbb{F}_p}^0$? These are clearly just the closures of the irreducible components of the KR-strata $Sh_{t_\lambda}^0$, as λ ranges over the Weyl-orbit $W\mu$. A priori, each of these maximal KR-strata might be the (disjoint) union of several irreducible components, all having the same dimension.

COROLLARY 13.4. *In the Siegel or “fake” unitary case, $Sh_{\mathbb{F}_p}^0$ is equidimensional, and the number of irreducible components is at least $\#W\mu$.*

In the Siegel case, a much more precise statement has been established by C.-F. Yu [Yu], answering in the affirmative a question raised in [deJ].

THEOREM 13.5 ([Yu]). *In the Siegel case with $K_p = K_p^{\mathbf{a}}$, each maximal KR-stratum $Sh_{t_\lambda}^0$ is irreducible. Hence $Sh_{\mathbb{F}_p}^0$ has exactly 2^n irreducible components. An analogous statement holds for any standard parahoric subgroup K_p .*

It is reasonable to expect that similar methods will apply to the “fake” unitary case to prove that the number of irreducible components in $Sh_{\mathbb{F}_p}^0$ is exactly $\#W\mu$. In fact, it would be interesting to determine whether this last statement remains true for any PEL Shimura variety attached to a group whose p -adic completion is unramified.

14. Appendix: Summary of Dieudonné theory and de Rham and crystalline cohomology for abelian varieties

This summary is extracted from some standard references – [BBM], [Dem], [Fon], [Il], [MaMe], [Me], and [O] – as well as from [deJ].

14.1. de Rham cohomology and the Hodge filtration. To an abelian scheme $a : A \rightarrow S$ of relative dimension g is associated the de Rham complex $\Omega_{A/S}^\bullet$ of \mathcal{O}_A -modules. We define the de Rham cohomology sheaves

$$R^i a_*(\Omega_{A/S}^\bullet).$$

The first de Rham cohomology sheaf

$$R^1 a_*(\Omega_{A/S}^\bullet)$$

is a locally free \mathcal{O}_S -module of rank $2g$. If S is the spectrum of a Noetherian ring R , then

$$H^1_{DR}(A/S) := H^1(A, \Omega^{\bullet}_{A/S}) = \Gamma(S, R^1 a_*(\Omega^{\bullet}_{A/S}))$$

is a locally free R -module of rank $2g$.

The Hodge-de Rham spectral sequence degenerates at E_1 ([BBM], §2.5), yielding the exact sequence

$$0 \rightarrow R^0 a_*(\Omega^1_{A/S}) \rightarrow R^1 a_*(\Omega^{\bullet}_{A/S}) \rightarrow R^1 a_*(\mathcal{O}_A) \rightarrow 0.$$

We define $\omega_A := R^0 a_*(\Omega^1_{A/S})$, a locally free sub- \mathcal{O}_S -module of rank g . The term $R^1 a_*(\mathcal{O}_A)$ may be identified with $\text{Lie}(\widehat{A})$, the Lie algebra of the dual abelian scheme \widehat{A}/S . It is also locally free of rank g . Thus we have the *Hodge filtration* on de Rham cohomology

$$0 \rightarrow \omega_A \rightarrow R^1 a_*(\Omega^{\bullet}_{A/S}) \rightarrow \text{Lie}(\widehat{A}) \rightarrow 0.$$

Recall that in our formulation of the moduli problem defining $\text{Sh}(\mathbf{G}, h)_{K_p}$, the important determinant condition refers to the Lie algebra $\text{Lie}(A)$, and not to $\text{Lie}(\widehat{A})$. Because of this it is convenient (although not, strictly-speaking, necessary) to work with the *covariant* analogue $M(A)$ of $R^1 a_*(\Omega^{\bullet}_{A/S})$. To define it, recall that for any \mathcal{O}_S -module N , we define the dual \mathcal{O}_S -module N^\vee by

$$N^\vee := \mathcal{H}om_{\mathcal{O}_S}(N, \mathcal{O}_S).$$

Let $M(A) := (R^1 a_*(\Omega^{\bullet}_{A/S}))^\vee$ be the dual of de Rham cohomology. This is a locally free \mathcal{O}_S -module of rank $2g$. By the proposition below, we can identify $\omega_A^\vee = \text{Lie}(A)$ and so the *Hodge filtration* on $M(A)$ takes the form

$$0 \rightarrow \text{Lie}(\widehat{A})^\vee \rightarrow M(A) \rightarrow \text{Lie}(A) \rightarrow 0.$$

It is sometimes convenient to denote $\mathbb{D}(A)_S := R^1 a_*(\Omega_{A/S})$ (this notation refers to crystalline cohomology, see [BBM], [II]).

PROPOSITION 14.1 ([BBM], Prop. 5.1.10). *There is a commutative diagram whose vertical arrows are isomorphisms*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Lie}(\widehat{A})^\vee & \longrightarrow & \mathbb{D}(A)_S^\vee & \longrightarrow & \omega_A^\vee \longrightarrow 0 \\ & & \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ 0 & \longrightarrow & \omega_{\widehat{A}} & \longrightarrow & \mathbb{D}(\widehat{A})_S & \longrightarrow & \text{Lie}(A) \longrightarrow 0. \end{array}$$

14.2. Crystalline cohomology. Let $k_r = \mathbb{F}_{p^r}$ be a finite field with ring of Witt vectors $W(k_r)$. The fraction field L_r of $W(k_r)$ is an unramified extension of \mathbb{Q}_p and its Galois group is the cyclic group of order r generated by the Frobenius element $\sigma : x \mapsto x^p$; note also that σ acts on Witt vectors by the rule $\sigma(a_0, a_1, \dots) = (a_0^p, a_1^p, \dots)$.

Let A be an abelian variety over k_r of dimension g . We have the *integral isocrystal* associated to A/k_r , given by the data

$$\mathbb{D}(A) = (H^1_{\text{crys}}(A/W(k_r)), F, V).$$

Here the crystalline cohomology group $H^1_{\text{crys}}(A/W(k_r))$ is a free $W(k_r)$ -module of rank $2g$, equipped with a σ -linear endomorphism F (“Frobenius”) and the σ^{-1} -linear endomorphism V (“Verschiebung”) which induce bijections on

$H_{\text{crys}}^1(A/W(k_r)) \otimes_{W(k_r)} L_r$. We have the identity $FV = VF = p$ (by definition of V), hence the inclusions of $W(k_r)$ -lattices

$$H_{\text{crys}}^1(A/W(k_r)) \supset FH_{\text{crys}}^1(A/W(k_r)) \supset pH_{\text{crys}}^1(A/W(k_r))$$

(as well as the analogous inclusions for V replacing F).

The endomorphism F has the property that $F^r = \pi_A$ on H_{crys}^1 , where π_A denotes the absolute Frobenius morphism of A relative to the field of definition k_r (on projective coordinates x_i for A , π_A induces the map $x_i \mapsto x_i^{p^r}$).

14.3. Relation with Dieudonné theory. The crystalline cohomology of A/k_r is intimately connected to the (contravariant) Dieudonné module of the p -divisible group $A[p^\infty] := \varinjlim A[p^n]$, the union of the sub-groupschemes $A[p^n] = \ker(p^n : A \rightarrow A)$. Recall that the classical contravariant Dieudonné functor $G \mapsto D(G)$ establishes an exact anti-equivalence between the categories

$$\left\{ p\text{-divisible groups } G = \varinjlim G_n \text{ over } k_r \right\}$$

and

$$\left\{ \text{free } W(k_r)\text{-modules } M = \varprojlim M/p^n M, \text{ equipped with operators } F, V \right\};$$

see [Dem], [Fon]. Here F and V are σ - resp. σ^{-1} -linear, inducing bijections on $M \otimes_{W(k_r)} L_r$.

The crystalline cohomology of A/k_r , together with the operators F and V , is the same as the Dieudonné module of the p -divisible group $A[p^\infty]$, in the sense that there is a canonical isomorphism

$$(14.3.1) \quad H_{\text{crys}}^1(A/W(k_r)) \cong D(A[p^\infty])$$

which respects the endomorphisms F and V on both sides, cf. [BBM]. Moreover, we have the following identifications

$$(14.3.2) \quad \mathbb{D}(A)_{k_r} := H_{\text{crys}}^1(A/W(k_r)) \otimes_{W(k_r)} k_r \cong H_{DR}^1(A/k_r) \cong D(A[p]).$$

The second isomorphism is due to Oda [O]; see below. The first isomorphism is a standard fact ([BBM]), but can also be deduced via Oda's theorem by reducing equation (14.3.1) modulo p : the exactness of the functor D implies that $D(A[p]) = D(A[p^\infty])/pD(A[p^\infty]) = D(A[p^\infty]) \otimes_{W(k_r)} k_r$. In particular, the k_r -vector space $H_{DR}^1(A/k_r)$ inherits endomorphisms F and V (σ - resp. σ^{-1} -linear).

The theorem of Oda [O] includes as well the relation between the Hodge filtration on the de Rham cohomology of A and a suitable filtration on the isocrystal $\mathbb{D}(A)$.

THEOREM 14.2 ([O], Cor. 5.11). *There is a natural isomorphism $\psi : \mathbb{D}(A)_{k_r} \xrightarrow{\sim} H_{DR}^1(A/k_r)$, and under this isomorphism, $V\mathbb{D}(A)_{k_r}$ is taken to ω_A . In particular there is an exact sequence*

$$0 \rightarrow V\mathbb{D}(A) \rightarrow \mathbb{D}(A) \rightarrow \text{Lie}(\widehat{A}) \rightarrow 0.$$

14.4. Remarks on duality. We actually make use of this in a dual formulation. Define $H = H(A)$ to be the $W(k_r)$ -linear dual of the isocrystal $\mathbb{D}(A)$

$$(14.4.1) \quad H(A) = \text{Hom}_{W(k_r)}(\mathbb{D}(A), W(k_r)),$$

and define $H_{L_r} = H(A)_{L_r} = H(A) \otimes_{W(k_r)} L_r$. Letting $\langle \cdot, \cdot \rangle : H \times \mathbb{D}(A) \rightarrow W(k_r)$ denote the canonical pairing, we define σ - resp. σ^{-1} -linear injections F resp. V on H (they are bijective on H_{L_r}) by the formulae

$$(14.4.2) \quad \langle Fu, a \rangle = \sigma \langle u, Va \rangle$$

$$(14.4.3) \quad \langle Vu, a \rangle = \sigma^{-1} \langle u, Fa \rangle$$

for $u \in H$ and $a \in \mathbb{D}(A) = H_{\text{crys}}^1(A/W(k_r))$.

Of course $H(A)_{k_r} := H \otimes_{W(k_r)} k_r$ is the k_r -linear dual of $\mathbb{D}(A)_{k_r}$, hence

$$H(A)_{k_r} = M(A/k_r).$$

LEMMA 14.3. *Let $S = \text{Spec}(k_r)$. Equip $H = \mathbb{D}(A)^\vee$ with operators F, V as in (14.4.2). Then the isomorphism*

$$\mathbb{D}(A)_{k_r}^\vee \xrightarrow{\sim} \mathbb{D}(\widehat{A})_{k_r}$$

of Proposition 14.1 is an isomorphism of $W(k_r)[F, V]$ -modules.

PROOF. From [Dem], Theorem 8 (p. 71), for a p -divisible group G with Serre dual G' (loc. cit., p. 46) there is a duality pairing in the category of $W(k_r)[F, V]$ -modules

$$\langle \cdot, \cdot \rangle : D(G') \times D(G) \rightarrow W(k_r)(-1)$$

where $W(k_r)(-1)$ is the isocrystal with underlying space $W(k_r)$ and σ -linear endomorphism $p\sigma$. That is, we have the identity

$$\langle F_{G'}x, F_Gy \rangle = p\sigma \langle x, y \rangle,$$

or

$$\langle F_{G'}x, y \rangle = \sigma \langle x, pF_G^{-1}y \rangle = \sigma \langle x, V_Gy \rangle.$$

There is a canonical identification

$$(A[p^\infty])' = \widehat{A}[p^\infty].$$

Now from the above pairing with $G = A[p^\infty]$ and (14.3.1) we deduce a duality pairing of $W(k_r)[F, V]$ -modules

$$\mathbb{D}(\widehat{A}) \times \mathbb{D}(A) \rightarrow W(k_r)(-1),$$

which induces the isomorphism of Proposition 14.1. The lemma follows from these remarks. \square

Applying Oda's theorem (14.2) to \widehat{A} and invoking the above lemma gives $\text{Lie}(\widehat{A})^\vee = VM(A/k_r)$, thus there is an exact sequence

$$0 \rightarrow VH(A) \rightarrow H(A) \rightarrow \text{Lie}(A) \rightarrow 0.$$

Thus, we have

COROLLARY 14.4. *Let F, V be the σ - resp. σ^{-1} -linear endomorphisms of $H = H(A)$ defined in (14.4.2). There is a natural isomorphism*

$$\frac{V^{-1}H}{H} = \sigma(\text{Lie}(A)).$$

Moreover, on H_{L_r} we have the identity $V^{-r} = \pi_A^{-1}$. (Comp. [Ko92], §10, 16.)

References

- [AG] F. Andreatta, E.Z. Goren, *Geometry of Hilbert modular varieties over totally ramified primes*, IMRN **33**, (2003), 1785-1835.
- [BB] A. Beilinson, I.N. Bernstein, P. Deligne, *Faisceaux Pervers*, Astérisque **100**, (1981).
- [BBM] P. Berthelot, L. Breen, W. Messing, *Théorie de Dieudonné cristalline*, vol. 2, LNM **930**, Springer 1982.
- [BR] D. Blasius, J.D. Rogawski, *Zeta functions of Shimura varieties*, in: *Motives*, vol. 2, Proc. Symp. Pure Math. **55.2** (1994), 525-571.
- [Bo] A. Borel, *Automorphic L-functions*, In: *Automorphic Forms, Representations and L-functions*, Proc. Sympos. Pure Math., vol. **33**, part 2, Amer. Math. Soc., Providence, RI, 1979, pp. 27-61.
- [BT1] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. I*, Inst. Hautes Études Sci. Publ. Math. **41** (1972), 5-251.
- [BT2] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. II*, Inst. Hautes Études Sci. Publ. Math. **60** (1984), 5-184.
- [Ca] P. Cartier, *Representations of p-adic groups: a survey* In: *Automorphic Forms, Representations and L-functions*, Proc. Sympos. Pure Math., vol. **33**, part 1, Amer. Math. Soc., Providence, RI, 1979, pp. 111-155.
- [CF] C.-L. Chai, G. Faltings, *Degeneration of abelian varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Band 22, Springer-Verlag (1990).
- [deJ] A.J. de Jong, *The moduli spaces of principally polarized abelian varieties with $\Gamma_0(p)$ -level structure*, J. Alg. Geom. **2** (1993), 667-688.
- [SGA7 XIII] P. Deligne, *Le formalisme des cycles évanescents*, SGA7, Exp. XIII.
- [Del] P. Deligne, *Travaux de Shimura*, Sem. Bourbaki 389. LNM **244**, Springer 1971.
- [Del2] P. Deligne, *Variétés de Shimura: Interprétation modulaire, et techniques de construction de modèles canoniques*, Proc. Symp. Pure Math. **33** (1979), part 2, 247-290.
- [Weil2] P. Deligne, *La conjecture de Weil II*, Publ. Math. IHES **52**, 137-252 (1980)
- [Dem] M. Demazure, *Lectures on p-divisible groups*, Springer LNM **302** (1972), 98 pp.
- [DP] P. Deligne, G. Pappas, *Singularities des espaces de modules de Hilbert, en les caractéristiques divisant le discriminant*, Composition Math. **90** (1994), 59-79.
- [FM] L. Fargues, E. Mantovan *Variétés de Shimura, espaces de Rapoport-Zink et correspondances de Langlands locales*, Astérisque **291** (2004), 331 pp.
- [Fon] J.-M. Fontaine, *Groupes p-divisibles sur les corps locaux*, Astérisque **47-48**, (1977), 262 pp.
- [Ga] D. Gaitsgory, *Construction of central elements in the affine Hecke algebra via nearby cycles*, Invent. Math. **144** (2001), 253-280.
- [Ge] A. Genestier, *Un modèle semi-stable de la variété de Siegel de genre 3 avec structures de niveau de type $\Gamma_0(p)$* , Compositio Math. **123** (2000), no. 3, 303-328.
- [GN] A. Genestier, B.C. Ngô, *Alcoves et p-rang des variétés abéliennes*, Ann. Inst. Fourier **52** (2002), 1665-1680.
- [GOo] E.Z. Goren, F. Oort, *Stratifications of Hilbert modular varieties*, J. Alg. Geom. **9**, (2000), no. 1, 111-154.
- [Go1] U. Görtz, *On the flatness of models of certain Shimura varieties of PEL-type*, Math. Ann. **321** (2001), no. 3, 689-727.
- [Go2] U. Görtz, *On the flatness of local models for the symplectic group*, Adv. Math. **176**, (2003), no.1, 89-115.
- [Go3] U. Görtz, *Topological flatness of local models in the ramified case*, math.AG/0205050.
- [Go4] U. Görtz, *Computing the alternating trace of Frobenius on the sheaves of nearby cycles on local models for GL_4 and GL_5* , J. of Algebra **278** (2004), no.1, 148-172.
- [GH] U. Görtz, T. Haines, *The Jordan-Hölder series for nearby cycles on some Shimura varieties and affine flag varieties*, preprint (2003). math.AG/0402143.
- [SGA7 I] A. Grothendieck, *Groupes de monodromie en géométrie algébrique*, Exp. I, (rédigé par P. Deligne), Springer LNM **288** (1972), 1-24.
- [H1] T. Haines, *The combinatorics of Bernstein functions*, Trans. Amer. Math. Soc. **353** (2001), 1251-1278.
- [H2] T. Haines, *Test functions for Shimura varieties: the Drinfeld case*, Duke Math. J. **106** (2001), no. 1, 19-40.

- [H3] T. Haines, *On connected components of Shimura varieties*, *Canad. J. Math.*, vol. **54** (2), 2002, pp. 352-395.
- [HKP] T. Haines, R. Kottwitz, and A. Prasad, *Iwahori-Hecke Algebras*, math.RT/0309168.
- [HN1] T. Haines, B.C. Ngô, *Nearby cycles for local models of some Shimura varieties*, *Compositio Math.* **133** (2002), no. 2, 117–150.
- [HN2] T. Haines, B.C. Ngô, *Alcoves associated to special fibers of local models*, *Amer. J. Math.* **124** (2002), no. 6, 1125–1152.
- [HN3] T. Haines, B.C. Ngô, *On the semi-simple local L -factors for some simple Shimura varieties*, in preparation.
- [HP] T. Haines, A. Pettet, *Formulae relating the Bernstein and Iwahori-Matsumoto presentations of an affine Hecke algebra*, *J. Algebra* **252** (2002), no. 1, 127–149.
- [HT] M. Harris, R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, *Annals of Math. Studies*, Princeton Univ. Press, 276 pp + vi, 2001.
- [Ha] R. Hartshorne, *Algebraic Geometry*, GTM **52**, Springer-Verlag, 1977.
- [Il] L. Illusie, *Report on crystalline cohomology*, *Proc. Symp. Pure Math.* **29** (1975), 459-478.
- [Ko84] R. Kottwitz, *Shimura varieties and twisted orbital integrals*, *Math. Ann.* **269** (1984), 287-300.
- [Ko85] R. Kottwitz, *Isocrystals with additional structure*, *Compositio Math.* **56** (1985), 201-220.
- [Ko90] R. Kottwitz, *Shimura varieties and λ -adic representations*, In: *Automorphic Forms, Shimura Varieties and L -functions*, part 1. *Perspectives in Mathematics 10*. Academic Press, 1990, pp. 161-209.
- [Ko92] R. Kottwitz, *Points on some Shimura varieties over finite fields*, *J. Amer. Math. Soc.* **5** (1992), 373-444.
- [Ko92b] R. Kottwitz, *On the λ -adic representations associated to some simple Shimura varieties*, *Invent. Math.* **108** (1992), 653-665.
- [Ko97] R. Kottwitz, *Isocrystals with additional structure. II*, *Compositio Math.* **109**, (1997), 255-339.
- [Ko03] R. Kottwitz, *On the Hodge-Newton decomposition for split groups*, *IMRN* **26** (2003), 1433-1447.
- [KR] R. Kottwitz, M. Rapoport, *Minuscule alcoves for GL_n and GSp_{2n}* , *Manuscripta Math.* **102** (2000), no. 4, 403–428.
- [KW] R. Kiehl, R. Weissauer, *Weil conjectures, Perverse sheaves, and ℓ -adic Fourier transform*, *Springer Erg. Math. 3 Folg.*, **42** (2001).
- [Lu] G. Lusztig, *Affine Hecke algebras and their graded versions*, *J. Amer. Math. Soc.* **2** (1989), 599–635.
- [Ma] B. Mazur, *Notes on étale cohomology of number fields*, *Ann. Sci. Ecole Norm. Sup.* **6** (1973), 521-556.
- [MaMe] B. Mazur, W. Messing, *Universal extensions and one dimensional crystalline cohomology*, *Springer LNM* **370** (1974), 134 pp.
- [Me] W. Messing, *The crystals associated to Barsotti-Tate groups*, *LNM* **264**, Springer 1962.
- [Mi] J.S. Milne, *Abelian Varieties*, p. 103-150, In: *Arithmetic Geometry*, ed. G. Cornell, J.H. Silverman, Springer-Verlag, 1986.
- [Mu] D. Mumford, *Abelian varieties*, Oxford University Press, 1974.
- [O] T. Oda, *The first de Rham cohomology group and Dieudonné modules*, *Ann. Scient. École Norm. Sup.*, 4 ser., t. 2, (1969), 63-135.
- [Oo] F. Oort, *Newton polygon strata in the moduli space of abelian varieties*, In: *Moduli of abelian varieties* (Texel Island, 1999), 417-440, *Progr. Math.*, **195**, Birkhäuser, Basel, 2001.
- [P] G. Pappas, *On the arithmetic moduli schemes of PEL Shimura varieties*, *J. Alg. Geom.* **9** (2000), 577-605.
- [PR1] G. Pappas, M. Rapoport, *Local models in the ramified case I. The EL-case*, *J. Alg. Geom.* **12** (2003), 107-145.
- [PR2] G. Pappas, M. Rapoport, *Local models in the ramified case II. Splitting models*, *Duke Math. J.* **127** (2005), no. 2, 193-250.
- [R1] M. Rapoport, *On the bad reduction of Shimura varieties*. *Automorphic Forms, Shimura Varieties, and L -functions*, Vol. II, (Ann Arbor, MI, 1988), 253-321, *Perspec. Math* **11**, Academic Press, Boston, MA, 1990.
- [R2] M. Rapoport, *A Guide to the reduction modulo p of Shimura varieties*, preprint (2002). math.AG/0205022.

- [RR] M. Rapoport, M. Richartz, *On the classification and specialization of F -isocrystals with additional structure*, *Compositio Math.* **103** (1996), 153-181.
- [RZ82] M. Rapoport, Th. Zink, *Über die lokale Zetafunktion von Shimuravarietäten. Monodromiefiltration und verschwindende Zyklen in ungleicher Charakteristik*, *Invent. Math.* **68** (1982), no.1, 21-101.
- [RZ] M. Rapoport, T. Zink, *Period domains for p -divisible groups*, *Ann. of Math. Stud.* **141**, Princeton University Press (1996).
- [Re1] H. Reimann, *The semi-simple zeta function of quaternionic Shimura varieties*, *Lecture Notes in Mathematics* **1657**, Springer-Verlag, Berlin, 1997, viii + 143 pp.
- [Re2] H. Reimann, *Reduction of quaternionic Shimura varieties at parahoric levels*, *Manuscripta Math.* **107**, (2002), no. 3, 355-390.
- [Re3] H. Reimann, *Cohomology of quaternionic Shimura varieties at parahoric levels*, *Manuscripta Math.* **107**, (2002), no. 3, 391-407.
- [Ta] J. Tate, *Number theoretic background*, In: *Automorphic Forms, Representations and L-functions*, *Proc. Sympos. Pure Math.*, vol. **33**, part 2, Amer. Math. Soc., Providence, RI, 1979, pp. 3 -26.
- [T] J. Tits, *Reductive groups over local fields*, In: *Automorphic Forms, Representations, and L-functions*, ed. A. Borel, W. Casselman, *Proc. Symp. Pure Math.* **33** (1979), Part I, 29-69.
- [W1] T. Wedhorn, *Ordinariness in good reductions of Shimura varieties of PEL-type*, *Ann. Sci. École Norm. Sup. (IV)*, **32**, (1999), no. 5, 575-618.
- [W2] T. Wedhorn, *The local Langlands correspondence for $GL(n)$ over p -adic fields*, *math.AG/0011210 v2*.
- [Wi] J.-P. Wintenberger, *Existence de F -cristaux avec structures supplémentaires*, *Adv. Math.* **190** (2005), no. 1, 196-224.
- [Yu] C.-F. Yu, *Irreducibility of the Siegel moduli spaces with parahoric level structure*, *IMRN*, **48** (2004), 2593-2597.
- [Z] Th. Zink, *Über die schlechte Reduktion einiger Shimuramannigfaltigkeiten*, *Compositio Math.* **45** (1981), 15-107.

MATHEMATICS DEPARTMENT, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742-4015
E-mail address: tjh@math.umd.edu