

TOPOLOGICAL TAMENESS OF MARGULIS SPACETIMES

SUHYOUNG CHOI AND WILLIAM GOLDMAN

Dedicated to the memory of Bill Thurston, our mentor

ABSTRACT. We show that Margulis spacetimes without parabolic holonomy are topologically tame. A Margulis spacetime is the quotient of the 3-dimensional Minkowski space by a free proper isometric action of the free group of rank ≥ 2 . We will use our particular point of view that the Margulis spacetime is a manifold-with-boundary with an \mathbb{RP}^3 -structure in an essential way. The basic tools are a bordification by a closed surface with an \mathbb{RP}^2 -structure and a free holonomy group, the work of Goldman, Labourie, and Margulis on geodesics in the Margulis spacetimes and 3-manifold topology.

CONTENTS

1. Introduction	2
2. Preliminaries	7
2.1. \mathbb{RP}^n -structures on manifolds	7
2.2. Projective \mathbf{S}^3	8
2.3. A matrix lemma	11
2.4. The space of properly convex sets and the joins	12
3. The oriented Lorentzian space and projective boosts	14
3.1. Oriented Lorentzian vector spaces	14
3.2. Null half-planes	14
3.3. Affine boosts	17

Date: June 10, 2013.

1991 Mathematics Subject Classification. Primary 57M50; Secondary 53C15, 53C50, 53A20.

Key words and phrases. Lorentzian manifolds, handlebody, geodesic flow, \mathbb{RP}^2 -structure, proper action.

The first author was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MEST) (No.2010-0027001). The second author gratefully acknowledges partial support from National Science Foundation grants DMS-070781 and DMS-1065965.

3.3.1. Oriented axes	17
3.4. Projective boosts	18
4. \mathbb{RP}^2 -surfaces and hyperbolic surfaces and their dynamics	19
4.1. \mathbb{RP}^2 -structures on surfaces	19
4.2. Complete hyperbolic surfaces and Fuchsian groups	20
4.3. The dynamic of geodesic flows on hyperbolic surfaces	21
5. \mathbb{RP}^2 -Schottky uniformizations	23
5.1. The construction of the Γ -invariant domain	23
5.2. The \mathbb{RP}^2 -surface	26
6. The dynamic core of a Margulis spacetime	29
6.1. Margulis invariants	29
6.2. Neutralized sections	30
6.3. Lifting the neutralized section to the coverings	32
6.4. Mapping the convex core	32
7. Γ acts properly on $E \cup \tilde{\Sigma}$.	33
7.1. The action of projective-boost automorphisms	34
7.2. The proper discontinuity of the action of Γ .	37
8. The proof of tameness.	46
9. Asymptotics of Margulis spacetimes	49
9.1. Supporting hyperplanes	49
9.2. Domains	52
10. Geometric finiteness	55
References	58

1. INTRODUCTION

A *complete affine manifold* is a quotient M of an n -dimensional real affine space E by a discrete group Γ of affine transformations acting properly. Equivalently, M is a manifold with a geodesically flat torsionfree affine connection. The purpose of this paper is topologically classify complete affine manifolds in dimension $n = 3$ when the fundamental group $\pi_1(M)$ is finitely generated and has no parabolic holonomy.

Those affine 3-manifolds where the fundamental groups are solvable were classified by Fried-Goldman [29] in the twentieth century. These include the case when the manifolds are compact.

In all other cases, M admits a parallel Lorentzian structure, and $\pi_1(M^3) \cong \Gamma$ is a free group. The existence of these manifolds was first demonstrated by Margulis [41], and later clarified by Drumm [22] using the geometric construction of *crooked planes*. In particular Drumm's

examples are homeomorphic to the interiors of *closed solid handlebodies*, that is, compact 3-manifolds-with-boundary obtained by attaching 1-handles to a closed 3-ball. We call such a manifold M *topologically tame*.

Charette, Drumm, Goldman, and Labourie [22], [23], [24] made contributions to understanding such spacetimes with parallel Lorentzian structures culminating in recent work [35] and [34]. The complete classification of such spacetimes is continuing with work of Charette, Drumm, Goldman, Margulis and Minsky [12], [13], [36].

The theory of compactifying open manifolds goes back to Browder, Levine and Livesay [10] and Siebenmann [47]. For 3-manifolds, Tucker [50] and Scott and Tucker [49] made an initial clarification. See also Meyers [44]. As a note, we state that complete hyperbolic 3-manifolds with finitely generated fundamental groups were shown to be tame by Agol, Calegari, and Gabai. See Bowditch [9] for details on this very large topic. We won't elaborate on this as the methods are completely different. Our proof is closer to the proof of the tameness of geometrically finite hyperbolic 3-manifolds due to Thurston [48]. (This is written up nicely by Epstein and Marden [26] and Marden [42] even earlier. See also Bonahon [7].)

Lately after the first version of this paper was uploaded, there is another independent approach to this question by J. Danciger, F. Guéritaud, and F. Kassel. Their approach uses the deformation of constant curvature Lorentzian manifolds and yields other results such as the deformability of Margulis spacetimes to anti-de Sitter spaces. Compared to their approach, our approach mostly concentrates on \mathbb{RP}^2 -structures and the compactification of the spacetime. We also obtain some asymptotic properties of the action.

Let V denote a 3-dimensional real vector space with an inner product of signature 2, 1 and a fixed orientation. The spacetime E is an affine space with underlying vector space V . A *Lorentzian isometry* is an automorphism of E preserving the Lorentzian inner product. Denote the group of orientation-preserving Lorentzian isometries by $\text{Isom}^+(E)$. The projectivization $\mathbb{P}(V)$ is defined as the quotient space

$$V - \{O\} / \sim \text{ where } v \sim w \text{ if and only if } v = sw \text{ for } s \in \mathbb{R} - \{0\}.$$

Of course, $\mathbb{P}(V)$ is identical with the real projective 2-space \mathbb{RP}^2 . Recall that we embed the hyperboloid model \mathbb{H}^2 of the hyperbolic plane to the disk that is the interior of the conic in \mathbb{RP}^2 determined by null vectors. Here $\text{PGL}(3, \mathbb{R})$ acts faithfully on \mathbb{RP}^2 as the group of collineations.

Define $\mathbb{S}(V)$ as

$$V - \{O\} / \sim \text{ where } v \sim w \text{ if and only if } v = sw \text{ for } s > 0.$$

Then the image \mathbb{S}_+ of the space of future timelike vectors identifies with the hyperbolic 2-plane \mathbb{H}^2 , which is basically the Beltrami-Klein model of the hyperbolic plane. Let \mathbb{S}_- denote the subspace of $\mathbb{S}(V)$ corresponding to past timelike vectors. The group of orientation preserving linear maps of V is denoted by $\mathrm{SO}(2, 1)$, a Lie group with two components one of which is the identity component $\mathrm{SO}(2, 1)^\circ$. The linear group $\mathrm{SO}(2, 1)^\circ$ acts faithfully on $\mathbb{H}^2 = \mathbb{S}_+$ as the orientation-preserving isometry group and $\mathrm{SO}(2, 1)$ acts so on $\mathbb{S}_+ \cup \mathbb{S}_-$ and acts on $\mathbb{S}(V)$ projectively where the action is induced from that on V .

Since $\mathrm{SO}(2, 1)$ injects to $\mathrm{PGL}(3, \mathbb{R})$ under projection $\mathrm{GL}(3, \mathbb{R}) \rightarrow \mathrm{PGL}(3, \mathbb{R})$, we consider that $\mathrm{SO}(2, 1)$ acts on $\mathbb{S}_+ = \mathbb{H}^2$ a subset of \mathbb{RP}^2 as well in a *projective manner*. Note that $\mathrm{SO}(2, 1)^\circ$ acts on \mathbb{S}_+ *honestly* as a subset of $\mathbb{S}(V)$.

Recall that there is a homomorphism

$$\mathcal{L} : \mathrm{Isom}^+(E) \rightarrow \mathrm{SO}(2, 1)$$

given by taking the linear part acting on $\mathbb{S}(V)$. We also denote by

$$\mathcal{L}' : \mathrm{Isom}^+(E) \xrightarrow{\mathcal{L}} \mathrm{SO}(2, 1) \hookrightarrow \mathrm{PGL}(3, \mathbb{R})$$

with the image acting on \mathbb{RP}^2 .

A Lorentzian transformation is said to be *parabolic* if its linear part has only eigenvalues 1 but is not identity. One that is neither parabolic nor identity has eigenvalues $\lambda, 1/\lambda, 1$. Such an element is called a *hyperbolic transformation* if $|\lambda| > 1$. A *positive hyperbolic transformation* is a hyperbolic transformation with only positive eigenvalues. Sometimes a positive hyperbolic transformation is called a *linear boost*.

Suppose that Γ is a finitely generated Lorentzian isometry group acting freely and properly on E . We assume that Γ is not amenable (i.e., not solvable). Since E/Γ is aspherical, the fundamental group Γ is torsion-free. Γ injects under \mathcal{L} to $\mathcal{L}(\Gamma) \subset \mathrm{SO}(2, 1)$ acting properly discontinuously and freely on $\mathbb{S}_+ \cup \mathbb{S}_-$. By Mess [43], Γ is a free group of rank ≥ 2 . (See [29] where this nontrivial fact was first discovered and Section 3 of [34] for details.) Thus, we restrict ourselves to the group actions of free groups of rank ≥ 2 in this paper. Then E/Γ is said to be a *Margulis spacetime*, and an element of Γ is said to be a *holonomy*.

Note that $\mathbb{H}^2/\mathcal{L}'(\Gamma)$ may be not orientable if $\mathcal{L}(\Gamma)$ is not a subset of $\mathrm{SO}(2, 1)^\circ$. However, $\mathbb{S}_+ \cup \mathbb{S}_-/\mathcal{L}'(\Gamma)$ is an orientable surface double-covering $\mathbb{H}^2/\mathcal{L}'(\Gamma)$. (See [13] for details.) The hyperbolic surface

$\mathbb{H}^2/\mathcal{L}'(\Gamma)$ is *geometrically finite* if Γ is nonelementary and has no parabolics.

In this paper we will be concerned with the cases without parabolic elements in Γ . The following conditions are equivalent conditions for a finitely generated Lorentzian isometry group Γ .

- E/Γ is a Margulis spacetime without parabolics holonomies.
- Γ is proper affine deformation of a geometrically finite Fuchsian group; i.e., $\mathbb{H}^2/\mathcal{L}'(\Gamma)$ is geometrically finite and Γ acts properly on E . (See [35] for definition.)
- Γ is a nonamenable Lorentzian isometry group acting properly on E without parabolic holonomy.
- Γ is a free Lorentzian isometry group of rank ≥ 2 acting properly and freely on E without parabolic holonomy.

In this case, we say that Γ is a *Margulis group without parabolics*.

It also follows from above that elements of Γ are either hyperbolic or parabolic or identity. (See [35]). Notice that the definitions of hyperbolic, positive hyperbolic, and parabolic for elements of Γ are the same as ones for $\mathcal{L}'(\Gamma)$ as a group of isometries of \mathbb{S}_+ . (If a hyperbolic element in $\text{SO}(2, 1)$ has a negative eigenvalue, then they should be $-\lambda, -1/\lambda, 1, 1$.)

An \mathbb{RP}^n -structure on a manifold is given by an atlas of charts to \mathbb{RP}^n with projective transition maps. Such geometric structures were first considered by Kuiper, Koszul, Benzécri and others in the 1950s and 1960s. Further developments can be followed in Goldman [33], Choi and Goldman [18], and Cooper, Long, and Thistlethwaite [19] and [20].

Theorem 1.1. *Let Γ be a finitely generated free group of rank ≥ 2 in $\text{SO}(2, 1)$ acting on $\mathbb{S}_+ \cup \mathbb{S}_-$ properly discontinuously and freely but without parabolics. Let $S := S(V)$ be the sphere of directions in V , which is a model for the universal covering space $\widehat{\mathbb{RP}^2} \rightarrow \mathbb{RP}^2$. Then there exists a Γ -invariant open domain $D \subset S(V)$ such that D/Γ is a closed surface Σ . Such a domain is unique up to the antipodal map A .*

These surfaces have the \mathbb{RP}^2 -structures on closed surfaces of genus g , $g \geq 2$, discovered by Goldman [31] in the late 1970s. Although their developing maps from the universal cover $\tilde{\Sigma}$ of some such surface Σ are not covering-spaces onto their images in \mathbb{RP}^2 , when lifted to the double-covering space $S(V) := \widehat{\mathbb{RP}^2} \rightarrow \mathbb{RP}^2$, the developing maps (remarkably) are covering-spaces onto open domains. The surface is a quotient of a domain in S by a group of projective automorphisms, an \mathbb{RP}^2 -analog of the standard Schottky uniformization of a Riemann surface as a \mathbb{CP}^1 -manifold as observed by Goldman.

An n -dimensional open manifold is said to be *tame* if it is homeomorphic to the interior of a compact n -manifold with boundary. A *handlebody* is a 3-dimensional manifold obtained from a 3-ball B^3 by attaching 1-handles to 3-dimensional balls. The topology of handlebodies and related objects form a central topic in the 3-dimensional manifold topology. See Hempel [38] for a thorough survey of this field of mathematics developed in the 20th century.

Theorem 1.2. *Let E be the standard Lorentzian space with the standard norm given by $x_0^1 - x_1^2 - x_2^2$, and let Γ be a Margulis group without parabolics. Then E/Γ is homeomorphic to the interior of a handlebody of genus r for the rank r of Γ as a free group.*

We define in this paper that a complete geodesic on a quotient space of a hyperbolic space or a Lorentzian space is *nonwandering* if it is bounded in both directions; that is, the closure of the forward part is compact and so is that of the backward part; i.e., the α -limit and the ω -limit are both nonempty and compact. This is the same as the term “recurrent” in [34] and [14], which does not agree with the common usage in the dynamical system theory whereas the recurrence set they discuss is the same as the nonwandering set as in Eberlein [25] and Katok and Hasselblatt [39]: the reason is that these are geodesics with both endpoints in the limit sets and the set of these geodesics coincide with the nonwandering set in [25] by Corollary 3.8 there.

The work of Goldman–Labourie–Margulis [35] on nonwandering geodesics on M implies tameness: A Margulis spacetime is geometrically finite in some sense since every closed geodesic is in a bounded neighborhood, which will explain our particular viewpoints on this in Section 9.

In Section 2, we review facts on hyperbolic surfaces, Fuchsian group actions, \mathbb{RP}^n -structures and geometry that we need. We compactify E by a real projective 2-sphere \mathbb{S} that is just the space of directions in E .

In Section 3, we review facts about the Lorentzian space E and affine boosts. The affine boosts will be extended to projective boosts in \mathbf{S}^3 .

In Section 4, we review real projective structures on surfaces and hyperbolic surfaces. We also discuss the dynamic of the geodesic flows on the unit tangent bundles of hyperbolic surfaces.

In Section 5, we prove Theorem 1.1. This will be done by finding an open domain $\tilde{\Sigma}$ in the boundary \mathbb{S} of E where Γ acts properly discontinuously and freely. $\tilde{\Sigma}/\Gamma$ is shown to be a closed \mathbb{RP}^2 -surface Σ . Such a surface was constructed by Goldman [31]; however, we realize it as the quotient of an open domain in the sphere \mathbb{S} , which is a union of two disks and infinitely many strips joining the two.

Now let Γ be as assumed in Theorem 1.2. In Section 6, we recall the work of Goldman, Labourie, and Margulis [35] on nonwandering geodesics on the Margulis spacetime. Their work in fact shows that all nonwandering spacelike geodesics in a Margulis spacetime are in a uniformly bounded part.

In Section 7, we prove the proper discontinuity of the action of the group Γ on $E \cup \tilde{\Sigma}$. We first show that for a fixed Lorentzian isometry g , we can push every compact convex subset of $E \cup \tilde{\Sigma}$ into an ϵ -neighborhood of the “attracting” segment of g in \mathbb{S} by g^i for sufficiently large i . (This is the key model of our proof.)

We next show that for any compact set $K \subset E \cup \tilde{\Sigma}$, there are only finitely many $g \in \Gamma$ so that $g(K) \cap K \neq \emptyset$. Suppose not. We find a sequence $\{g_i\}$ so that $g_i(K) \cap K \neq \emptyset$. Using the work [35], we can find an infinite sequence $\{g_i\}$ that behaves almost like $\{g^i\}$ in the dynamical sense as $i \rightarrow \infty$ up to small changes of stable and unstable planes. In other words, we can find a uniformly bounded sequence of coordinates to normalize g_i into fixed forms and these coordinates are uniformly convergent.

In Section 8, we prove the tameness of E/Γ , i.e., Theorem 1.2, using the classical 3-manifold topology as developed by Dehn and so on.

In Section 9, we show that there exists an invariant set of hyperplanes asymptotic to the limit set Λ of Γ on $\text{Cl}(\mathbb{S}_+)$. This gives us two disjoint Γ -invariant closed convex domains in E asymptotic to Λ and its antipodal set Λ_- in \mathbb{S} . (This is a generalization of a result of Mess [43].)

In Section 10, we also define “geometric finiteness” in this setting and show that there exists a topological core of a Margulis spacetime that contains all spacelike nonwandering geodesics.

This work was started from the question of Goldman on the \mathbb{RP}^2 -bordification of the Margulis spacetime during the Newton Institute Workshop “Representations of Surface Groups and Higgs Bundles” held in Oxford, March 14-18, 2011. The authors thank the Newton Institute and the Institut Henri Poincaré where parts of this work were carried out. Finally, we thank Bill Thurston without whose teaching we could not have accomplished many of the things in this paper.

2. PRELIMINARIES

2.1. \mathbb{RP}^n -structures on manifolds. The projective space \mathbb{RP}^n is given by $\mathbb{P}(\mathbb{R}^{n+1})$. The general linear group is sent

$$\text{GL}(n+1, \mathbb{R}) \rightarrow \text{PGL}(n+1, \mathbb{R}) := \text{GL}(n+1, \mathbb{R}) / \sim$$

where two linear maps L_1 and L_2 are equivalent if $L_1 = sL_2$ for $s \in \mathbb{R} - \{0\}$. A map of an open subset of \mathbb{RP}^n to one of \mathbb{RP}^n is *projective* if it is a restriction of an element of $\mathrm{PGL}(n+1, \mathbb{R})$.

More generally, a nonzero linear map $L : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ becomes a *projective endomorphism*

$$\hat{L} : \mathbb{P}(\mathbb{R}^{n+1}) - \mathbb{P}(N) \rightarrow \mathbb{P}(\mathbb{R}^{n+1})$$

where N is the kernel of L and $\mathbb{P}(N)$ is the image of $N - \{O\}$ in $\mathbb{P}(\mathbb{R}^{n+1})$ defined by $\hat{L}([\vec{v}]) = [L(\vec{v})]$. (See Section 5.1 of Benzécri [6].) The space of projective endomorphisms equals the projectivization $\mathbb{P}(\mathrm{Mat}_{n+1}(\mathbb{R}))$ of the linear space $\mathrm{Mat}_{n+1}(\mathbb{R})$ of all linear maps $\mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$. The space $\mathbb{P}(\mathrm{Mat}_{n+1}(\mathbb{R}))$ forms a compactification of the group $\mathrm{PGL}(n+1, \mathbb{R})$ as observed first by Kuiper [40] and developed by Benzécri [6].

The projective geometry is given by a pair $(\mathbb{RP}^n, \mathrm{PGL}(n+1, \mathbb{R}))$. An \mathbb{RP}^n -structure on a manifold M is given by a maximal atlas of charts to \mathbb{RP}^n with projective transition maps. The manifold M with an \mathbb{RP}^n -structure is said to be an \mathbb{RP}^n -manifold. A *projective map* for two \mathbb{RP}^n -manifolds M and N is a map $f : M \rightarrow N$ so that $\phi \circ f \circ \psi^{-1}$ is projective whenever it is defined where ϕ is a chart for N and ψ^{-1} is a local inverse of a chart for M .

An \mathbb{RP}^n -structure on a manifold M gives us an immersion $\mathrm{dev} : \tilde{M} \rightarrow \mathbb{RP}^n$ that is equivariant with respect to a homomorphism

$$h : \pi_1(M) \rightarrow \mathrm{PGL}(n+1, \mathbb{R}) : \text{ that is,}$$

$$\mathrm{dev} \circ \gamma = h(\gamma) \circ \mathrm{dev} \text{ for every } \gamma \in \pi_1(M).$$

Here, dev is called a *developing map* and h is called the *holonomy homomorphism*. (dev, h) is only determined up to an action of $\mathrm{PGL}(n+1, \mathbb{R})$ where

$$g(\mathrm{dev}, h(\cdot)) = (g \circ \mathrm{dev}, gh(\cdot)g^{-1}) \text{ for } g \in \mathrm{PGL}(n+1, \mathbb{R}).$$

Conversely, such a pair (dev, h) will give us an \mathbb{RP}^n -structure since dev gives us charts that have projective transition maps. (See [16] for details.)

2.2. Projective \mathbf{S}^3 . We can identify \mathbf{S}^n with $\mathbb{S}(\mathbb{R}^{n+1})$. There exists a quotient map $q_1 : \mathbf{S}^n \rightarrow \mathbb{RP}^n$ given by sending a unit vector to its equivalence class. This is a double-covering map, and it induces an \mathbb{RP}^n -structure on \mathbf{S}^n . The group $\mathrm{Aut}(\mathbf{S}^n)$ of projective automorphisms is isomorphic to the group $\mathrm{SL}_\pm(n+1, \mathbb{R})$ of linear maps of determinant equal to ± 1 with the quotient homomorphism $\mathrm{SL}_\pm(n+1, \mathbb{R}) \rightarrow \mathrm{PGL}(n+1, \mathbb{R})$ with the kernel $\pm \mathbf{I}$. It will be convenient to think of these as matrices in $\mathrm{SL}_\pm(n+1, \mathbb{R})$.

This space is again an open dense subspace of $\mathbb{S}(\text{Mat}_n(\mathbb{R}))$. We call the element of $\mathbb{S}(\text{Mat}_n(\mathbb{R}))$ a *projective endomorphism* of \mathbf{S}^n . Each element corresponds to a map $\hat{L} : \mathbb{S}(\mathbb{R}^{n+1}) - \mathbb{S}(N) \rightarrow \mathbb{S}(\mathbb{R}^{n+1})$ where N is the kernel of a linear map L in $\text{Mat}_n(\mathbb{R})$ and $\mathbb{S}(N)$ is its image in \mathbf{S}^n and defined by

$$\hat{L}([\vec{v}]) = [L(\vec{v})]$$

for $\vec{v} \in \mathbb{R}^{n+1}$.

For an open domain \mathcal{D} in \mathbf{S}^n and a discrete group Γ in $\text{Aut}(\mathbf{S}^n)$ acting on it properly discontinuously and freely, \mathcal{D}/Γ has an \mathbb{RP}^n -structure since $q_1|_{\mathcal{D}}$ gives an immersion and the homomorphism

$$\Gamma \subset \text{Aut}(\mathbf{S}^n) \rightarrow \text{PGL}(n+1, \mathbb{R})$$

gives the associated holonomy homomorphism.

In this paper, we will study objects on $\mathbf{S}^3 = \mathbb{S}(\mathbb{R}^4) = \mathbb{S}(\mathbb{R} \oplus \mathbb{R}^3)$. A *homogeneous coordinate system* of \mathbf{S}^3 is given by setting a point p of \mathbf{S}^3 be given coordinates $[x : y : z : t]$ where $\frac{(x,y,z,t)}{|(x,y,z,t)|}$ represents p as a unit vector in \mathbf{S}^3 . Again

$$[x : y : z : t] \sim [x' : y' : z' : t'] \text{ iff } (x, y, z, t) = \lambda(x', y', z', t') \text{ for } \lambda > 0.$$

There is the antipodal map

$$\mathcal{A} : \mathbf{S}^3 \rightarrow \mathbf{S}^3 \text{ given by } [v] \rightarrow [-v].$$

Given a subset or a point K of \mathbf{S}^3 , we denote by K_- the set of antipodal points of points of K .

A *subspace* of \mathbf{S}^3 is a subset defined by a system of linear equations in \mathbb{R}^4 . A *line* is a 1-dimensional subspace. A singleton is not a subspace. A *projective geodesic* is an arc in a line. We will be using the standard Riemannian metric on \mathbf{S}^3 to be denoted by $\mathbf{d}_{\mathbf{S}^3}$. Notice that the geodesic structure of \mathbf{S}^3 is the same as the projective one. Thus, we will use the same term “geodesics” for both types of geodesics. Geodesics and totally geodesic subspaces are all subsets of lines or subspaces in \mathbf{S}^3 . A projective automorphism sends these to themselves, while it does not preserve any metric. A pair of *antipodal points* on \mathbf{S}^3 is the equivalence class $[v]$ of a nonzero vector v and one $[-v]$ of $-v$. A segment connecting two antipodal points is precisely a segment of $\mathbf{d}_{\mathbf{S}^3}$ -length π .

Here, we denote by $\overline{pqp_-}$ the unique closed segment in \mathbf{S}^3 with endpoints p and p_- passing through q not equal to p or p_- . By \overline{pq} , we denote the unique closed segment in \mathbf{S}^3 with endpoints p and q not containing p_-, q_- provided $p \neq q$ and $p \neq q_-$. The notation \overline{pq}° or $\overline{pqp_-}^\circ$ indicates the interior of the segment.

The Lorentzian space E is the affine space \mathbb{R}^3 equipped with a non-degenerate bilinear form \mathbf{B} of signature $1, 1, -1$. If we choose the origin in E , we obtain the Lorentzian vector space V .

One can think of \mathbb{R}^3 to be identical with E and the complement of a subspace of codimension-one in \mathbb{RP}^3 , so called the complete affine subspace. $\mathbb{R}^3 = \mathsf{E}$ lifts to a subspace of \mathbf{S}^3 double-covering \mathbb{RP}^3 . The closure of the lifted E is a standard 3-dimensional hemisphere \mathcal{H} , and E identifies with the open hemisphere \mathcal{H}° . The boundary \mathbf{S}^2 of E is the subspace of codimension-one identifiable with the projective 2-sphere \mathbb{S} of directions in \mathbb{R}^3 . (We identify E with the open affine space given by $t > 0$ in \mathbf{S}^3 from now on for convenience.)

The sphere \mathbb{S} corresponds to the hyperplane given by $t = 0$ and the origin of E is given by $[0 : 0 : 0 : 1]$, and by choosing O . For this system of coordinates, the point $(x, y, z) \in \mathsf{E}$ is given coordinates $[x : y : z : 1]$.

\mathbb{S} also has a homogeneous coordinate system $[x, y, z]$ assigned to a vector p in \mathbb{S} of unit length 1 if $\frac{(x, y, z)}{\|(x, y, z)\|} = p$. In the larger coordinate system $[x : y : z]$ is identical with $[x : y : z : 0]$.

To summarize, the subsets of \mathbf{S}^3 indentified in this paper are:

$$\begin{aligned}\mathbf{S}^3 &= \{[x : y : z : t] | x, y, z, t \in \mathbb{R}, (x, y, z, t) \neq (0, 0, 0, 0)\} \\ \mathcal{H} &= \{[x : y : z : t] | x, y, z, t \in \mathbb{R}, t \geq 0, (x, y, z, t) \neq (0, 0, 0, 0)\} \\ \mathbb{R}^3 = \mathsf{E} = \mathcal{H}^\circ &= \{[x, y, z, t] | x, y, z, t > 0 \in \mathbb{R}\} \\ &= \{[x : y : z : 1] | x, y, z \in \mathbb{R}\} \\ \mathbb{S} &= \{[x : y : z : 0] | x, y, z \in \mathbb{R}, (x, y, z, t) \neq (0, 0, 0, 0)\}.\end{aligned}$$

We will also be using a fixed Euclidean metric d_E on E compatible with the affine coordinate system. It is of great importance that projective geodesics, spherical geodesics, and Euclidean geodesics on E are the same ones up to parametrizations.

The group $\text{Isom}^+(\mathsf{E})$ of orientation-preserving Lorentzian transformations of E is in the group $\text{Aff}^+(\mathsf{E})$ of orientation-preserving affine transformations of E . $\text{Aff}^+(\mathsf{E})$ identifies with the subgroup of $\text{Aut}(\mathbf{S}^3)$ of orientation-preserving elements acting on $\mathbb{R}^3 = \mathsf{E}$, preserving \mathbb{S} , since any affine transformation of \mathbb{R}^3 extends to a projective automorphism of \mathbb{RP}^3 and hence an automorphism of \mathbf{S}^3 analytically. Often such a transformation is represented by

$$\begin{pmatrix} A & \vec{v} \\ 0 & 1 \end{pmatrix} \in \text{GL}(3, \mathbb{R})$$

a representative matrix in $\text{Aut}(\mathbf{S}^3)$ with A a nonsingular 2×2 -matrix in $\text{SO}(2, 1)$ and requiring the last diagonal diagonal entry 1. It is represented as an affine transformation

$$\vec{x} \mapsto A\vec{x} + \vec{v} \text{ for } \vec{x} \in E.$$

The elements of respective groups are called *projective automorphisms* of *Lorentzian* type and ones of *affine* type or just *Lorentzian isometries* or *affine transformations*. A *bi-Lipschitz map* of \mathbf{S}^3 is a homeomorphism $f : \mathbf{S}^3 \rightarrow \mathbf{S}^3$ so that

$$C^{-1}d_{\mathbf{S}^3}(x, y) \leq d_{\mathbf{S}^3}(f(x), f(y)) \leq Cd_{\mathbf{S}^3}(x, y) \text{ for } x, y \in \mathbf{S}^3$$

where $C > 0$ is independent of x, y . Elements of $\text{Aut}(\mathbf{S}^3)$ including ones extending the Lorentzian isometries are bi-Lipschitz maps of \mathbf{S}^3 .

Finally, for the purpose of drawing figures only, we map E to the unit 3-ball in \mathbb{R}^3 by the map

$$[x : y : z : 1] \rightarrow \frac{(x, y, z)}{\sqrt{x^2 + y^2 + z^2 + 1}}.$$

This will identify \mathbb{S} with the sphere of radius 1 in \mathbb{R}^3 and as a codimension-one totally geodesic subspace of \mathbf{S}^3 , represented stereographically.

We will denote by $\text{bd}A$ or $\text{bd}_X A$ the topological boundary of a subset A of some topological space X and by $\text{int}A$ or $\text{int}_X A$ the topological interior. If A is a topological manifold with boundary, we will denote the manifold boundary by ∂A and we will denote the manifold interior by A° .

2.3. A matrix lemma. We denote the standard vector by

$$\vec{e}_1 = (1, 0, 0, 0), \vec{e}_2 = (0, 1, 0, 0), \vec{e}_3 = (0, 0, 1, 0), \vec{e}_4 = (0, 0, 0, 1),$$

and denote

$$\begin{aligned} \mathbf{e}_1 &= [\vec{e}_1], \mathbf{e}_2 = [\vec{e}_2], \mathbf{e}_3 = [\vec{e}_3], \mathbf{e}_4 = [\vec{e}_4], \\ \mathbf{e}_{1-} &= [-\vec{e}_1], \mathbf{e}_{2-} = [-\vec{e}_2], \mathbf{e}_{3-} = [-\vec{e}_3], \mathbf{e}_{4-} = [-\vec{e}_4]. \end{aligned}$$

Lemma 2.1 (Uniform convergence). *Let \mathbf{v}_i^j for $j = 1, 2, 3, 4$ be a sequence of points of \mathbf{S}^3 . Suppose that $\mathbf{v}_i^j \rightarrow \mathbf{v}_\infty^j$ for each j and mutually distinct points $\mathbf{v}_\infty^1, \dots, \mathbf{v}_\infty^4$ formed by independent vectors in \mathbb{R}^4 . Then we can choose a sequence h_i of elements of $\text{Aut}(\mathbf{S}^3)$ so that*

- $h_i(\mathbf{v}_i^j) = \mathbf{e}_j$,
- h_i and h_i^{-1} are represented by uniformly bounded matrices,
- the sequence $\{h_i\}$ converges to h_∞ uniformly for h_∞ and
- $\{h_i^{-1}\}$ converges to h_∞^{-1} uniformly both under C^s -topology for every $s \geq 0$.

Proof. Let \vec{v}_i^j represent the unit norm vector corresponding to v_i^j . We find $\hat{h}_i \in \text{Aut}(\mathbf{S}^3)$ so that $\hat{h}_i(\mathbf{e}_j) = v_i^j$ for each i . We set $M_i(\vec{e}_i) = \vec{v}_i^j$; that is, the column vectors of M_i equals $\vec{v}_i^1, \vec{v}_i^2, \vec{v}_i^3, \vec{v}_i^4$. Then clearly, the sequence $\{M_i\}$ converges to a 4×4 -matrix M_∞ of nonzero determinant. Therefore, the sequence $\{M_i^{-1}\}$ of inverse matrices converges to M_∞^{-1} .

We let h_i be the projective automorphism in $\text{Aut}(\mathbf{S}^3)$ induced by M_i^{-1} , let h_∞ be the one corresponding to M_∞^{-1} , and this satisfies the properties we need. \square

2.4. The space of properly convex sets and the joins. Let $\mathbf{d}_{\mathbf{S}^3}\text{-diam}(A)$ denote the diameter of the set, that is, the supremum of the set of distances between every pair of points of a subset A of \mathbf{S}^n . Given a pair of points or subsets A and B in \mathbf{S}^n , we define the infimum distance

$$\mathbf{d}_{\mathbf{S}^3}(A, B) = \inf\{\mathbf{d}_{\mathbf{S}^3}(x, y) | x \in A, y \in B\},$$

and the supremum distance

$$\mathbf{d}_{\mathbf{S}^3}^s(A, B) = \sup\{\mathbf{d}_{\mathbf{S}^3}(x, y) | x \in A, y \in B\} = \mathbf{d}_{\mathbf{S}^3}\text{-diam}(A \cup B).$$

If $\mathbf{d}_{\mathbf{S}^3}(A, B) > \epsilon$ for $\epsilon > 0$, then we say that A is *bounded away* from B by ϵ or conversely.

An ϵ -neighborhood $N_\epsilon(A)$ for a number $\epsilon > 0$ of a subset A of \mathbf{S}^3 is the set of points of $\mathbf{d}_{\mathbf{S}^3}$ -distances less than ϵ from some points of A . We define the geometric Hausdorff distance $\mathbf{d}_{\mathbf{S}^3}^H(A, B)$ between two compact subsets A and B of \mathbf{S}^3 to be

$$\inf\{\epsilon \geq 0 | B \subset N_\epsilon(A) \text{ and } A \subset N_\epsilon(B)\}.$$

A sequence of compact sets $\{K_i\}$ in \mathbf{S}^3 converges or converges geometrically to a compact subset K if for every $\epsilon > 0$, there exists N so that for $i > N$, $\mathbf{d}_{\mathbf{S}^3}^H(K, K_i) < \epsilon$ or equivalently

$$K \subset N_\epsilon(K_i) \text{ and } K_i \subset N_\epsilon(K).$$

We will simply write $\{K_i\} \rightarrow K$ for K_i and K compact subsets of \mathbf{S}^3 .

The following are commonly known facts with elementary proofs.

Proposition 2.2.

- The space of compact subsets of a compact Hausdorff space with this metric is compact Hausdorff also and hence every sequence has a convergent subsequence.
- given a convergent sequence $\{K_i\}$ of compact subsets of \mathbf{S}^3 , if a sequence $\{J_i\}$ is such that $J_i \subset K_i$, then any geometric limit of a subsequence of $\{J_i\}$ is a subset of the geometric limit of $\{K_i\}$.

- If we have $\{K_i^j\} \rightarrow K^j$ for each $j = 1, \dots, m$, then

$$\bigcup_{j=1}^n K_i^j \rightarrow \bigcup_{j=1}^n K^j$$

holds.

Note that

$$\mathbf{d}_{\mathbf{S}^3}(A, B) \leq \mathbf{d}_{\mathbf{S}^3}^H(A, B) \leq \mathbf{d}_{\mathbf{S}^3}^s(A, B),$$

and three could be all distinct.

Suppose that two compact subsets A and B of \mathbf{S}^3 have no pairs of points $x \in A, y \in B$ so that x and y are antipodal, i.e., $A \cap \mathcal{A}(B) = \emptyset$. Then the *join* $J(A, B)$ is the union of all segments of $\mathbf{d}_{\mathbf{S}^3}$ -lengths $< \pi$ with single endpoints in A and the other ones in B . In set theoretic terms, we let

$$(1) \quad J(A, B) = q_1(\{c_1 \vec{u}_1 + c_2 \vec{u}_2 \mid c_1, c_2 \geq 0, \vec{u}_1 \in q_1^{-1}(A), \vec{u}_2 \in q_1^{-1}(B)\}).$$

When A is a great circle in \mathbf{S}^3 and B is a point $\notin A$, every segment from B to A is of $\mathbf{d}_{\mathbf{S}^3}$ -length $< \pi - \epsilon$ for a fixed $\epsilon > 0$. In this case, $J(A, B)$ is a *2-dimensional hemisphere*, a totally geodesic disk with geodesic boundary in \mathbf{S}^3 , a convex set.

If A and B are compact and has no antipodal pair of a point of A and a point B , then $J(A, B)$ is well-defined and is also compact as we can easily show that $J(A, B)$ is closed.

If A and B are convex and $A \cup B$ has no pair of antipodal points, then $J(A, B)$ is defined and also properly convex. The proof follows by reverting back to \mathbb{R}^4 by (1). Also, for a projective automorphism $g \in \text{Aut}(\mathbf{S}^3)$, we obtain

$$gJ(A, B) = J(g(A), g(B)).$$

Lemma 2.3. *Let A_i and B_i be a sequence of properly convex compact subsets of \mathbf{S}^3 geometrically converging to properly convex compact sets A and B respectively. Suppose that $A \cap \mathcal{A}(B) = \emptyset$. Then $J_i = J(A_i, B_i)$ is defined for $i > I_0$ for sufficiently large I_0 and the sequence $\{J_i\}_{i>I_0}$ converges to the join J of A and B .*

Proof. Clearly, A_i and B_i have no antipodal pair of points for sufficiently large i as $A_i \subset N_\epsilon(A), B_i \subset N_\epsilon(B)$ for sufficiently large i . One can parameterize each geodesic segment connecting a point of A_i to that of B_i by arclength in $\mathbf{d}_{\mathbf{S}^3}$. From this, the lemma follows. \square

3. THE ORIENTED LORENTZIAN SPACE AND PROJECTIVE BOOSTS

3.1. Oriented Lorentzian vector spaces. Let (V, \cdot, Det) denote a three-dimensional *oriented Lorentzian* \mathbb{R} -vector space. That is, $V \cong \mathbb{R}^3$ and is given a symmetric bilinear form

$$\mathbf{B} : V \times V \longrightarrow \mathbb{R}$$

defined by $(v, u) \mapsto v \cdot u$ of index 1 and a nondegenerate alternating trilinear form

$$V \times V \times V \longrightarrow \mathbb{R}$$

defined by $(v, u, w) \mapsto \text{Det}(v, u, w)$. Its automorphism group is the special orthogonal group $\text{SO}(2, 1)$.

The Lorentzian structure divides S into three open domains S_+, S_0 , and S_- separated by two conics $\text{bd}_S S_+$ and $\text{bd}_S S_-$. Let $S_+ \subset S$ denote the set of future timelike directions; its boundary $\text{bd}_S S_+$ consists of future null directions. Similarly, let $S_- \subset S$ denote the set of past timelike directions with boundary $\text{bd}_S S_-$ consisting of past null directions. Both S_+ and S_- are cells, interchanged by \mathcal{A} . The set S_0 of spacelike directions forms an \mathcal{A} -invariant annulus bounded by the disjoint union $\text{bd}_S S_+ \coprod \text{bd}_S S_-$.

Recall that S_+ is the Beltrami-Klein model of the hyperbolic plane where $\text{SO}(2, 1)^\circ$ acts as the orientation-preserving isometry group. Here the metric geodesics are precisely the projective geodesics and vice versa. (We will use the same term “geodesics” for both types of geodesics for ones in S_+ and later S_- .) The geodesics in S_+ are straight arcs and $\text{bd}_S S_+$ forms the ideal boundary of S_+ . For a finitely generated discrete subgroup Γ in $\text{SO}(2, 1)^\circ$, S_+/Γ has a complete hyperbolic structure as well as an \mathbb{RP}^2 -structure with the compatible geodesic structure.

Alternatively, we can use S_- as the model and S_-/Γ has a complete hyperbolic structure as well as an \mathbb{RP}^2 -structure. Since S_+ and S_- are properly convex, these are examples of properly convex \mathbb{RP}^2 -surfaces.

3.2. Null half-planes. Let \mathcal{N} denote the *nullcone* in V , that is, the collection of all vectors $v \in V$ with $v \cdot v = 0$. Its projectivization $\mathbb{P}(\mathcal{N} - \{O\})$ consists of all *null lines* in V . If $v \in \mathcal{N} - \{O\}$, then its orthogonal complement v^\perp is a *null plane* which contains the line $\mathbb{R}v$. The line $\mathbb{R}v$ separates v^\perp into two half-planes.

The orientation on V determines an $\text{SO}(2, 1)$ -invariant way to uniquely associate a component of $v^\perp - \mathbb{R}v$ to $\mathbb{R}v$ as follows. Since $v \in \mathcal{N}$ holds, its direction $[v]$ lies in either $\text{bd}_S S_+$ or $\text{bd}_S S_-$. Choose an arbitrary element u of S_+ or S_- respectively, so that the directions of v and u both lie in $\text{Cl}(S_+)$ or $\text{Cl}(S_-)$ respectively. (For example $u = (0, 0, \pm 1)$ would

be sufficient.) Define the *null half-plane* $\mathcal{W}(\mathbf{v})$ (or the *wing*) associated to \mathbf{v} as:

$$\mathcal{W}(\mathbf{v}) := \{\mathbf{w} \in \mathbf{v}^\perp \mid \text{Det}(\mathbf{v}, \mathbf{w}, \mathbf{u}) > 0\}.$$

Since the Lorentzian product $\mathbf{u} \times \mathbf{v}$ is in $\mathcal{W}(\mathbf{v})$, the directions in the wing satisfy the right-hand rule. Since $\mathcal{W}(\mathbf{v}) = \mathcal{W}(\mathcal{A}(\mathbf{v}))$ holds, the null half-plane $\mathcal{W}(\mathbf{v})$ depends only on $\mathbb{R}\mathbf{v}$. The corresponding set of directions is the open arc

$$\varepsilon(\mathbf{v}) := [\mathcal{W}(\mathbf{v})]$$

in \mathbb{S}_0 joining $[\mathbf{v}]$ to $\mathcal{A}[\mathbf{v}] = [-\mathbf{v}]$. Since \mathbf{v}^\perp is tangent to \mathcal{N} , the arc $\varepsilon(\mathbf{v})$ is tangent to $\text{bd}_{\mathbb{S}}\mathbb{S}_+$. The orientation of $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ as the boundary orientation of \mathbb{S}_+ agrees with the orientation of $\varepsilon(\mathbf{v})$ away from $[\mathbf{v}]$.

The corresponding map $[\mathbf{v}] \mapsto \varepsilon(\mathbf{v})$ is a $\text{SO}(2, 1)$ -equivariant map

$$\text{bd}_{\mathbb{S}}\mathbb{S}_+ \rightarrow \mathcal{S}$$

where \mathcal{S} denotes the set of half-arcs of form $\varepsilon(\mathbf{v})$ for $\mathbf{v} \in \text{bd}_{\mathbb{S}}\mathbb{S}_+$. The arcs $\varepsilon([\mathbf{v}])$ for $\mathbf{v} \in \text{bd}_{\mathbb{S}}\mathbb{S}_+$ foliate \mathbb{S}_0 obtained from an \mathbf{S}^1 -action fixing $[\pm 1 : 0 : 0]$ where \mathbf{S}^1 is a subgroup of $\text{SO}(2, 1)$. Let us call the foliation \mathcal{F} .

Hence \mathbb{S}_0 has a $\text{SO}(2, 1)$ -equivariant quotient map

$$\Pi : \mathbb{S}_0 \rightarrow \mathbb{P}(\mathcal{N} - \{O\}) \cong \mathbf{S}^1$$

and $\varepsilon([\mathbf{v}]) = \Pi^{-1}([\mathbf{v}])$ for each $\mathbf{v} \in \mathcal{N} - \{O\}$.

Consider a future-pointing null vector of Euclidean length $\sqrt{2}$:

$$\mathbf{n}_\theta := \begin{bmatrix} \cos(\theta) \\ \sin(\theta) \\ 1 \end{bmatrix} \quad \text{for } \theta \in [0, 2\pi).$$

Then $\mathcal{W}(\mathbf{n}_\theta)$ consists of all

$$\mathbf{p}_\theta(t, s) := t\mathbf{n}_\theta + s \begin{bmatrix} -\sin(\theta) \\ \cos(\theta) \\ 0 \end{bmatrix}$$

where $t \in \mathbb{R}$ and $s > 0$. The arc $c_\theta := \varepsilon(\mathbf{n}_\theta)$ on the sphere of directions \mathbb{S} is parametrized by unit vectors

$$\frac{1}{\sqrt{2}}\mathbf{p}_\theta(t, \sqrt{1-t^2})$$

as $-1 < t < 1$. (As an element of \mathbf{S}^3 , its homogeneous coordinates equal

$$[\frac{1}{\sqrt{2}}\mathbf{p}_\theta(t, \sqrt{1-t^2}) : 0]$$

where $\frac{1}{\sqrt{2}}\mathbf{p}_\theta(t, \sqrt{1-t^2})$ is used as the last three coordinates.)




FIGURE 1. The tangent geodesics to disks \mathbb{S}_+ and \mathbb{S}_- in the unit sphere S imbedded in \mathbb{R}^3 .




FIGURE 2. The tangent geodesics to disks \mathbb{S}_+ and \mathbb{S}_- in the stereographically projected S from $(0, 0, -1)$. Radial arcs are geodesics. The inner circle represents the boundary of \mathbb{S}_+ . The arcs of form $\varepsilon(x)$ for $x \in \text{bd}_{\mathbb{S}}\mathbb{S}_+$ are leaves of the foliation \mathcal{F} on \mathbb{S}_0 .

The other component of $v^\perp - \mathbb{R}v$ could also be used as a wing $\mathcal{W}(v)$, but for the opposite orientation on V . Alternatively, these *negatively oriented null half-planes* are the images of the positively oriented null half-planes under the antipodal map \mathcal{A} (which reverses orientation in dimension three). This phenomenon appears in the theory of crooked planes, where one must choose a class of orientations for crooked planes, as well as the Margulis invariant, where the choice of orientation plays a crucial role.

Since we have no need to consider negatively oriented null half-planes, we henceforth restrict our attention to positively oriented null half-planes in this paper.

3.3. Affine boosts.

Definition 3.1. An isometry of E is an (*affine*) *boost* if its linear part is a positive hyperbolic element of $\mathrm{SO}(2, 1)$ and it acts freely on E .

Suppose that $\gamma \in \mathrm{Isom}^+(E)$ is an affine boost. Then γ preserves a unique line which we denote $\mathrm{Axis}(\gamma)$. (Margulis calls this line C_γ .) Furthermore $\mathrm{Axis}(\gamma)$ is spacelike and the restriction of γ to $\mathrm{Axis}(\gamma)$ is a nonzero translation. In a suitable coordinate system, we can take $\mathrm{Axis}(\gamma)$ to be the z -axis, in which case γ is the affine transformation:

$$p \xrightarrow{\gamma} \begin{bmatrix} e^l & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-l} \end{bmatrix} p + \begin{bmatrix} 0 \\ \alpha \\ 0 \end{bmatrix}$$

where

$$\begin{bmatrix} e^l & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-l} \end{bmatrix}$$

is the linear part $\mathcal{L}(\gamma)$ of γ .

3.3.1. Oriented axes. As observed by Margulis, the axis of an affine boost γ admits a *canonical* orientation, induced by the orientation of E .

Let $g = \mathcal{L}(\gamma) \in \mathrm{SO}(2, 1)$ be positive hyperbolic. Then since g is an isometry, and unimodular, the eigenvalues of g are $\lambda, 1, \lambda^{-1}$ where $\lambda > 1$. The eigenspaces for eigenvalues λ, λ^{-1} are null (since g is an isometry); choose respective eigenvectors $v_+(g), v_-(g) \in V$ with the same causal character (that is, they are either both future-pointing or both past-pointing). The 1-eigenspace $\mathrm{Fix}(g) \subset V$ is spacelike and orthogonal to $v_+(g)$ and $v_-(g)$.

Definition 3.2. The *neutral eigenvector* of g is the unique unit-spacelike vector $v_0(g)$ such that $\{v_+(g), v_0(g), v_-(g)\}$ is a positively oriented basis of V . (See [24] for details.)

Since $\mathrm{Axis}(\gamma)$ is parallel to $\mathrm{Fix}(\mathcal{L}(g))$, the neutral eigenvector $\mathcal{L}(g)^0$ defines an orientation on $\mathrm{Axis}(\gamma)$. Denote the corresponding *oriented geodesic* by $\widehat{\mathrm{Axis}}(\gamma)$.

The weak-stable plane $W^{wu}(\gamma) \subset E$ is the affine subspace containing $\mathrm{Axis}(\gamma)$ and parallel to $v_-(g)$. Note that $\gamma^n|W^{wu}(\gamma)$ acts on a point p of $W^{wu}(\gamma)$ so that

$\mathbf{d}_{\mathbf{S}^3}(g^n(p), [v_-(g) : 0]) \rightarrow 0$ and $\mathbf{d}_{\mathbf{S}^3}(g^n(p), \mathrm{Axis}(\gamma)) \rightarrow 0$ as $n \rightarrow \infty$.

(Notice $W^{wu}(\gamma)$ is a null-space.)

3.4. Projective boosts.

Definition 3.3. A *projective boost* is a collineation of \mathbf{S}^3 defined by the direct sum of a positive hyperbolic element of $\mathrm{SL}(3, \mathbb{R})$:

$$\begin{bmatrix} e^l & 0 \\ 0 & e^{-l} \end{bmatrix}$$

and a nontrivial unipotent element of $\mathrm{SL}(3, \mathbb{R})$:

$$\begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix},$$

where $\alpha \neq 0$, such that the three fixed points lie on the ideal plane \mathbb{S}_∞ .

Thus a projective boost is defined by the 4×4 matrix where the ideal plane subspace is defined by the fourth coordinate:

$$\begin{bmatrix} e^l & 0 & 0 & 0 \\ 0 & 1 & 0 & \alpha \\ 0 & 0 & e^{-l} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

The three fixed points on \mathbb{P}_∞ are:

$$x^+ := [1 : 0 : 0 : 0], \quad x^0 := [0 : 1 : 0 : 0], \quad x^- := [0 : 0 : 1 : 0]$$

in homogeneous coordinates.

Such a collineation induces the affine isometry of E defined by:

$$v \xrightarrow{\gamma} \begin{bmatrix} e^l & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-l} \end{bmatrix} v + \begin{bmatrix} 0 \\ \alpha \\ 0 \end{bmatrix}$$

The subspace invariant under the unipotent direct summand defines a projective line, which we call the *spine*:

$$\sigma := [0 : * : 0 : *]$$

that contains the axis.

Thus, any conjugate element in $\mathrm{Aut}(\mathbf{S}^3)$ is also said to be a *projective boost*. We denote by $x^+(\gamma), x^0(\gamma), x^-(\gamma)$, and $\sigma(\gamma)$ the points going to x^+, x^0, x^- and the subspace σ under the coordinate change of γ to make its matrix into the above form. The span $\mathcal{W}^{wu}(\gamma)$ of $x^-(\gamma) \cup \sigma(\gamma)$ is the subspace of \mathbf{S}^3 extending the weak-stable plane $W^{wu}(\gamma) \subset E$. Points on $W^{wu}(\gamma) \subset E$ has $x^0(\gamma)$ as a limit point along the line $\sigma(\gamma) \setminus \{x^0(\gamma)\}$. (Here $\sigma(\gamma)$ is the subspace spanned by $\mathrm{Axis}(\gamma)$ in E .)

Let $\{\gamma_n\}$ be a sequence of projective boosts in $\mathrm{Aut}(\mathbf{S}^3)$. Since $\mathbb{S}(\mathrm{Mat}_4(\mathbb{R}))$ is compact, we can extract a convergent subsequence always. Suppose that $\lim_{n \rightarrow +\infty} \gamma^n$ is a projective endomorphism γ_∞ in $\mathbb{S}(\mathrm{Mat}_4(\mathbb{R}))$ of

rank one in $\mathbb{S}(\text{Mat}_4(\mathbb{R}))$. (One could have a rank 2 limit also in general.) Then the undefined hyperplane of γ_∞ equals a subspace \mathcal{W}_∞^{wu} of codimension-one and whose image equals a pair of points x_∞^+ and x_∞^- where

$$\mathcal{W}^{wu}(\gamma_n) \rightarrow \mathcal{W}_\infty^{wu}, x^+(\gamma_n) \rightarrow x_\infty^+.$$

More generally, if $\{\gamma_n\}$ is a sequence in $\text{Aut}(\mathbf{S}^3)$ converging to a projective endomorphism γ_∞ in $\mathbb{S}(\text{Mat}_4(\mathbb{R}))$ of rank 1, we can choose a subsequence of $\{\gamma_n\}$ so that the corresponding subsequence of the attracting fixed points $\{x^+(\gamma_n)\}$ converges to the image $I(\gamma_\infty)$ and the corresponding subsequence of the extended weak-stable subspaces $\mathcal{W}^{wu}(\gamma_n)$ converges to the undefined subspace $U(\gamma_\infty)$. (Note that we can have a limit of rank 2 if we do not have $\{\mathbf{v}(\gamma_i)/\lambda(\gamma_i)\} \rightarrow 0$.)

4. \mathbb{RP}^2 -SURFACES AND HYPERBOLIC SURFACES AND THEIR DYNAMICS

4.1. \mathbb{RP}^2 -structures on surfaces. Let ϑ correspond to an element A of $\text{SL}(3, \mathbb{R})$ diagonalizable with three distinct positive eigenvalues. Let ϑ' be the element of $\text{Aut}(\mathbf{S}^2)$ induced by A . Then ϑ' has six fixed points a, a_-, r, r_-, s, s_- and three great circles l_1, l_2 , and l_3 so that

$$\{a, r, a_-, r_-\} \subset l_1, \{r, s, r_-, s_-\} \subset l_2, \{a, s, a_-, s_-\} \subset l_3.$$

There are eight invariant open triangles that are components of $\mathbf{S}^2 - l_1 \cup l_2 \cup l_3$. Let Δ be one with vertices a, r, s . Then

$$(\Delta \cup \overline{ar}^o \cup \overline{as}^o) / \langle \vartheta' \rangle, (\Delta \cup \overline{ar}^o \cup \overline{rs}^o) / \langle \vartheta' \rangle$$

are both examples of compact annuli. The \mathbb{RP}^2 -surface projectively diffeomorphic to these are said to be an *elementary annulus*.

Let Δ' be an adjacent triangle sharing \overline{ar}^o with Δ in the boundary. Then

$$(\Delta \cup \Delta' \cup \overline{ar}^o \cup \overline{as}^o \cup \overline{as_-}^o) / \langle \vartheta' \rangle, (\Delta \cup \Delta' \cup \overline{ar}^o \cup \overline{rs}^o \cup \overline{rs_-}^o) / \langle \vartheta' \rangle$$

are examples of compact annuli again. The \mathbb{RP}^2 -surfaces projectively diffeomorphic to these are said to be π -annuli. They are the union of two elementary annuli meeting at a boundary component. (See [16] and [17] for more details.)

Let \tilde{S} be the universal cover of an \mathbb{RP}^2 -surface S with a developing map $\text{dev} : \tilde{S} \rightarrow \mathbb{RP}^2$ and a holonomy homomorphism h . For a circle \mathbf{S}^1 , a *closed geodesic* $c : \mathbf{S}^1 \rightarrow S$ is a closed curve where $\text{dev} \circ \tilde{c}$ is a straight arc in \mathbb{RP}^2 for a lift $\tilde{c} : \mathbb{R} \rightarrow \tilde{S}$ of c . A closed geodesic is *principal* if for a lift \tilde{c} to the universal cover \tilde{S} , $\text{dev} \circ \tilde{c}$ is an embedding to a straight arc connecting an attracting and a repelling fixed point of $h(\gamma)$ for a deck transformation γ of \tilde{S} satisfying $\tilde{c}(t + 2\pi) = \gamma \circ \tilde{c}(t)$. (Here γ

is said to be the *corresponding* deck transformation of \tilde{c} and c , and it exists uniquely.)

A *properly convex* domain in \mathbb{RP}^2 is a bounded convex domain of an affine subspace in \mathbb{RP}^2 . An \mathbb{RP}^2 -surface is *properly convex* if it is a quotient of a properly convex domain in \mathbb{RP}^2 by a properly discontinuous and free action of a subgroup of $\mathrm{PGL}(3, \mathbb{R})$.

A disjoint collection of simple closed geodesics c_1, \dots, c_m *decomposes* an \mathbb{RP}^2 -surface S into subsurfaces S_1, \dots, S_n if each S_i is the closure of a component of $S - \bigcup_{i=1, \dots, m} c_i$ where we do not allow a curve c_i to have two one-sided neighborhoods in only one S_j for some j .

In [17], we proved:

Theorem 4.1. *Let Σ be a closed orientable \mathbb{RP}^2 -surface with principal geodesic or empty boundary and $\chi(\Sigma) < 0$. Then Σ has a collection of disjoint simple closed principal geodesics decomposing Σ into properly convex \mathbb{RP}^2 -surfaces with principal geodesic boundary and of negative Euler characteristic and/or π -annuli with principal geodesic boundary.*

4.2. Complete hyperbolic surfaces and Fuchsian groups. Let G be a discrete subgroup of projective automorphisms of \mathbb{S}_+ in $\mathrm{Aut}(\mathbb{S})$. \mathbb{S}_+ has a complete Hilbert metric defined using cross-ratios and is isometric with the complete hyperbolic plane, forming the so-called Beltrami-Klein model. (See [11].) We assume that G has no elliptic or parabolic elements. Here complete geodesics of the hyperbolic metrics are maximal straight lines in \mathbb{S}_+ and vice versa.

We suppose that G is nonelementary: that is, it does not act on a pair of points in $\mathrm{Cl}(\mathbb{S}_+)$. Then \mathbb{S}_+/G has a complete hyperbolic metric induced from the Hilbert one. \mathbb{S}_+/G is *geometrically finite* in the sense that it contains a compact surface bounded by closed geodesics that is the deformation retract of it.

Let Λ denote the limit set of an orbit $G(x)$ for a point $x \in \mathbb{S}_+$, which is independent of the choice of x , as is well-known from the classical Fuchsian group theory. We know from the Fuchsian group theory that Λ is a Cantor subset of $\mathrm{bd}_{\mathbb{S}} \mathbb{S}_+$ and of Lebesgue measure zero and the set of fixed points of nonidentity elements of G is dense in Λ . Let us define $\tilde{\Sigma}_+ := \mathbb{S}_+/G$. We define $\tilde{\Sigma}'_+$ as $\mathrm{Cl}(\mathbb{S}_+) - \Lambda$. (See Chapter 8 of Beardon [4] for the classical Fuchsian group theory used here.) Let \mathcal{J} be the set indexing the boundary components of $\tilde{\Sigma}'_+$. We denote the boundary components of $\tilde{\Sigma}'_+$ by $\mathbf{b}_i, i \in \mathcal{J}$.

$$\tilde{\Sigma}'_+ = \mathbb{S}_+ \cup \bigcup_{i \in \mathcal{J}} \mathbf{b}_i = \mathrm{Cl}(\mathbb{S}_+) - \Lambda.$$

Then defining $\Sigma'_+ = \tilde{\Sigma}'_+/G$, we obtain a compact surface containing open surface Σ_+ compactifying it with boundary components diffeomorphic to circles.

These surfaces are both examples of \mathbb{RP}^2 -surfaces as they are quotients of open domains by projective automorphism groups.

Note that for each element $g \in G$, either $g(\mathbf{b}_i) \cap \mathbf{b}_i = \emptyset$ or $g(\mathbf{b}_i) = \mathbf{b}_i$ holds with g in the cyclic group generated by the deck transformation corresponding to \mathbf{b}_i and the boundary component of Σ'_+ corresponding to it.

We obtain a convex domain Ω_+ closed in \mathbb{S}_+ bounded by a union of straight segments \mathbf{l}_i for $i \in \mathcal{J}$ with same endpoints as \mathbf{b}_i . That is, \mathbf{l}_i are geodesics in \mathbb{S}_+ connecting ideal endpoints of \mathbf{b}_i . Here, \mathbf{l}_i is $\langle \mathbf{g}_i \rangle$ -invariant for a unique primitive element of $\mathbf{g}_i \in \Gamma$ where \mathbf{g}_i acts along the orientation of \mathbf{l}_i . Since Γ acts as a geometrically finite Fuchsian group on \mathbb{S}_+ without parabolics, the classical Fuchsian group theory tells us that Ω_+/Γ is a compact hyperbolic surface with geodesic boundary that is the convex hull of \mathbb{S}_+/Γ and homeomorphic to Σ'_+ . Thus,

$$\mathbb{S}_+ - \Omega_+^o = \coprod_{i \in \mathcal{J}} D_i$$

where D_i is a convex open domain $\text{bd}_{\mathbb{S}} D_i = \text{Cl}(\mathbf{b}_i) \cup \mathbf{l}_i$ for $i \in \mathcal{J}$ that covers an open annulus in Σ'_+ .

Since \mathbf{l}_i covers a simple closed geodesic in Σ''_+ , it follows that

$$(2) \quad \begin{aligned} \mathbf{l}_i &= \mathbf{l}_j, i = j \in \mathcal{J} \text{ or } \mathbf{l}_i \cap \mathbf{l}_j = \emptyset, i, j \in \mathcal{J}, \\ g(\mathbf{l}_i) &= \mathbf{l}_i \text{ for } g = \mathbf{g}_i^n, n \in \mathbb{Z} \text{ or } g(\mathbf{l}_i) \cap \mathbf{l}_i = \emptyset, g \in \Gamma. \end{aligned}$$

4.3. The dynamic of geodesic flows on hyperbolic surfaces. We recall that Γ is isomorphic to a free group of finite rank ≥ 2 , and \mathbb{S}_+/Γ is a complete genus \tilde{g} hyperbolic surface with b ideal boundary components. Assume that $\mathcal{L}(\Gamma) \subset \text{SO}(2, \mathbb{R})^o$. For convenience, denote by Σ_+ the interior of the surface Σ'_+ from now on. That is, $\Sigma_+ = \mathbb{S}_+/\Gamma$.

The Fuchsian Γ -action on the boundary $\text{bd}_{\mathbb{S}} \mathbb{S}_+$ of the standard disk \mathbb{S}_+ in \mathbb{S} forms a *discrete convergence group*: For every sequence g_j of mutually distinct elements of Γ , there is a subsequence g_{j_k} and two (not necessarily distinct) points a, b in the circle $\text{bd}_{\mathbb{S}} \mathbb{S}_+$ such that

- the sequences $g_{j_k}(x) \rightarrow a$ locally uniformly in $\text{bd}_{\mathbb{S}} \mathbb{S}_+ - \{b\}$, that is, uniformly in any compact subset of $\text{bd}_{\mathbb{S}} \mathbb{S}_+ - \{b\}$, and
- $g_{j_k}^{-1}(y) \rightarrow b$ locally uniformly on $\text{bd}_{\mathbb{S}} \mathbb{S}_+ - \{a\}$ respectively as $k \rightarrow \infty$.

(See [1] for details.) We remark that

$$(3) \quad a, b \in \Lambda.$$

For later purpose, we say that a is an *attractor point* and b is a *repellor point* of the sequence $\{g_{j_k}\}$.

Lemma 4.2. *Let g_{j_k} be a sequence satisfying the above convergence group properties. Suppose that the attractor point a is distinct from the repellor point b . Let a_i and r_i denote the attracting fixed point and the repelling fixed point of g_i respectively. Then it follows that*

$$(4) \quad \{a_{j_k}\} \rightarrow a \text{ and } \{r_{j_k}\} \rightarrow b.$$

Proof. One can use the fact that if a closed arc I is send to its interior I° by a continuous map f , then the fixed points are in the image $f(l)$. (See also ϵ -hyperbolicity in p.256 of [2]. Since $a \neq b$, these transformations are uniformly ϵ -hyperbolic.) \square

Let $\mathbb{U}\mathbb{S}_+$ be the unit tangent vector bundle over \mathbb{S}_+ , and let $\mathbb{U}(\mathbb{S}_+/\Gamma) = \mathbb{U}\Sigma_+$ be the unit tangent vector bundle over $\mathbb{S}_+/\Gamma = \Sigma_+$. By following the geodesics in Σ_+ , we obtain a geodesic flow

$$\Phi : \mathbb{U}\Sigma_+ \times \mathbb{R} \rightarrow \mathbb{U}\Sigma_+.$$

A *geodesic current* is a Borel probability measure on $\mathbb{U}(\mathbb{S}_+/\Gamma)$ invariant under the geodesic flow which is supported on a union of nonwandering geodesics. The subspace $\mathcal{C}(\mathbb{S}_+/\Gamma)$ composed of geodesic currents supported on unions of closed geodesics is dense in the space of all geodesic currents with the weak-* topology.

Let $\mathbb{U}_{\text{rec}}\Sigma_+$ denote the set of unit tangent vectors of nonwandering geodesics on Σ_+ . We recall

Lemma 4.3. *Let Σ_+ be as above. Then*

- $\mathbb{U}_{\text{rec}}\Sigma_+ \subset \mathbb{U}\Sigma_+$ is a connected compact geodesic flow invariant set and is a subset of the compact set $\mathbb{U}\Sigma_+''$.
- The inverse image $\mathbb{U}_{\text{rec}}\mathbb{S}_+$ of $\mathbb{U}_{\text{rec}}\Sigma_+$ in $\mathbb{U}_{\text{rec}}\mathbb{S}_+$ is precisely the set of geodesics with both endpoints in Λ .
- The set of pairs of fixed points for all $g \in G$ is dense in $\Lambda \times \Lambda$.

Proof. The first item is in Lemma 1.2 in [35] and the second item is in the proof of the same lemma. The connectedness is also proved in Lemma 1.3 in [35]. The third item follows from the fact that the set of closed geodesic orbits are dense in $\mathbb{U}_{\text{rec}}\Sigma_+$ as the flow is Anosov. \square

Let Δ denote the diagonal of $\Lambda \times \Lambda$. Such pairs $(\lambda_u, \lambda_s) \in \Lambda \times \Lambda - \Delta$ correspond to nonwandering geodesics in the quotient hyperbolic surface. Thus, the set of closed geodesics is dense in the set of nonwandering geodesics by Lemma 4.3.

We have a compactification picture of the above phenomena: For a nonelementary Fuchsian group $\Gamma_0 \subset \mathrm{PGL}(2, \mathbb{R})$, the closure in the projective space $\mathbb{P}(\mathrm{Mat}_2(\mathbb{R}))$ equals $\Gamma_0 \cup (\Lambda \times \Lambda)$, where $\Lambda \subset \mathrm{bd}_{\mathbb{S}} \mathbb{S}_+$ is the limit set of Γ_0 . Here we could write

$$\mathbb{P}(\mathrm{Mat}_2(\mathbb{R})) = \mathrm{PGL}(2, \mathbb{R}) \cup (\mathbb{RP}^1 \times \mathbb{RP}^1)$$

where $\mathbb{RP}^1 \times \mathbb{RP}^1$ corresponds to the set of projective equivalence classes of rank-one 2×2 real matrices. (Projective equivalence classes of such matrices are completely determined by their kernels and images, which are *arbitrary* lines through the origin in \mathbb{R}^2 .) The kernel of a rank one 2×2 matrix is the undefined set for the corresponding singular projective transformation, and the singular projective transformation is constant on the complement of this point in the one-dimensional real projective space.

5. \mathbb{RP}^2 -SCHOTTKY UNIFORMIZATIONS

Goldman constructed an \mathbb{RP}^2 -surface Σ with a free holonomy group in [31]. The holonomy group is in the image $\mathrm{PSO}(2, 1)$ of $\mathrm{SO}(2, 1)$ in $\mathrm{PGL}(3, \mathbb{R})$. We will show that a domain in \mathbb{S} regularly covers Σ .

As above, let Γ be a Margulis group without parabolics. (We will use the notation of Section 4.2 letting $G = \Gamma$.) We obtain an exact sequence

$$1 \rightarrow \Gamma' \rightarrow \Gamma \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 1$$

where Γ' is the subgroup of Γ of index ≤ 2 so that $\mathcal{L}(\Gamma') \subset \mathrm{SO}(2, 1)^\circ$.

5.1. The construction of the Γ -invariant domain. To begin, we assume $\mathcal{L}(\Gamma) \subset \mathrm{SO}(2, 1)^\circ$ initially and so Γ acts on \mathbb{S}_+ . Then \mathbb{S}_+/Γ is a complete genus \tilde{g} hyperbolic surface with b ideal boundary components where $b \geq 1$ and \tilde{g} is just some integer ≥ 0 . In other words, \mathbb{S}_+/Γ can be compactified by adding boundary components to a compact surface Σ'_+ with b boundary components. The universal cover $\tilde{\Sigma}'_+$ of Σ'_+ is identified with the union of a domain \mathbb{S}_+ and a collection of open arcs in $\mathrm{bd}_{\mathbb{S}} \mathbb{S}_+$. (At the moment, we do not require that Γ acts properly discontinuously or freely on E .)

Let Λ denote the limit set of an orbit $\Gamma(x)$ for $x \in \mathbb{S}_+$. We can identify $\tilde{\Sigma}'_+$ as $\mathrm{Cl}(\mathbb{S}_+) - \Lambda$. Let \mathcal{J} be the set indexing the boundary components of $\tilde{\Sigma}'$. We denote the boundary components of $\tilde{\Sigma}'_+$ by $\mathbf{b}_i, i \in \mathcal{J}$.

$$\tilde{\Sigma}'_+ = \mathbb{S}_+ \cup \bigcup_{i \in \mathcal{J}} \mathbf{b}_i = \mathrm{Cl}(\mathbb{S}_+) - \Lambda.$$

We will be using the same symbols for Γ its extension to \mathbf{S}^3 and its element g and its extension to \mathbf{S}^3 . We identify \mathbb{S} with the boundary of \mathcal{H} , called a *boundary sphere*. Γ acts on the standard circles $\text{bd}_{\mathbb{S}}\mathbb{S}_{\pm}$ and the interiors \mathbb{S}_{\pm} in \mathbb{S} . Then each nonidentity element g of Γ has an attracting fixed point a in $\text{bd}_{\mathbb{S}}\mathbb{S}_+$, a repelling fixed point r in $\text{bd}_{\mathbb{S}}\mathbb{S}_+$, and a saddle-type fixed point s in \mathbb{S}_0 . Of course, their antipodes in \mathbb{S} are also an attracting fixed point a_- in $\text{bd}_{\mathbb{S}}\mathbb{S}_-$ and a repelling fixed point r_- in $\text{bd}_{\mathbb{S}}\mathbb{S}_-$ and a saddle-type fixed point s_- in \mathbb{S}_0 respectively.

Since g acts on $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ and fixes a and r , we obtain that g acts on the two 1-dimensional subspaces tangent to $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ at a and r . Hence, $\{s, s_-\}$ is their intersection. Therefore, we deduce that

$$(5) \quad \{s, s_-\} \subset \varepsilon(a) \cup \varepsilon(r)$$

as $\varepsilon(a)$ and $\varepsilon(r)$ are disjoint halves of these 1-dimensional subspaces.

We now denote by $\tilde{\Sigma}'_-$ the domain of antipodal points of $\tilde{\Sigma}'_+$ in \mathbb{S}_- ; that is, $\tilde{\Sigma}'_- = \mathcal{A}(\tilde{\Sigma}')$. We note that Γ acts properly discontinuously and freely on $\tilde{\Sigma}'_-$ by antipodality, and $\tilde{\Sigma}'_-/\Gamma$ is a compact \mathbb{RP}^2 -surface with geodesic boundary diffeomorphic to Σ'_+ . Denote it by Σ'_- . Also, let Ω_- be the set of antipodal points to Ω_+ . Then Γ acts properly discontinuously and freely on Ω_+ and Ω_- . Denote the respective compact quotient surfaces with geodesic boundary by Σ''_+ and Σ''_- .

For each point x of $\text{bd}_{\mathbb{S}}\mathbb{S}_+$, we recall that $\varepsilon(x)$ is an open segment of length π with end points x and $-x$ tangent to $\text{bd}_{\mathbb{S}}\mathbb{S}_+$. The open arcs $\varepsilon(x)$ for $x \in \text{bd}_{\mathbb{S}}\mathbb{S}_+$ give leaves of the foliation \mathcal{F} in \mathbb{S}_0 . Let us give an induced orientation on Σ'_+ and hence an induced orientation on each \mathbf{l}_i for $i \in \mathcal{J}$. Let p_i and q_i denote the forward and backward end points of \mathbf{l}_i . We draw a segment $s_i = \varepsilon(p_i)$. We also draw a segment $t_i = \varepsilon(q_i)$. Then \mathbf{l}_i, s_i, t_i , and $\mathbf{l}_{i,-}$ bound an open disk D_i invariant under $\langle \mathbf{g}_i \rangle$, which we call a *strip*. We denote by \mathcal{R}_i the open strip union with \mathbf{l}_i and $\mathbf{l}_{i,-}$; that is,

$$\mathcal{R}_i = D_i \cup \mathbf{l}_i \cup \mathbf{l}_{i,-}, \text{bd}_{\mathbb{S}}D_i = \mathbf{l}_i \cup s_i \cup t_i \cup \mathbf{l}_{i,-}.$$

(See Figures 1, 3, and 2 for the pictures of these arcs tangent to \mathbb{S}_+ and \mathbb{S}_- .)

Using \mathcal{F} , we obtain:

Proposition 5.1. *The strips \mathcal{R}_i and \mathcal{R}_j are disjoint for $i \neq j$ where $i, j \in \mathcal{J}$.*

Proof. Since s_i, t_i, s_j, t_j are all leaves of \mathcal{F} corresponding to distinct points of $\text{bd}_{\mathbb{S}}\mathbb{S}_+$, the mutual disjointness of these arcs follows. (See Figure 3.) s_i does not intersect t_i, s_j, t_j for $i \neq j, i, j \in \mathcal{J}$, and t_i does not intersect s_i, s_j, t_j for $i \neq j$. The elements \mathbf{l}_i and $\mathbf{l}_{i,-}$ for $i \in \mathcal{J}$

are mutually disjoint by (2). Thus the boundary $\text{bd}_{\mathbb{S}} \mathcal{R}_i$ and $\text{bd}_{\mathbb{S}} \mathcal{R}_j$ for $i \neq j$ are disjoint. Hence, the conclusion follows. \square




FIGURE 3. The quotient spaces of the two “lune” regions are π -annuli stereographically projected.

Proposition 5.2. *Given \mathcal{R}_i for $i \in \mathcal{J}$ and $\gamma \in \Gamma$, we either have*

$$\gamma(\mathcal{R}_i) = \mathcal{R}_i \text{ or } \gamma(\mathcal{R}_i) \cap \mathcal{R}_i = \emptyset.$$

In the former case, $\gamma = \mathbf{g}_i^n$, $n \in \mathbb{Z}$ for the deck transformation \mathbf{g}_i corresponding to \mathbf{l}_i . Furthermore, $\mathcal{R}_i/\langle \mathbf{g}_i \rangle$ is a compact annulus with geodesic boundary that decomposes into two π -annuli or four elementary annuli along some collection of simple closed geodesics.

Proof. Suppose that $\mathcal{R}_i \cap \gamma(\mathcal{R}_i) \neq \emptyset$ holds. Then $\mathbf{l}_i \cap \gamma(\mathbf{l}_i) \neq \emptyset$ holds since the arcs t_i are mutually disjoint. Hence, (2) implies the result.

There is a subspace l of dimension 1, a geodesic circle, containing \mathbf{l}_i and $\mathbf{l}_{i,-}$ in \mathbb{S} . l and t_i bound a closed disk D_1 and if we remove t_i and the fixed points at the ends of \mathbf{l}_i , this region covers a π -annulus with principal boundary components. The similar statement is true for the disk bounded by l and s_i . Therefore, $A_i = \mathcal{R}_i/\langle \mathbf{g}_i \rangle$ is a union of two compact π -annuli. In other words, $l - \text{Cl}(\mathbf{l}_i) - \text{Cl}(\mathbf{l}_{i,-})$ maps to a simple closed principal geodesic decomposing A_i into two π -annuli with principal boundary. (See Figure 3.) Also, each π -annulus decomposes into two elementary annuli. (See Section 4.1.) \square

We say that for $i, j \in \mathcal{J}$, the annulus $\mathcal{R}_i/\langle \mathbf{g}_i \rangle$ is *equivalent* to $\mathcal{R}_j/\langle \mathbf{g}_j \rangle$ if $\mathcal{R}_j = g(\mathcal{R}_i)$ and $g\mathbf{g}_i g^{-1} = \mathbf{g}_j^\pm$ for $g \in \Gamma$. Thus, in fact, there are only \tilde{b} equivalence classes of annuli of above form.

We define

$$\mathcal{A}_i := \mathcal{R}_i \cap \mathbb{S}_0 = \bigcup_{x \in \mathbf{b}_i} \varepsilon(x) \text{ for } i \in \mathcal{J}.$$

We note that $\mathcal{A}_i \subset \mathcal{R}_i$ for each $i \in \mathcal{J}$. We finally define

$$\begin{aligned} \tilde{\Sigma} &= \tilde{\Sigma}'_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{R}_i \cup \tilde{\Sigma}'_- \\ &= \tilde{\Sigma}'_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{A}_i \cup \tilde{\Sigma}'_- \\ (6) \quad &= \Omega_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{R}_i \cup \Omega_- \\ (7) \quad &= \mathbb{S} - \bigcup_{x \in \Lambda} \text{Cl}(\varepsilon(x)). \end{aligned}$$

an open domain in \mathbb{S} . Since the collection whose elements are of form \mathcal{R}_i mapped to itself by Γ , we showed that Γ acts on this open domain.

5.2. The \mathbb{RP}^2 -surface. First, $\tilde{\Sigma}$ does not contain any fixed point: Suppose that $g \in \Gamma$ acts on \mathbf{l}_i for some $i \in \mathcal{J}$. Then g acts on \mathcal{R}_i and the four attracting and repelling fixed points are the vertices of \mathcal{R}_i and two saddle-type fixed points are in s_i or t_i . Hence, they are outside $\tilde{\Sigma}$.

Suppose that $g \in \Gamma - \{\text{Id}\}$ does not act on any of the boundary components. Then it has four attracting and repelling fixed points $a, r \in \text{bd}_{\mathbb{S}} \mathbb{S}_+$ and $a_-, r_- \in \text{bd}_{\mathbb{S}} \mathbb{S}_-$. By (5), saddle-type fixed points s and s_- are either on a great segment $\varepsilon(a)$ or the other one $\varepsilon(r)$ tangent to $\text{bd}_{\mathbb{S}} \mathbb{S}_+$ at a and r respectively. Since $a, r \in \Lambda$ holds, the fixed points are outside $\tilde{\Sigma}$ by (7).

Let $G \subset \text{SO}(2, 1)^\circ$ be a subgroup whose elements preserve orientation on \mathbb{S}_+ , and any open domain \mathcal{D} an open domain on \mathbb{S} upon which where G acts freely and properly-discontinuously. Then \mathcal{D}/G has an induced orientation from \mathbb{S}_+ .

Theorem 5.3. *Let Γ be an orientation-preserving finitely generated group in $\text{SO}(2, 1)^\circ$ without parabolics acting freely and properly discontinuously on \mathbb{S}_+ (or equivalent on \mathbb{S}_-) isomorphic to a free group of finite rank ≥ 2 . Let \mathbf{b} be the number of ideal boundary components of \mathbb{S}_+/Γ . Then*

- *there exists an open domain \mathcal{D} in \mathbb{S} where Γ acts properly discontinuously and freely, and \mathcal{D}/Γ is homeomorphic to a closed surface of genus g , $g \geq 2$.*
- *As an \mathbb{RP}^2 -surface \mathcal{D}/Γ decomposes along simple closed principal geodesics into a union of b annuli A_1, \dots, A_b and two convex*




FIGURE 4. A figure representing an actual free group action with some of the arcs s_i and t_i drawn in point-plots on stereographically projected sphere as in Figure 2.

\mathbb{RP}^2 -surfaces Σ''_+ and Σ''_- . Each annulus A_i , $i = 1, \dots, b$ decomposes into two π -annuli.

- Γ also acts properly discontinuously and freely on $\mathcal{D}_- := \mathcal{A}(\mathcal{D})$, and \mathcal{D}_-/Γ is a closed \mathbb{RP}^2 -surface diffeomorphic to \mathcal{D}/Γ .
- \mathcal{D}_- also decomposes into Σ''_+ and Σ''_- and b annuli A'_i .
- The antipodal map $\mathcal{A} : \mathbb{S} \rightarrow \mathbb{S}$ induces a projective diffeomorphism between \mathcal{D}/Γ and \mathcal{D}_-/Γ .

Proof. We begin by letting $\mathcal{D} = \tilde{\Sigma}$ obtained as above. We showed that Γ has no fixed point on \mathcal{D} . Let $\text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ denote the closure of \mathcal{A}_i in \mathcal{D} for each $i \in \mathcal{J}$.

Now Γ acts properly discontinuously on $\tilde{\Sigma}'_+$ and $\tilde{\Sigma}'_-$ and each $g \in \Gamma$ sends $\text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ to $\text{Cl}_{\mathcal{D}}(\mathcal{A}_j)$ for $j \neq i$ or g is in the infinite cyclic subgroup of Γ acting properly discontinuously on $\text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ if $g(\text{Cl}_{\mathcal{D}}(\mathcal{A}_i)) = \text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$. Hence, Γ acts properly discontinuously on $\bigcup_{i \in \mathcal{J}} \text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$.

For any compact set K in $\tilde{\Sigma}$, we consider the compact sets

$$K_1 := K \cap \tilde{\Sigma}'_+, K_{-1} := K \cap \tilde{\Sigma}'_-, \text{ and } K_0 := K \cap \bigcup_{i \in \mathcal{J}} \text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$$

since these are closed. We have $K = K_+ \cup K_- \cup K_0$.

Since each of $\text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ has an open neighborhood O_i forming a mutually disjoint collection of open sets covering K_0 , it follows that K_0 meets only finitely many $\text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$.

If $g(K) \cap K \neq \emptyset$, we have $g(K_i) \cap K_i \neq \emptyset$ for some $i = -1, 0, 1$. Since Γ acts properly discontinuously on $\tilde{\Sigma}'_+$ and $\tilde{\Sigma}'_-$ and $\bigcup_{i \in \mathcal{J}} \text{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ respectively, there exist only finitely many elements g of Γ so that $g(K_i) \cap K_i \neq \emptyset$ for each i , $i = -1, 0, 1$. We showed that Γ acts properly discontinuously and freely on \mathcal{D} .

Since $\tilde{\Sigma}'_+/\Gamma$ and $\tilde{\Sigma}'_-/\Gamma$ are compact surfaces, and $\bigcup_{i \in \mathcal{J}} \text{Cl}_{\mathcal{D}}(\mathcal{A}_i)/\Gamma$ is a union of finitely many compact annuli, it follows that \mathcal{D}/Γ is a closed surface.

For the second item, \mathcal{D}/Γ decomposes along the images of the union of $\mathbf{l}_i, \mathbf{l}_{i,-}, i \in \mathcal{J}$ to Σ''_+ and Σ''_- and annuli with boundary components that are images of the union of \mathbf{l}_i and $\mathbf{l}_{i,-}$, $i \in \mathcal{J}$. Proposition 5.2 implies that these annuli are obtained from two π -annuli.

The third item follows since the antipodal map \mathcal{A} conjugates Γ to itself and $\mathcal{A}(\mathcal{D}) = \mathcal{D}_-$. Let κ denote the induced diffeomorphism $\Sigma \rightarrow \mathcal{D}_-/\Gamma$ where κ sends Σ''_- to Σ''_+ and vice versa. κ sends annuli in Σ to ones in \mathcal{D}_-/Γ but they do not share subdomains in the universal cover. (Actually \mathcal{R}_i goes to its antipodal image which is distinct from \mathcal{R}_i for each $i \in \mathcal{J}$.)

The rest follows from this observation. \square

The proof of Theorem 1.1. Now, we consider the case when Γ is a subgroup of $\text{SO}(2, 1)$ in general. Recall that Γ' is the subgroup of Γ of index ≤ 2 so that $\mathcal{L}(\Gamma') \subset \text{SO}(2, 1)^\circ$. We construct \mathcal{D}' for Γ' by Theorem 5.3. Let ψ be the element of $\Gamma - \Gamma'$. Then ψ is orientation-preserving but $\psi(\mathbb{S}_+) = \mathbb{S}_-$. Since ψ preserves orientation, ψ acts on $\coprod_{i \in \mathcal{J}} \mathcal{A}_i$ as constructed above for Γ' as well. Thus, ψ acts on \mathcal{D}' and hence Γ acts on \mathcal{D}' .

Recall from the introduction that $\mathcal{L}'(\Gamma)$ is a free group of rank ≥ 2 acting freely on \mathbb{S}_+ as a subgroup of $\text{PGL}(3, \mathbb{R})$ and $\mathbb{S}_+/\mathcal{L}'(\Gamma)$ is a nonorientable complete hyperbolic surface. Since $\mathcal{L}'(\Gamma)$ acts freely on

\mathbb{S}_+ , it follows that ψ goes to an element of $\mathrm{PGL}(3, \mathbb{R})$ that acts freely on \mathbb{S}_+ and on $\bigcup_{i \in \mathcal{J}} \mathbf{b}_i$. Hence, ψ as an element of $\mathrm{Aut}(\mathbf{S}^3)$ acts without fixed point on

$$\bigcup_{i \in \mathcal{J}} \bigcup_{x \in \mathbf{b}_i} \varepsilon(x) = \coprod_{i \in \mathcal{J}} \mathcal{A}_i$$

since ψ does not act on any fiber of the fibration. Hence ψ has no fixed point on

$$\tilde{\Sigma}'_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{A}_i \cup \tilde{\Sigma}'_-.$$

\mathcal{D}'/Γ' double-covers a closed surface \mathcal{D}/Γ .

Finally, it is straightforward to verify the same statements for $\mathcal{A}(\mathcal{D})$. Replacing with this set is equivalent to replacing $\mathrm{Cl}(\varepsilon(x))$ with $\mathrm{Cl}(\varepsilon(x)_-)$ for every appropriate point x in (7).

The uniqueness follows from Theorem R of [31] since for a given Fuchsian holonomy, a closed \mathbb{RP}^2 -surface is obtained by construction called grafting. Grafting is classified by a word of free semigroup of rank two for each boundary component of Σ'_+ . We can easily show that only two cases for each boundary component result in imbedded domains for the universal cover and developing maps in all other cases are immersions that are not imbeddings. (See also [18].) \square

6. THE DYNAMIC CORE OF A MARGULIS SPACETIME

This section briefly summarizes the relevant results of Goldman, Labourie, and Margulis in [35] and [34].

6.1. Margulis invariants. Let Γ be a Margulis group without any parabolic elements and with the property $\mathcal{L}(\Gamma) \subset \mathrm{SO}(2, 1)^\circ$. Γ acts on \mathbb{S}_+ as a geometrically finite Fuchsian group.

An element $g \in \Gamma - \{\mathrm{I}\}$ is hyperbolic as it is not parabolic or elliptic. We choose so that $\{\mathbf{v}_+(g), \mathbf{v}_0(g), \mathbf{v}_-(g)\}$ are in the standard orientation of \mathbb{R}^3 ; or $\mathbf{v}_-(g) \times \mathbf{v}_+(g) = \mathbf{v}_0(g)$. (See Definition 3.2.) We recall the Margulis invariant $\mu : \Gamma - \{\mathrm{I}\} \rightarrow \mathbb{R}$

$$\mu(g) := \mathbf{B}(gx - x, \mathbf{v}_0(g)) \text{ for } g \in \Gamma - \{\mathrm{I}\}, x \in \mathsf{E},$$

which is independent of the choice of x in E .

If Γ acts freely and properly on E , then Margulis invariants of non-identity elements are all positive or all negative by the opposite sign lemma. (See [35].) By choosing the opposite orientation of E , we can change the all signs of the invariants. We henceforth assume that the Margulis invariants of nonidentity elements are all positive.

In the second main theorem of [35], they extend the Margulis invariants to the geodesic current space on the hyperbolic space \mathbb{S}_+/Γ , that is, diffused Margulis invariants for currents.

The Margulis invariants of elements of $\mathcal{C}(\mathbb{S}_+/\Gamma)$ are positive by the second main theorem in [35], and the subspace of currents supported closed geodesics with total measure 1 is precompact in $\mathcal{C}(\mathbb{S}_+/\Gamma)$. It follows that there exists a global $C > 0$ independent of $g \in \Gamma$ so that

$$(8) \quad 1/C \leq \frac{\mu(g)}{l_{\mathbb{S}_+}(g)} \leq C \text{ for } g \in \Gamma - \{\text{I}\}$$

where $l_{\mathbb{S}_+}(g)$ denotes the minimum of the hyperbolic distances between x and $g(x)$ for $x \in \mathbb{S}_+$ where g is considered as an element of $\text{PGL}(3, \mathbb{R})$ acting on $\mathbb{S}_+ \subset \mathbb{RP}^2$. (This equals the length of the closed geodesic represented by g . See also Corollary of [35].)

We remark that the positivity of the diffused Margulis-invariants is equivalent to the proper discontinuity of the action of Γ on E which is the main result of Goldman, Labourie, and Margulis [35].

Goldman, Minsky, and Margulis [36] show that the positivity of the Margulus-invariants for closed curves is not sufficient to obtain the proper discontinuity of Γ on E by finding counter-examples when Σ_+ is a one-holed torus but shows that the positivity of diffused Margulis invariants laminations on surfaces is sufficient.

6.2. Neutralized sections. We note that Γ acts on $\mathbb{U}\mathbb{S}_+$ as a deck transformation group over $\mathbb{U}\Sigma_+$. An element $\gamma \in \Gamma$ goes to the differential map $D\gamma : \mathbb{U}\mathbb{S}_+ \rightarrow \mathbb{U}\mathbb{S}_+$. They in [34] constructed a flat affine bundle \mathbf{E} over the unit tangent bundle $\mathbb{U}\Sigma_+$ of Σ_+ . They took the quotient $\mathsf{E} \times \mathbb{U}\mathbb{S}_+$ by the diagonal action given by $\gamma(v, x) = (\gamma(v), D\gamma(x))$ for a deck transformation $\gamma \in \Gamma$ of the cover $\mathbb{U}\mathbb{S}_+$ of $\mathbb{U}\Sigma_+$.

The cover of \mathbf{E} is denoted by $\hat{\mathbf{E}}$ and is identical with $\mathsf{E} \times \mathbb{U}\mathbb{S}_+$. We denote by

$$\pi_{\mathsf{E}} : \hat{\mathbf{E}} = \mathsf{E} \times \mathbb{U}\mathbb{S}_+ \rightarrow \mathsf{E}$$

the projection.

We define \mathbf{V} as the quotient space of $\mathsf{V} \times \mathbb{U}\mathbb{S}_+$ under the diagonal action defined by $\gamma(v, x) = (\mathcal{L}(\gamma)(v), D\gamma(x))$ for a deck transformation $D\gamma$ of the cover $\mathbb{U}\mathbb{S}_+$ of $\mathbb{U}\Sigma_+$.

A *neutral section* of \mathbf{V} is a $\text{SO}(2, 1)$ -invariant section which is parallel along geodesic flow of $\mathbb{U}\Sigma_+$. A neutral section $\nu : \mathbb{U}\Sigma_+ \rightarrow \mathbf{V}$ arises from a graph of the $\text{SO}(2, 1)$ -invariant map

$$\tilde{\nu} : \mathbb{U}\mathbb{S}_+ \rightarrow \mathsf{V}$$

with the image in the space of unit spacelike vectors in V : $\tilde{\nu}$ is defined by sending a unit vector u in $\mathbb{U}\mathbb{S}_+$ to the normalization of the Lorentzian cross-product $\rho(u) \times \alpha(u)$ of the null vectors $\rho(u)$ and $\alpha(u)$ respectively representing the starting point and the ending point in $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ of the hyperbolic geodesic tangent to u in \mathbb{S}_+ . (See Section 4.2 of [35] for details.)

Lemma 6.1. *Suppose that u is a unit tangent vector to a geodesic where a deck transformation γ acts on along the orientation given by u . Then*

- $\rho(u)$ and $\alpha(u)$ respectively correspond in \mathbb{S}_+ a repelling fixed point and an attracting fixed point on $\text{bd}_{\mathbb{S}}\mathbb{S}_+$,
- $\tilde{\nu}(u)$ is an eigenvector of $\mathcal{L}(\gamma)$ and corresponds to the saddle-type fixed point of γ in \mathbb{S}_0 .
- γ acts on a unique spacelike line in E in the direction of $\tilde{\nu}(u)$ as a translation by the vector $\mu(\gamma)\tilde{\nu}(u)$ for the Margulis invariant μ .

Proof. Since the geodesic is invariant, the endpoints are attracting or repelling fixed points. The Lorentzian cross-product of the attracting eigenvector and the repelling eigenvector is clearly another eigenvector. The final item is from an interpretation of the Margulis invariants. \square

Let ∇ be the flat connection on E as a bundle over $\mathbb{U}\mathbb{S}_+$ with fiber isomorphic to E induced from the product structure of $\tilde{E} = E \times \mathbb{U}\mathbb{S}_+$. (See Sections 3.2 and 3.3 of [35].) We induce a flow

$$\tilde{\Phi} : E \times \mathbb{U}\mathbb{S}_+ \times \mathbb{R} \rightarrow E \times \mathbb{U}\mathbb{S}_+$$

by letting the flow be constant in the second factor and be the geodesic flow on $\mathbb{U}\mathbb{S}_+$. This induces an Anosov flow

$$E \times \mathbb{R} \rightarrow E.$$

Using this Anosov geodesic flow of E in Lemma 8.4 of [35], Goldman, Labourie, and Margulis show that there exists a *neutralized section*, that is, a section $\mathcal{N} : \mathbb{U}_{\text{rec}}\Sigma_+ \rightarrow E$ satisfying

$$(9) \quad \nabla_X \mathcal{N} = f\nu$$

where X is the vector field of geodesic flow on $\mathbb{U}_{\text{rec}}\Sigma_+$ and f is a real valued function defined on $\mathbb{U}_{\text{rec}}\Sigma_+$. Lemma 3 of [34] shows that we can choose \mathcal{N} so that $f > 0$ by changing \mathcal{N} to $\mathcal{N} + g\nu$ for a suitable real valued $g : \mathbb{U}_{\text{rec}}\Sigma_+ \rightarrow \mathbb{R}$.

6.3. Lifting the neutralized section to the coverings. Let $\mathbb{U}_{\text{rec}}\mathbb{S}_+$ denote the inverse image of $\mathbb{U}_{\text{rec}}\Sigma_+$ in $\mathbb{U}\mathbb{S}_+$. We take a compact fundamental domain F in $\mathbb{U}_{\text{rec}}\mathbb{S}_+$ of $\mathbb{U}_{\text{rec}}\Sigma_+$, and then \mathcal{N} lifts on F to a map $\tilde{\mathcal{N}}|F$ to $\tilde{\mathbf{E}}$ so that for each side pairing $g_S \in \Gamma$ of F for a side S of F ,

$$(10) \quad \begin{aligned} \tilde{\mathcal{N}} \circ g_S(\vec{u}) &= g_S \circ \tilde{\mathcal{N}}(\vec{u}) \\ &= (g_S(\pi_{\mathbf{E}} \circ \tilde{\mathcal{N}}(\vec{u})), g_S(\vec{u})) \text{ for every } \vec{u} \in F \end{aligned}$$

holds since \mathcal{N} is obtained from an arbitrary lift satisfying this equation and deforming it (see the proof of Lemma 8.3 of [35]). Thus, we find the section $\tilde{\mathcal{N}} : \mathbb{U}_{\text{rec}}\mathbb{S}_+ \rightarrow \tilde{\mathbf{E}}$ lifting \mathcal{N} satisfying

$$(11) \quad \tilde{\mathcal{N}} \circ \gamma = \gamma \circ \tilde{\mathcal{N}}$$

for each deck transformation γ of $\mathbb{U}\mathbb{S}_+ \rightarrow \mathbb{U}\mathbb{S}_+$.

Let \mathbf{u}' be a unit tangent vector to a closed geodesic in Σ_+ . The geodesic flow line α on $\mathbb{U}_{\text{rec}}\Sigma_+$ tangent to \mathbf{u}' will return to \mathbf{u}' after a period. Suppose that a deck transformation γ in $\mathbb{U}_{\text{rec}}\mathbb{S}_+$ corresponds to a closed geodesic in Σ_+ , and acts on a geodesic l_γ in \mathbb{S}_+ containing a unit vector \mathbf{u} . Let $\overrightarrow{l}_\gamma$ denote the subspace of unit tangent vectors to l_γ in the γ -direction. Since $\tilde{\mathcal{N}}$ is a lift of \mathcal{N} ,

$$\pi_{\mathbf{E}} \circ \tilde{\mathcal{N}}| \overrightarrow{l}_\gamma$$

maps to a line parallel to $\tilde{\nu}(\mathbf{u})$ up to an action of $\mathcal{L}(\Gamma)$ by (9). Since γ acts on the image of this line in \mathbf{E} by (11), and $\mathcal{L}(\gamma)$ has $\tilde{\nu}(\mathbf{u})$ as an eigenvector by Lemma 6.1,

$$\pi_{\mathbf{E}} \circ \tilde{\mathcal{N}}| \overrightarrow{l}_\gamma$$

is a line parallel to $\tilde{\nu}(\mathbf{u})$.

Equation (9) also lifts to

$$(12) \quad \nabla_X \tilde{\mathcal{N}} = f \tilde{\nu}$$

where X is the unit-vector field of geodesic flow on $\mathbb{U}\mathbb{S}_+$, ∇ is the flat connection on $\mathbf{E} \times \mathbb{U}\mathbb{S}_+$, and f is a positive-valued function. Thus, we conclude that the geodesic flow on $\mathbb{U}_{\text{rec}}\mathbb{S}_+$ is conjugated to one on the image subset of $\mathbb{U}\mathbf{E}$ by a map associated with $\pi_{\mathbf{E}} \circ \tilde{\mathcal{N}}$. (We note here that the flow speed changes by the diffused Margulis invariants.)

6.4. Mapping the convex core. Theorem 1 of [34] shows that the image of a geodesic γ of \mathbb{S}_+ mapping to a nonwandering geodesic in Σ_+ under $\pi_{\mathbf{E}} \circ \tilde{\mathcal{N}}$ is a spacelike geodesic in \mathbf{E} mapping to a nonwandering geodesic in \mathbf{E}/Γ .

Let $\mathbb{U}\mathbf{E}$ denote the space of unit spacelike vectors at each point of \mathbf{E} and $\mathbb{U}(\mathbf{E}/\Gamma)$ the space of unit spacelike vectors at each point of \mathbf{E}/Γ .

Denote by $\mathbb{U}_{\text{rec}}\mathsf{E}$ in $\mathbb{U}\mathsf{E}$ the inverse image of the subset of $\mathbb{U}_{\text{rec}}(\mathsf{E}/\Gamma)$ composed of unit tangent spacelike vectors tangent to nonwandering spacelike geodesics.

Denote by \overrightarrow{yx}^o the open line segment connecting two points $x, y \in \text{bd}_{\mathbb{S}}\mathbb{S}_+$ with the orientation towards x .

We give the topology of the space $\mathcal{G}_{\text{rec}}\mathbb{S}_+$ of oriented geodesics in \mathbb{S}_+ mapping to nonwandering oriented geodesics in Σ_+ by the quotient topology from the unit tangent bundle $\mathbb{U}_{\text{rec}}\mathbb{S}_+$. Similarly, we give the topology of the space $\mathcal{G}_{\text{rec}}\mathsf{E}$ of spacelike oriented geodesics in E mapping to nonwandering geodesics in E/Γ by the quotient topology from the space $\mathbb{U}_{\text{rec}}\mathsf{E}$. (As usual, these are standard topological structures on subspaces of complete geodesics in \mathbb{S}_+ and E .)

A *bounded subset* of $\mathcal{G}_{\text{rec}}\mathbb{S}_+$ is a set of geodesics passing a bounded set in $\mathbb{U}\mathbb{S}_+$, and a *bounded subset* of $\mathcal{G}_{\text{rec}}\mathsf{E}$ is a set of spacelike geodesics passing a bounded set in $\mathbb{U}\mathsf{E}$ under the Euclidean metric d_E .

Proposition 6.2. *The map $\tilde{\mathcal{N}}$ induces a continuous function $\mathcal{N} : \mathcal{G}_{\text{rec}}\mathbb{S}_+ \rightarrow \mathcal{G}_{\text{rec}}\mathsf{E}$ where*

- For the attracting fixed point a and the repelling one r of g ,

$$\widehat{\text{Axis}}(g) = \mathcal{N}(\overrightarrow{ra}^o).$$

- the bounded set of elements of $\mathcal{G}_{\text{rec}}\mathbb{S}_+$ maps to a bounded set in $\mathcal{G}_{\text{rec}}\mathsf{E}$.
- Finally, the map is surjective.

Proof. For a flow segment in $\mathbb{U}_{\text{rec}}\mathbb{S}_+$, $\pi_{\mathsf{E}} \circ \tilde{\mathcal{N}}$ sends it to a spacelike line segment in E . This induces \mathcal{N} . The first item is discussed in Section 6.3.

The next statement follows by the fact that \mathcal{N} is induced by the continuous map $\pi_{\mathsf{E}} \circ \tilde{\mathcal{N}}$ which sends a compact set to a compact set.

Clearly our map is surjective from the set of geodesics in \mathbb{S}_+ mapping to closed ones in \mathbb{S}_+/Γ to the set of geodesics in E mapping to closed ones in E/Γ since any a closed geodesic correspond to a unique nontrivial element of Γ in the one-to-one manner. The surjectivity of \mathcal{N} now follows by Theorem 1 of [34] by the density of closed geodesics and the continuity of \mathcal{N} . \square

7. Γ ACTS PROPERLY ON $\mathsf{E} \cup \tilde{\Sigma}$.

Let Γ be a proper affine deformation of a convex cocompact Fuchsian group of rank ≥ 2 . We assume initially that $\mathcal{L}(\Gamma) \subset \text{SO}(2, 1)^o$ and \mathbb{S}_+/Γ is a complete genus $\tilde{\mathbf{g}}$ hyperbolic surface with \mathbf{b} ideal boundary components. As above, we assume that the Margulis invariants of nonidentity elements are all positive.




FIGURE 5. The action of a Lorentzian isometry \hat{g} on the 3-hemisphere \mathcal{H} where the boundary sphere \mathbb{S} is the unit sphere with center $(0, 0, 0)$ here.

7.1. The action of projective-boost automorphisms. Recall that \mathbf{S}^3 is the quotient $\mathbb{R}^4 - \{O\} / \sim$ where \sim is given by $v \sim w$ iff $v = sw$ for $s > 0$. Recall that homogeneous coordinate system of \mathbf{S}^3 is given by setting a point p of \mathbf{S}^3 to have coordinates $[x : y : z : t]$ where (x, y, z, t) divided by its norm represent p as a unit vector in \mathbf{S}^3 . We identified E with the open affine space given by $t > 0$. We use the coordinates so that \mathbb{S} corresponds to the hyperplane given by $t = 0$ and the origin O of E is given by e_0 . The closure of E is the 3-dimensional hemisphere \mathcal{H} with $\partial\mathcal{H} = \mathbb{S}$. For this system of coordinates, $(x, y, z) \in E$ is given homogeneous coordinates $[x : y : z : 1]$.

A projective automorphism g that is of form

$$(13) \quad \begin{bmatrix} \lambda & 0 & 0 & 0 \\ 0 & 1 & 0 & k \\ 0 & 0 & \frac{1}{\lambda} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \lambda > 1, k \neq 0$$

under a homogeneous coordinate system of \mathbf{S}^3 is a projective-boost. In this coordinate system, g acts on an open 3-dimensional hemisphere to be identified with E given by $t > 0$, has an invariant axis $\text{Axis}(g)$ given by $x = 0, z = 0$, given by $x = 0$ and one U given by $z = 0$ containing it. On S , g contracts vectors along a direction by a constant $1/\lambda$ and g expands the vectors along a direction by a constant λ on U assuming $\lambda > 1$. The subspace S given by $x = 0$ is the stable subspace, and the subspace U given by $z = 0$ is the unstable subspace. (See Section 3.4.)

Let η_+ be the segment $\overline{\mathbf{e}_1 \mathbf{e}_2 \mathbf{e}_{1-}}$, “the attracting arc”. Let η_- denote the segment $\overline{\mathbf{e}_3 \mathbf{e}_{2-} \mathbf{e}_{3-}}$, “the repelling arc”. These have $\mathbf{d}_{\mathbf{S}^3}$ -lengths equal to π . The sphere \mathbf{S}_0^2 is called an *unstable sphere*.

We will prove the following lemma for a properly convex K because the proof is much simpler. But this is true for general compact subset K .

Lemma 7.1. *Let $g_{\lambda,k}$ denote the projective boost on \mathcal{H} defined by the above matrix in (13) for a homogeneous coordinate system with coordinate functions t, x, y, z in the given order and let \mathbf{S}_0^2 denote the subspace given by $x = 0$. We assume that $k \geq 0, \lambda > 0$. Then as $\lambda, k \rightarrow +\infty$ where $k/\lambda \rightarrow 0$, the following statements hold:*

- (a) $g_{\lambda,k}|_{\mathcal{H} - \mathbf{S}_0^2}$ converges in the compact open topology to a projective endomorphism Π_0 given by sending $[x : y : z : t]$ to \mathbf{e}_1 or \mathbf{e}_{1-} where the sign depends on the sign of x . ($t \geq 0$ always.)
- (b) $g_{\lambda,k}|(\mathbf{S}_0^2 \cap \mathcal{H}) - \eta_-$ converges in the compact open topology to a projective endomorphism Π_1 given by sending $[0 : y : z : t]$ to \mathbf{e}_2 .
- (c) For a properly convex compact set K in $\mathcal{H} - \eta_-$, we can choose a subsequence of $\{g_{\lambda,k}(K)\}$ that converges to either
 - (i) $\{\mathbf{e}_2\}$ if $K \subset \text{Cl}(U) - \eta_-$,
 - (ii) the point \mathbf{e}_1 or \mathbf{e}_{1-} if $K \cap \text{Cl}(U) = \emptyset$,
 - (iii) one of the segments

$$\overline{\mathbf{e}_1 \mathbf{e}_2}, \overline{\mathbf{e}_{1-} \mathbf{e}_2},$$

when K meets only one component of $\mathcal{H} - \text{Cl}(U)$, or
(iv) $\eta_+ = \overline{\mathbf{e}_1 \mathbf{e}_2 \mathbf{e}_{1-}}$.

(d) Finally, the convergences here are uniform independent of K provided $\mathbf{d}_{\mathbf{S}^3}(K, \eta_-) \geq \delta$ in cases (ii), (iii), and (iv) or

$$\mathbf{d}_{\mathbf{S}^3}(K, \text{Cl}(U)) > \delta$$

in case (i) for each fixed $\delta > 0$.

Proof. The first two items follow by normalizing the above matrix by diving it by the maximal norm of the entries.

For any properly convex compact subset K' in $\mathcal{H} - \mathbf{S}_0^2$, $\{g_{\lambda,k}(K')\}$ converges to one of $\{\mathbf{e}_1\}$ or $\{\mathbf{e}_{1-}\}$ as $\lambda, k \rightarrow \infty$ while $k/\lambda \rightarrow 0$ by the first item. Also, the case for $K' \subset \text{Cl}(U)$, the proof is similar as $k \rightarrow +\infty$.

So, we now suppose that $K' \cap \mathbf{S}_0^2 \neq \emptyset$. Suppose first that K' meets both components of $\mathcal{H} - \mathbf{S}_0^2$. For a properly convex compact subset K' in $\mathcal{H} - \eta_-$, let K'_1 be the nonempty properly convex compact set $K' \cap \mathbf{S}_0^2 - \eta_-$. We can take a closed 2-dimensional hemisphere H with boundary $\partial H \subset \mathbb{S}$ so that H is the topological boundary in \mathcal{H} of a convex open domain $H' \subset \mathcal{H}$ disjoint from η_- and containing K' where we can freely and slightly enlarge subspaces and take joins. (For example, we can choose H to be a plane in E separating K' and η_- containing \mathbf{e}_3 and \mathbf{e}_{3-} in the boundary.) Then the following statements hold:

- there exist a properly convex compact domain D_1 in a component of $H' - \mathbf{S}_0^2$ with $x > 0$ and another one D_2 in a component of $H' - \mathbf{S}_0^2$ with $x < 0$ so that K' is in the join J of D_1 and D_2 that can be chosen disjoint from η_- .
- For $K_1 := J \cap \mathbf{S}_0^2 \supset K'_1$, the join J_1 of K_1 and D_1 and the join J_2 of K_1 and D_2 satisfy $J = J_1 \cup J_2$.

For example, we may obtain D_1 and D_2 by removing from K' an ϵ - $\mathbf{d}_{\mathbf{S}^3}$ -neighborhood of \mathbf{S}_0^2 for small $\epsilon > 0$ and making them slightly larger so that the join of D_1 and D_2 contains K' .

By the second item of Lemma 7.1, $\{g_{\lambda,k}(K_1)\}$ converges to \mathbf{e}_2 as $\lambda, k \rightarrow \infty$. Since

$$\{g_{\lambda,k}(D_1)\} \rightarrow \{\mathbf{e}_1\} \text{ and } \{g_{\lambda,k}(D_2)\} \rightarrow \{\mathbf{e}_{1-}\}$$

as $\lambda, k \rightarrow \infty$ by above,

- $\{g_{\lambda,k}(J_1)\} \rightarrow s_1 = \overline{\mathbf{e}_1 \mathbf{e}_2}$ in \mathbb{S} and
- $\{g_{\lambda,k}(J_2)\} \rightarrow s_2 = \overline{\mathbf{e}_{1-} \mathbf{e}_2}$ in \mathbb{S} hold by Lemma 2.3 respectively.

Hence, we conclude $\{g_{\lambda,k}(J)\} \rightarrow \eta_+$ as $\lambda, k \rightarrow \infty$.

For each $\epsilon > 0$,

$$g_{\lambda,k}(K') \subset N_\epsilon(\eta_+) = N_\epsilon(s_1 \cup s_2)$$

holds whenever λ and k are sufficiently large, and there are points arbitrarily close to both points \mathbf{e}_1 and \mathbf{e}_{1-} . As K' is convex and connected, the geometric limit of $\{g_{\lambda,k}(K')\}$ exists as $\lambda, k \rightarrow \infty$ and equals η_+ . (See Figure 5 also.)

So, we now suppose that $K' \cap \mathbf{S}_0^2 \neq \emptyset$ and that K' meets only one component of $\mathcal{H} - \mathbf{S}_0^2$. Then the arguments are simpler and $\{g_{\lambda,k}(K')\}$ converges to $\overline{\mathbf{e}_1 \mathbf{e}_2}$ or $\overline{\mathbf{e}_{1-} \mathbf{e}_2}$ as $\lambda, k \rightarrow \infty$.

The final item (d) for cases (i) and (ii) can be obtained by easy estimations. For (d)(iii) and (d)(iv), we apply the results for (i) or (ii) to D_1 and D_2 and K_1 . The uniform convergences for the joins J_1 and J_2 follow by estimations also. \square

We remark that our proof of Proposition 7.2 is a generalization of the above one.

7.2. The proper discontinuity of the action of Γ . Recall from Theorem 5.3 that we can construct

$$(14) \quad \tilde{\Sigma} := \mathbb{S}_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{R}_i \cup \mathbb{S}_- \subset \mathbb{S},$$

that is a Γ -invariant open domain with an infinitely generated fundamental group and covers a closed \mathbb{RP}^2 -surface Σ , a union of Σ'_+ and Σ'_- and annuli A_i , $i = 1, \dots, b$.

Proposition 7.2. *Let Γ be a Margulis group without parabolic holonomy. Let $\tilde{\Sigma}$ be as determined above by (14). Assume $\mathcal{L}(\Gamma) \subset \text{SO}(2, 1)^\circ$. Then Γ acts freely and properly discontinuously on $\mathsf{E} \cup \tilde{\Sigma}$ as a group of projective automorphisms of \mathbf{S}^3 .*

Proof. Since Γ acts freely and properly discontinuously on $\mathbb{R}^3 = \mathsf{E}$ and so on the constructed $\tilde{\Sigma}$ separately, Γ acts freely on the union.

Suppose that there exists a sequence $\{g_i\}$ of elements of Γ and a compact subset K of $\mathsf{E} \cup \tilde{\Sigma}$ so that

$$(15) \quad g_i(K) \cap K \neq \emptyset \text{ for all } i.$$

We will show that actually the sequence $\{g_i\}_{i \in \mathbb{Z}^+}$ has to be a finite sequence for the set of positive integers \mathbb{Z}^+ .

We prove by contradiction: Suppose that $\{g_i\}$ is an infinite sequence and $g_i \neq g_j$ whenever $i \neq j$. Then (15) holds universally in this section even if we take any subsequences.

Recall that Γ is a convergence group acting on $\text{bd}\mathbb{S}_+$. Let a_i and r_i denote the attracting fixed point and the repelling fixed point of g_i respectively. We can choose a subsequence g_{j_k} satisfying the convergence group property (see Section 4.3). For the attracting fixed point a_{j_k} and

the repelling fixed point r_{j_k} in $\text{bd}\mathbb{S}_+$ of g_{j_k} , (4) shows $\{a_{j_k}\} \rightarrow a$ and $\{r_{j_k}\} \rightarrow b$ for $a, b \in \text{bd}\mathbb{S}_+$ provided $a \neq b$.

(I) *First, we will consider the case when $a \neq b$:*

(I)(i) *We collect some convergence properties of [35]:* Now letting $\{g_i\}$ denote the subsequence, we note that each g_i has an attracting fixed point a_i and a repelling fixed point r_i in $\text{bd}_{\mathbb{S}}\mathbb{S}_+$. By our conditions, $\{a_i\} \rightarrow a$ and $\{r_i\} \rightarrow b$ hold by Lemma 4.2.

We define the Lorentzian cross-product

$$\nu := \frac{\beta \times \alpha}{|||\beta \times \alpha|||}$$

of nonzero vectors α and β corresponding to a and b respectively. A Lorentzian isometry element g_i acts as a translation on a unique space-like line $\text{Axis}(g_i)$ in the direction ν_i of eigenvalue 1. Let α_i and ρ_i denote the null vectors in the directions of a_i and r_i respectively so that $\alpha_i \rightarrow \alpha$ and $\rho_i \rightarrow \rho$ hold. We compute

$$\nu_i := \frac{\rho_i \times \alpha_i}{|||\rho_i \times \alpha_i|||}; \text{ i.e., } \nu_i = \nu_0(g_i).$$

Thus, $[\nu_i] \rightarrow [\nu] \in \mathbb{S}$ holds. Since $\{a_i\} \rightarrow a$ holds, the sequence $a_i[\nu_i]a_{i,-} = \text{Cl}(\varepsilon(a_i))$ converges to a segment $\overline{a[\nu]a_-} = \text{Cl}(\varepsilon(a))$. (We can assume without loss of generality that

$$|||\rho_i \times \alpha_i||| = 1, |||\beta \times \alpha||| = 1.)$$

Since the geodesics with end points a_i, r_i pass the bounded part K'' of the unit tangent bundle of \mathbb{S}_+ , it follows that

$$\text{Cl}(\text{Axis}(g_i)) = \text{Cl}(\mathcal{N}(\overrightarrow{r_i a_i^o})) \rightarrow \text{Cl}(\mathcal{N}(\overrightarrow{ba^o}))$$

by the continuity of \mathcal{N} of Proposition 6.2. Hence, each $\text{Axis}(g_i)$ passes a point p_i , and $\{p_i\}$ forms a convergent sequence in \mathbb{E} . By choosing a subsequence, we assume without loss of generality that $p_i \rightarrow p_\infty$ for $p_\infty \in \mathbb{E}$.

Denote $U_i := \mathcal{W}^{wu}(g_i)$, which is the span of $\text{Axis}(g_i)$ and r_i . To conclude (I)(i), we obtained a sequence $\{g_i\}$ satisfying the properties:

$$\begin{aligned} & \{\alpha_i\} \rightarrow \alpha, \{\beta_i\} \rightarrow \beta, \{\nu_i = \rho_i \times \alpha_i\} \rightarrow \nu, \\ & \{a_i\} \rightarrow a, \{r_i\} \rightarrow b, \{[\nu_i]\} \rightarrow [\nu], \{p_i\} \rightarrow p_\infty, \\ & \{\text{Cl}(\text{Axis}(g_i))\} \rightarrow \overline{[\nu]p_\infty[-\nu]}, \\ (16) \quad & \{U_i := \mathcal{W}^{wu}(g_i)\} \rightarrow U_\infty. \end{aligned}$$

The last property follows easily. We will be fixing the property of $\{g_i\}$ in this section here.

(I)(ii) *The next important step of the proof is that the coordinate changes required for g_i so that it becomes of form in (13) are uniformly bounded:*

We now introduce $h_i \in \text{Aut}(\mathbf{S}^3)$ coordinatizing \mathbf{S}^3 for each i . We choose h_i so that

$$(17) \quad h_i(a_i) = \mathbf{e}_1, h_i([\nu_i]) = \mathbf{e}_2, h_i(b_i) = \mathbf{e}_3, \text{ and } h_i(p_i) = \mathbf{e}_4,$$

and $\text{Axis}(g_i)$ is sent to

$$\overline{\mathbf{e}_2 \mathbf{e}_4 \mathbf{e}_2}^o.$$

By (16) and Lemma 2.1, $\{h_i\}$ can be chosen so that $\{h_i\}$ converges to a bi-Lipschitz map $h \in \text{Aut}(\mathbf{S}^3)$, uniformly in C^s -sense for any integer $s \geq 0$.

We can consider $\mathbf{v}_0(g_i) = \nu_i$ as a tangent vector at the origin O in E . The linearization $\mathcal{L}(h_i)$ of h_i restricted to E sends the unit-norm vector $\mathbf{v}_0(g_i)$ to a vector parallel to $(0, 1, 0)$. By post-composing with a projective map Φ_i fixing \mathbf{e}_i , $i = 1, \dots, 4$, we further modify h_i so that the differential of h_i sends the tangent vector $\mathbf{v}_0(g_i)$ at p_i to the tangent vector $(0, 1, 0)$ at \mathbf{e}_4 . Since $\{\mathbf{v}_0(g_i)\}$ is a convergent sequence of vectors, we can still assume that h_i is convergent in the above sense to $h \in \text{Aut}(\mathbf{S}^3)$.

The sequence $\{h_i\}$ is *uniformly bi-Lipchitz* in $\mathbf{d}_{\mathbf{S}^3}$; There exists a fixed positive constant C so that

$$(18) \quad \begin{aligned} C^{-1}\mathbf{d}_{\mathbf{S}^3}(h_i(x), h_i(y)) &\leq \mathbf{d}_{\mathbf{S}^3}(x, y) \leq C\mathbf{d}_{\mathbf{S}^3}(h_i(x), h_i(y)) \\ C^{-1}\mathbf{d}_{\mathbf{S}^3}(h_i^{-1}(x), h_i^{-1}(y)) &\leq \mathbf{d}_{\mathbf{S}^3}(x, y) \leq C\mathbf{d}_{\mathbf{S}^3}(h_i^{-1}(x), h_i^{-1}(y)) \end{aligned}$$

for all $x, y \in \mathbf{S}^n$, $i = 1, 2, \dots$.

The sequence of each quadruple of coordinate functions converges to a quadruple of coordinate functions of a new adopted coordinate system h where

$$(19) \quad h(a) = \mathbf{e}_1, h(b) = \mathbf{e}_3, h([\nu]) = \mathbf{e}_2 \text{ and } h(p) = \mathbf{e}_4$$

hold. Moreover, at the same time, we can assume that the linearization of the affine map h_i in E sends the vector ν_i to $(0, 1, 0)$ in V and, similarly, the linearization of h sends the vector ν to $(0, 1, 0)$ which follows as in the proof of Lemma 2.1 up to multiplying by a sequence of bounded linear maps of E to the sequence.

We can conjugate by a sequence of h_i of orientation-preserving bi-Lipschitz maps considered as an element of $\text{Aut}(\mathbf{S}^3)$ so that $h_i g_i h_i^{-1}$ is

of form

$$(20) \quad \hat{g}_i := \begin{bmatrix} \lambda(g_i) & 0 & 0 & 0 \\ 0 & 1 & 0 & k(g_i) \\ 0 & 0 & \frac{1}{\lambda(g_i)} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\lambda(g_i) > 1$ and the orientation preserving h_i is given by sending \mathbf{e}_1 to a_i , \mathbf{e}_2 to $[\nu_i]$, \mathbf{e}_3 to r_i , and \mathbf{e}_4 to p_i .

We note that

$$(21) \quad g_i = h_i^{-1} \circ \hat{g}_i \circ h_i;$$

hence, g_i acts like the standard form matrix in (20) up to coordination by h_i .

Let us define the compact set $\hat{\Lambda} := \mathcal{H} - (\mathsf{E} \cup \tilde{\Sigma})$, so-called stretched limit set. Moreover, $\hat{\Lambda} = \bigcup_{z \in \Lambda} \text{Cl}(\varepsilon(z))$ for the limit set Λ of Γ . We recall from Lemma 7.1:

- the unstable sphere \mathbf{S}_0^2 given by $x = 0$,
- the segment $\eta_+ := \overline{\mathbf{e}_1 \mathbf{e}_2 \mathbf{e}_{1-}}$ and
- the segment $\eta_- := \overline{\mathbf{e}_3 \mathbf{e}_{2-} \mathbf{e}_{3-}}$.

Then

- $h_i^{-1}(\eta_-) = \text{Cl}(\varepsilon(r_i))$ is the segment $\overline{r_i[-\nu_i]r_{i-}} \subset \hat{\Lambda}$ and
- $h_i^{-1}(\eta_+) = \text{Cl}(\varepsilon(a_i))$ is the segment $\overline{a_i[\nu_i]a_{i-}} \subset \hat{\Lambda}$.

(I)(iii) Now we estimate the entries of the above matrices.

We claim that in this coordinate system h_i

$$(22) \quad \mu(g_i) = k(g_i) :$$

we chose x in $\mathbf{B}(g_i(x) - x, \mathbf{v}_0(g_i))$ to be on $\text{Axis}(g_i)$ and the linearization $\mathcal{L}(h_i)$ of h_i restricted to E sends $g_i(x) - x$ and $\mathbf{v}_0(g_i)$ respectively to a vector parallel to $(0, 1, 0)$ and the vector $(0, 1, 0)$. Since the Lorentzian norm $\|\mathbf{v}_0(g_i)\| = 1$ holds, we obtain

$$g_i(x) - x = \mu(g_i)\mathbf{v}_0(g_i) \text{ and } \mathcal{L}(h_i)(g_i(x) - x) = \mu(g_i)(0, 1, 0).$$

$$k(g_i) = \mathcal{L}(h_i)(g_i(x) - x) \cdot (0, 1, 0)$$

implies (22).

Hence the positivity of the Margulis invariant is equivalent to $k(g_i) > 0$ as the coordinate change is orientation-preserving where $k(g_i)$ depends on the coordinate system.

The hyperbolic metric on \mathbb{S}_+ gives us the closed geodesic c_i in Σ_+ represented by g_i the hyperbolic length

$$l_{\mathbb{S}_+}(g_i) = \left| \log \frac{\lambda(g_i)}{\frac{1}{\lambda(g_i)}} \right| = 2|\log \lambda(g_i)|,$$

where we used cross ratios.

Recall that $\{g_i\}$ has a property (16):

Lemma 7.3. *For g_i obtained as above in (I)(i), and conjugating g_i by h_i as defined above to the form of (20), we have*

$$(23) \quad \lambda(g_i) \rightarrow +\infty, \quad k(g_i) \rightarrow +\infty, \quad \text{and} \quad \frac{k(g_i)}{\lambda(g_i)} \rightarrow 0.$$

Proof. Since $\{g_i\}$ restricts to a sequence of mutually distinct hyperbolic isometries in \mathbb{S}_+ with sequences of fixed points $\{a_i\} \rightarrow a$ and $\{r_i\} \rightarrow b$ and $a \neq b$, it follows that $\{\lambda(g_i)\} \rightarrow \infty$. (Suppose not. Then we see that $\{g_i\}$ is a bounded subset of $\mathrm{PSL}(2, \mathbb{R})$ by conjugating a_i and b_i to the standard position in $\mathrm{bd}\mathbb{S}_+$ by a sequence of bounded matrices in $\mathrm{PSL}(2\mathbb{R})$. Since Γ is discrete and $\mathcal{L}'|\Gamma$ is injective, this implies $\{g_i\}$ is finite. However, we assumed otherwise.)

By (8), we have $\mu(g_i) \rightarrow \infty$. Hence, the second limit follows by (22).

By the main corollary of [35], $\mu(g_i) \leq C'l(g_i)$ and hence $\mu(g_i) \leq 2C'|\log \lambda(g_i)|$ hold; we obtain the final limit. \square

Note that without the right conjugating h_i , we may not obtain the above.

(I)(iv) *We go back to our original compact set K from the beginning of the proof and cover it by compact convex sets that we can understand under the action of g_i :* Recall from (I)(ii) that we may assume by choosing subsequences of g_i satisfying

- $\{U_i\} \rightarrow U_\infty$, the *limit unstable sphere* a great 2-dimensional sphere again,
- $\{h_i^{-1}(\eta_-)\} \rightarrow \eta_-^\infty = \mathrm{Cl}(\varepsilon(b)) \subset \hat{\Lambda}$, and
- $\{h_i^{-1}(\eta_+)\} \rightarrow \eta_+^\infty = \mathrm{Cl}(\varepsilon(a)) \subset \hat{\Lambda}$.

Here

- η_+^∞ is a segment $\overline{a[\nu]a_-} \subset \hat{\Lambda}$ and
- η_-^∞ is a segment $\overline{b[-\nu]b_-} \subset \hat{\Lambda}$.

They are all tangent to $\mathrm{bd}\mathbb{S}_+$.

Note that K is a subset of $\mathcal{H} - \bigcup_{i \in \mathbb{Z}_+} h_i^{-1}(\eta_-)$ since $K \subset E \cup \tilde{\Sigma} = \mathcal{H} - \hat{\Lambda}$. There are two components H_a and H_{a_-} of $\mathcal{H} - U_\infty$ containing a or a_- respectively. The strategy is to cover K by balls bounded away from U_∞ and by balls that meets U_∞ precisely at the centers. These two can be understood easily by the use of Lemma 7.1 and conjugation by h_i that form a bounded uniformly convergent change of coordinates. We explain details:

(I)(iv-a) First, we try to obtain the covering. Since $\{U_i\}$ converges to U_∞ , we will cover K in the following manner:

- Let $K_\infty := K \cap U_\infty$. Assume that it is not empty. This is a compact subset of an open 3-dimensional hemisphere in $U_\infty - \eta_-^\infty$. We cover K_∞ by the interiors $\{\tilde{B}_i^o | i \in I_1\}$ of a finitely many balls $\tilde{B}_i \subset E \cup \tilde{\Sigma}$ with center in U_∞ for $i \in I_1$ of small radius $\epsilon_0 > 0$ in $\mathcal{H} - \hat{\Lambda}$. (We assume that balls \hat{B}_i of radius $2\epsilon_0$ with the same centers are still in $\mathcal{H} - \hat{\Lambda}$.)
- We split the balls and push the components B_i^+, B_i^- of $\tilde{B}_i - U_\infty$ for each $i \in I_1$ using a parameter of smooth maps preserving the foliation in \mathcal{H} by perpendicular segments to U_∞ so that
 - the images $C_i^+ \subset H_a, C_i^- \subset H_{a-}$ are still compact and convex,
 - for the join $J(C_i^+, C_i^-)$ of the each pair of two halves C_i^+, C_i^- , a so-called dumbbell, is still in $\hat{B}_i \subset E \cup \tilde{\Sigma}$, by displacing exclusively in the $2\epsilon_0$ -balls, and
 - $\{J(C_i^+, C_i^-)^o | i \in I_1\}$ covers K_∞ .

There is a positive number δ_0 so that

$$(24) \quad \mathbf{d}_{S^3}(C_i^+, U_\infty), \mathbf{d}_{S^3}(C_i^-, U_\infty) > \delta_0 \text{ for all } i \in I_1 \text{ and}$$

$$(25) \quad K_\infty \subset \bigcup_{i \in I_1} J(C_i^+, C_i^-)^o \subset E \cup \tilde{\Sigma}.$$

- We obtain a compact subset

$$K' := K - \bigcup_{i=1, \dots, m_0} J(C_i^+, C_i^-)^o \subset E \cup \tilde{\Sigma} - U_\infty = \mathcal{H} - \hat{\Lambda} - U_\infty.$$

Let $\epsilon_{K'} := \mathbf{d}_{S^3}(U_\infty, K')$. We can cover the set by $\epsilon_{K'}/2$ - \mathbf{d}_{S^3} -balls B_i for $i \in I_2$ where I_2 is a finite set and

$$(26) \quad \mathbf{d}_{S^3}(B_i, U_\infty) \geq \frac{\epsilon_{K'}}{2}, i \in I_2.$$

We divide I_2 into I_2^+ and I_2^- so that $j \in I_2^+$ if and only if $B_j \subset H_a$. We denote by $C_{j,i}^{(0)}$ for $j \in I_1$, the set $U_i \cap J(C_j^+, C_j^-)$.

(I)(iv-b) Now we study the balls that are bounded away from U_∞ . Since $U_i \rightarrow U_\infty$,

- There exists N_0 such that for $i > N_0$, there exists $\delta > 0$ such that

$$(27) \quad \mathbf{d}_{S^3} \left(U_i, \bigcup_{j \in I_2^+} B_j \cup \bigcup_{j \in I_2^-} B_j \cup \bigcup_{j \in I_1} C_j^+ \cup \bigcup_{j \in I_1} C_j^- \right) > \delta$$

by (24) and (26) since $U_i \rightarrow U_\infty$.

- For $i > N_0$, since we have that $h_i(U_i) = \mathbf{S}_0^2$,

$$(28) \quad \mathbf{d}_{\mathbf{S}^3} \left(\mathbf{S}_0^2, h_i \left(\bigcup_{j \in I_2^+} B_j \right) \cup h_i \left(\bigcup_{j \in I_2^-} B_j \right) \cup h_i \left(\bigcup_{j \in I_1} C_j^+ \right) \cup h_i \left(\bigcup_{j \in I_1} C_j^- \right) \right) > C^{-1}\delta.$$

- Now we act by \hat{g}_i . For any $\epsilon'' > 0$, there exists N'_0 so that for $i > N'_0$,

$$(29) \quad \mathbf{d}_{\mathbf{S}^3}^s \left(\{\mathbf{e}_1, \mathbf{e}_{1-}\}, \hat{g}_i \circ h_i \left(\bigcup_{j \in I_2^+} B_j \cup \bigcup_{j \in I_2^-} B_j \cup \bigcup_{j \in I_1} C_j^+ \cup \bigcup_{j \in I_1} C_j^- \right) \right) < \epsilon''$$

holds by Lemma 7.1 (d).

- Hence, for every $\epsilon'' > 0$, we have an integer $N'_0 > N_0$ so that for $i > N'_0$, the images of these sets under g_i go into the ϵ'' - $\mathbf{d}_{\mathbf{S}^3}$ -neighborhoods of a or a_- by (18). That is,

$$(30) \quad \mathbf{d}_{\mathbf{S}^3}^s \left(\{a, a_-\}, g_i \left(\bigcup_{j \in I_2^+} B_j \cup \bigcup_{j \in I_2^-} B_j \cup \bigcup_{j \in I_1} C_j^+ \cup \bigcup_{j \in I_1} C_j^- \right) \right) < \epsilon''.$$

Thus, we conclude

$$(31) \quad g_i \left(\bigcup_{j \in I_2^+} B_j \cup \bigcup_{j \in I_2^-} B_j \cup \bigcup_{j \in I_1} C_j^+ \cup \bigcup_{j \in I_1} C_j^- \right) \rightarrow \{a, a_-\}, \{a\}, \text{ or } \{a_-\}.$$

If $K_\infty = \emptyset$, then the reader can proceed to (I)(iv-d) now using $\epsilon_K = \mathbf{d}_{\mathbf{S}^3}(U_\infty, K)$. Thus, assume $K_\infty \neq \emptyset$ from now on.

(I)(iv-c) Now, we study the balls meeting U_∞ :

We have

$$(32) \quad \mathbf{d}_{\mathbf{S}^3} \left(\bigcup_{i \in I_1} J(C_i^+, C_i^-), \hat{\Lambda} \right) > \delta'$$

for some $\delta' > 0$ since $\bigcup_{i \in I_1} \hat{B}_i$ and $\hat{\Lambda}$ are both compact subsets of \mathcal{H} and are disjoint. Since $\text{Cl}(\varepsilon(r_i))$ is a subset of the compact set $\hat{\Lambda}$,

$$\mathbf{d}_{\mathbf{S}^3}(C_{j,i}^{(0)}, \text{Cl}(\varepsilon(r_i))) > \delta' \text{ for a fixed } \delta' > 0.$$

Thus,

$$\mathbf{d}_{\mathbf{S}^3}(h_i(C_{j,i}^{(0)}), \eta_{i-}) > C^{-1}\delta' > 0$$

holds for all j . Furthermore, the sequence of images under $\{\hat{g}_i\}$ of $h_i(C_{j,i}^{(0)})$ then converges to \mathbf{e}_2 by Lemma 7.1 uniformly with respect to

j . Hence, for every $\epsilon > 0$,

$$\mathbf{d}_{\mathbf{S}^3}^s(\hat{g}_i \circ h_i(C_{j,i}^{(0)}), \mathbf{e}_2) < \epsilon$$

holds if $i > N_0''$ for some N_0'' independent of j . By (18),

$$\mathbf{d}_{\mathbf{S}^3}^s(g_i(C_{j,i}^{(0)}), [\nu_i]) < C\epsilon$$

holds. Therefore, the sequence $\{g_i(C_{j,i}^{(0)})\}_{i \in \mathbb{Z}_+}$ converges to $\{[\nu]\}$ by (21).

Now, the sequence

- $\{g_i(J(C_+^j, C_{j,i}^{(0)}))\}_{i \in \mathbb{Z}_+} \rightarrow \overline{a[\nu]}$ by Lemma 2.3 and (30) and the above convergence since

$$g_i(J(C_+^j, C_{j,i}^{(0)})) = J(g_i(C_+^j), g_i(C_{j,i}^{(0)}))$$

for each i, j .

- Similarly, $\{g_i(J(C_-^j, C_{j,i}^{(0)}))\}_{i \in \mathbb{Z}_+} \rightarrow \overline{[\nu]a_-}$.

Since $J(C_+^j, C_-^j) = J(C_+^j, C_{j,i}^{(0)}) \cup J(C_-^j, C_{j,i}^{(0)})$,

$$\{g_i(J(C_+^j, C_-^j))\}_{i \in \mathbb{Z}_+} \rightarrow \eta_+^\infty = \overline{a[\nu]} \cup \overline{[\nu]a_-}$$

for each $j \in I_1$ holds by Proposition 2.2.

(I)(iv-d) We now come to the last step of the proof in case (I). If $g_i(K) \cap K \neq \emptyset$ for all i , then there has to be some fixed pair of balls B and B' in the above finite collection $I_1 \cup I_2$ of balls so that $g_i(B) \cap B' \neq \emptyset$ holds for infinitely many i . However,

$$\{g_i(B)\} \rightarrow \eta_+^\infty, \{a\}, \text{ or } \{a_-\} \subset \mathbb{S} \text{ with } \eta_+^\infty \cap \tilde{\Sigma} = \emptyset.$$

Thus, for each $\epsilon > 0$ and every ball B in the collection $I_1 \cup I_2$, there exists an integer I_0 such that

$$N_\epsilon(\eta_+^\infty) \supset g_i(B) \text{ if } i > I_0$$

by definition of the geometric convergence. However, $N_\epsilon(\eta_+^\infty) \cap B' = \emptyset$ holds for any ball B' in $I_1 \cup I_2$ for sufficiently small ϵ , and we obtain a contradiction.

(II) Now we consider the $a = r$ case. Choose an element $\gamma_0 \in \Gamma$ so that $\gamma_0(a) \neq a$ as Γ is not an elementary Fuchsian group acting on \mathbb{S}_+ .

Consider the sequence $\{\gamma_0 g_i\}$. For each $\epsilon > 0$ and sufficiently large i and a precompact neighborhood U in $\text{bd}_{\mathbb{S}} \mathbb{S}_+ - \{a\}$.

$$\gamma_0 g_i(U) \subset N_\epsilon(\gamma_0(a))$$

holds. Now consider the sequence $g_i^{-1}\gamma_0^{-1}$. Let V be a precompact neighborhood in $\text{bd}_{\mathbb{S}}\mathbb{S}_+ - \{\gamma_0(a)\}$. Since $\gamma_0^{-1}(V)$ is a precompact interval in $\text{bd}_{\mathbb{S}}\mathbb{S}_+ - \{a\}$ and for arbitrary $\epsilon > 0$,

$$g_i^{-1}(\gamma_0^{-1}(V)) \subset N_\epsilon(a) \text{ for } i > I_0$$

holds for some large I_0 . Thus, $\gamma_0(a)$ is the attractor point of $\{\gamma_0 g_i\}$ and a is the repellor point.

Since $\gamma_0 g_i(K \cup \gamma_0(K)) \cap (K \cup \gamma_0(K)) \neq \emptyset$ for infinitely many i , we are reduced to the case where $a \neq r$ by replacing g_i with $\gamma_0 g_i$ and K with another compact set $K \cup \gamma_0(K) \subset E \cup \tilde{\Sigma}$. Again, we obtain contradiction. \square

Remark 7.4. Let Γ be given an equivalence relation \sim of conjugacy. We can replace the condition of the proper-discontinuity of the action of Γ in E by the condition (*) given by

$$\begin{aligned} \mu(g) &> 0 \text{ for } g \in \Gamma - \{I\} \\ \mu : \Gamma - \{I\} / \sim &\rightarrow [0, \infty) \text{ is proper, and} \\ \mathcal{N} &\text{ is continuous.} \end{aligned}$$

which seemingly is slightly weaker to the condition of (8). Then we can show that this condition implies that Γ acts properly discontinuously on $E \cup \tilde{\Sigma}$.

Let Γ be a free group of rank ≥ 2 acting freely on E without parabolics. We can show the equivalence of the following conditions:

- (a) Γ acts properly and freely on E .
 - (b) The Margulis invariants of Γ satisfy equation (8) and \mathcal{N} is continuous.
 - (c) Our condition(*).
 - (d) Γ acts properly and freely on $E \cup \tilde{\Sigma}$.
- (b) \Rightarrow (c) and (d) \Rightarrow (a) are trivial. (a) \Rightarrow (b) is in [35], and (c) \Rightarrow (d) is in the proof of Proposition 7.2.

Remark 7.5. In the proof of Proposition 7.2, once we obtain some covering of K by compact convex balls, instead of the above precise ones, then we can just use Lemma 7.1 and using how U_i intersect with each of the convex balls and extracting subsequences, we can still obtain the proof. This is the shorter proof that the authors used in the earlier preprint.

8. THE PROOF OF TAMENESS.

Using the above notations, we note that $\tilde{\Sigma}/\Gamma$ is a closed surface of genus g and forms the boundary of the 3-manifold $M := (\mathsf{E} \cup \tilde{\Sigma})/\Gamma$ by Proposition 7.2 provided $\mathcal{L}(\Gamma) \subset \mathrm{SO}(2, 1)^o$. Therefore, M is a 3-manifold in general since Γ has an index ≤ 2 subgroup Γ' with the property $\mathcal{L}(\Gamma') \subset \mathrm{SO}(2, 1)^o$ and the fact that Γ acts freely on $\mathsf{E} \cup \tilde{\Sigma}$.

We now show that M is compact. The key idea is to isotopy some spheres into E and they have to bound compact 3-balls. We first assume that $\mathcal{L}(\Gamma) \subset \mathrm{SO}(2, 1)^o$ so that Γ acts on \mathbb{S}_+ honestly.

Proposition 8.1. *Each simple closed curve γ in $\tilde{\Sigma}$ bounds a simple disk in $\mathsf{E} \cup \tilde{\Sigma}$. Let c be a simple closed curve in Σ that is homotopically trivial in M . Then c bounds an imbedded disk in M .*

Proof. This is just Dehn's lemma. \square

We can find a collection of disjoint simple curves γ_i , $i \in \mathcal{J}'$, on $\tilde{\Sigma}$ for an index set \mathcal{J}' so that the following hold:

- $\bigcup_{i \in \mathcal{J}'} \gamma_i$ is invariant under Γ .
- $\bigcup_{i \in \mathcal{J}'} \gamma_i$ cuts $\tilde{\Sigma}$ into a union of open pair-of-pants P_k , $k \in K$, for an index set K . The closure of each P_k is a closed pair-of-pants.
- $\{P_k\}_{k \in K}$ is a Γ -invariant set.
- Under the covering map $\pi : \tilde{\Sigma} \rightarrow \tilde{\Sigma}/\Gamma$, each γ_i for $i \in \mathcal{J}'$ maps to a simple closed curve in a one-to-one manner and each P_k for $k \in K$ maps to an open pair-of-pants as a homeomorphism.

This is achieved by finding arcs in \mathbb{S}_+/Γ cutting it into disks. Recalling that \mathbb{S}_+/Γ is an open surface compactified to the compact surface Σ'_+ with boundary, we obtain a system of geodesic segments $\hat{\alpha}_i$, $i \in I$ for a finite set I , Σ'_+ cutting it into a union of disks each of which is bounded by six arcs, alternating triple of which are arcs coming from $\partial\Sigma'_+$ and the other alternating triples are geodesic segments ending $\partial\Sigma'_+$. We can assume that each $\hat{\alpha}_i$ is imbedded in \mathbb{S}_+/Γ cutting it into finitely many hexagons so that each ideal boundary component of \mathbb{S}_+/Γ meets $\hat{\alpha}_i$ and $\hat{\alpha}_j$ for some pair i, j , $i \neq j$.

This gives a system of geodesic arcs α_j , $j \in \mathcal{J}'$, in \mathbb{S}_+ for some infinite index set \mathcal{J}' which decomposes \mathbb{S}_+ into hexagons and forms a Γ -invariant set. We define $\alpha_{j-} := \mathcal{A}(\alpha_j)$ in \mathbb{S}_- for $j \in \mathcal{J}'$. We connect each endpoint $\delta_1(\alpha_j)$ and $\delta_2(\alpha_j)$ of α_j with its antipodal endpoint of α_{j-} by $\varepsilon(x)$.

$$\gamma_j := \alpha_j \cup \alpha_{j-} \cup \varepsilon(\delta_1(\alpha_j)) \cup \varepsilon(\delta_2(\alpha_j)), \quad j \in \mathcal{J}'.$$

(These form *crooked circles*.) We do this for each arc and obtain the above system of simple closed curves. By construction, $\{\gamma_i\}_{i \in J'}$ maps to a system of disjointly imbedded curves $\hat{\gamma}_1, \dots, \hat{\gamma}_{3g-3}$, and each of them has a simple closed lift in $\tilde{\Sigma}$ and hence has a trivial holonomy. See Figure 6. Also, the collection $\hat{\gamma}_1, \dots, \hat{\gamma}_{3g-3}$ decomposes the closed surface $\tilde{\Sigma}/\Gamma$ of genus g for some $g \geq 2$ into $2g-2$ pairs of pants P'_1, \dots, P'_{2g-2} . (Here g depends on the genus \tilde{g} of \mathbb{S}_+/Γ and the number of ideal boundary components.)

By trivial holonomy and Dehn's lemma, each $\hat{\gamma}_i$ bounds a disk D_i in M . By the 3-manifold topology of disk exchanges, we can choose D_1, \dots, D_{3g-3} to be mutually disjoint. (See [38].)

By combinatorics of the situation, the Cauchy completion of each component of $M - D_1 - \dots - D_{3g-3}$ is homeomorphic to a compact 3-ball in as it is bounded by a 2-sphere that is a union of a pair-of-pants P'_j and three of D_1, \dots, D_{3g-3} : We can lift the sphere into $E \cup \tilde{\Sigma}$ and being a sphere which can be pushed inside E , it bounds a compact 3-ball disjoint from other disks.

The image of the 3-ball is a compact 3-ball in M as well. Since M is a union of the closure of the components that are 3-balls identified with one another in disjoint disks, M is a compact 3-manifold. This implies also that M is homeomorphic to a solid handlebody of genus g ; that is, our manifold $(E \cup \tilde{\Sigma})/\Gamma$ is a compact 3-manifold and so its interior E/Γ is tame. This completes the proof of Theorem 1.2 provided $\mathcal{L}(\Gamma) \subset SO(2, 1)^\circ$. (Also, it follows that the rank of Γ equals g .)

In the general case when $\mathcal{L}(\Gamma)$ is not a subgroup of $SO(2, 1)^\circ$, the above work applies to $(E \cup \tilde{\Sigma})/\Gamma'$ double-covering $(E \cup \tilde{\Sigma})/\Gamma$. By Theorem 5.2 of Chapter 5 of [38], M is homeomorphic to a handlebody since Γ is a free group and $(E \cup \tilde{\Sigma})/\Gamma$ is an aspherical compact 3-manifold.

A *crooked plane* is a closed disk embedded in E constructed as follows: Take two null vectors v_1 and v_2 and respective parallel lines L_1 and L_2 meeting at a point x . Then they are on a timelike plane P . Take all timelike lines on P from x and take their union. It is a union of two cones C_1 and C_2 with vertex at x . We obtain two null half-planes $\mathcal{W}(v_1)$ and $\mathcal{W}(v_2)$. A crooked plane is the union $C_1 \cup C_2 \cup \mathcal{W}(v_1) \cup \mathcal{W}(v_2)$. (See [21] for details.)

A disk in E is an *almost crooked plane* if it agrees with a crooked plane in the complement of its compact subset. Also, its immersed or imbedded image is said to be *crooked plane* as well.

Proposition 8.2. *Let $\tilde{\Sigma}$ and Γ be as above. The Margulis spacetime E/Γ with a free fundamental group has a system of disks with boundary*




FIGURE 6. The arcs in \mathbb{S}_+ and an example of $\hat{\gamma}_i$ in the bold arcs.

in $E \cup \tilde{\Sigma}/\Gamma$ so that the closures of components of the complement are compact 3-balls and the disks are almost crooked planes.

Proof. First, assume $\mathcal{L}(\Gamma) \subset SO(2, 1)^o$. Each of our disks has the boundary of a crooked plane. In the Dehn's lemma, one can arbitrarily assign the tubular neighborhood of the boundary of each disk as long as it is transversal to $\tilde{\Sigma}/\Gamma$. Thus, following the above discussions, we obtain the proof in this case.

Suppose that $\mathcal{L}(\Gamma)$ is not in $SO(2, 1)^o$. Then we take a quotient $\Sigma' := \mathbb{S}_+/\mathcal{L}'(\Gamma)$ given by the projective action. The nonorientable Σ' admits a decomposition into hexagons as above. We obtain the induced decomposition on $\mathbb{S}_+ \cup \mathbb{S}_-$ and we extend the above construction to this situation and obtain the crooked circles and crooked planes.

As above $E \cup \tilde{\Sigma}/\Gamma'$ double-covers $E \cup \tilde{\Sigma}/\Gamma$ with the deck transformation group generated by a projective automorphism ϕ of order two. By choosing a collection for the quotient $\mathbb{S}_+/\mathcal{L}'(\Gamma)$ as in the beginning of the proof of Proposition 8.1, we may assume without loss of generality that ϕ sends each of the above crooked circles to itself or to a disjoint one by our construction. We can modify each almost crooked plane D not changing a neighborhood of ∂D in D so that $D \cap \phi(D) = \emptyset$ using the proof of Theorem 3 of [37]. Hence, we obtain a collection of ϕ -equivariant almost crooked planes that cuts M into 3-balls. \square

Corollary 8.3. *A Margulis spacetime without parabolics has a finite sided fundamental polyhedron in E .*

Proof. We take a union of finitely many of the 3-balls obtained by cutting along the disks in E . \square

9. ASYMPTOTICS OF MARGULIS SPACETIMES

Let N be a submanifold in a projective manifold M and is closed. If each boundary point of N has a convex open neighborhood B where $B - B \cap N$ is a convex open domain, then N is said to have *concave boundary*. Let Γ be a Margulis group without parabolics satisfying $\mathcal{L}(\Gamma) \subset \mathrm{SO}(2, 1)^o$. We show the existence of two convex hypersurfaces in E asymptotic to $\mathrm{bd}_{\mathbb{S}}\mathbb{S}_+$ and $\mathrm{bd}_{\mathbb{S}}\mathbb{S}_-$ respectively. We will use this to show that a concave boundary submanifold contains all the closed spacelike geodesics in E/Γ in the next section.

9.1. Supporting hyperplanes. Let \mathcal{H} be the set of pairs (x, h) where x is a point of the limit set Λ in $\mathrm{bd}_{\mathbb{S}}\mathbb{S}_+$ and a 2-dimensional hemisphere h in \mathcal{H} containing x in its boundary ∂h . We give the Hausdorff convergence topology to this space. γ in Γ acts by sending (x, h) to $(\gamma(x), \gamma(h))$.

Proposition 9.1. *For each point x of the limit set Λ of Γ in \mathbb{S} in $\mathrm{bd}_{\mathbb{S}}\mathbb{S}_+$, there exists a unique 2-dimensional hemisphere h_x depending continuously on x so that h_x contains the fixed line $\mathrm{Axis}(g) \subset \mathbb{R}^3$ of an element of $g \in \Gamma$ if x is an attracting fixed point of g . Moreover, there exists a map $\tau : \Lambda \rightarrow \mathcal{H}$ sending x to (x, h_x) that is continuous and the map is Γ -equivariant; that is,*

$$\tau(\gamma(x)) = (\gamma(x), \gamma(h_x)) \text{ for } \tau(x) = (x, h_x), \gamma \in \Gamma.$$

Proof. If x is an attracting fixed point of $\gamma \in \Gamma$, we let h_x be the 2-hemisphere containing the invariant line $\mathrm{Axis}(\gamma) \subset \mathsf{E}$ of γ and x as the stabilizer subgroup of Γ is infinite cyclic and is uniquely determined by x by the elementary Fuchsian group theory. Then h_x is γ -equivariant and ∂h_x contains x, x_-, s, s_- in \mathbb{S} for the saddle-type fixed points s and s_- and the antipode x_- of x . Thus, τ is defined for the set of attracting fixed points of Γ .

The inverse image in $\mathbb{U}\mathbb{S}_+$ of $\mathbb{U}_{\mathrm{rec}}(\mathbb{S}_+/\Gamma)$ is precisely the set of geodesics with endpoints in Λ . (See Lemma 4.3 and the proof of Lemma 1.3 of [35].) Recall that $\Lambda^* = \Lambda \times \Lambda - \Delta$ consists of elements (x, y) where x is the ending point and y is the starting point of a nonwandering geodesic l .

For convenience, we use the notation from Proposition 6.2

$$L(x, y) := \mathcal{N}(\overrightarrow{yx})$$

with the orientation forgotten. Using Proposition 6.2, we define a function $\tau(x, y) : \Lambda^* \rightarrow \mathcal{H}$ by

$$(x, y) \mapsto (x, h(x, y))$$

for the hemisphere $h(x, y) = \mathcal{H} \cap \mathbf{S}^2$ for the span \mathbf{S}^2 of x and $L(x, y)$. Note here that $L(x_g, y_g) = \text{Axis}(g)$ if x_g and y_g are fixed points of g in $\text{bd}_{\mathbb{S}_+} \mathbb{S}_+$ ordered appropriately, and $h(x, y)$ is contained in the unstable subspace of g .

Notice that $\tau(a, y)$ is independent of y , $y \in \Lambda - \{a\}$ if a is an attracting fixed point of an element of Γ with the repelling fixed point r in Λ and equals $\tau(a, r)$ by the main result of Charette, Goldman, and Jones [14]: $L(a, y)$ has to lie in the unstable plane $h(a, r)^o$ the line $L(a, r)$ going to a closed geodesic in $L(a, y)$ has to go to a line spiralling to the closed geodesic in E/Γ . This fact is subsumed under Lemma 9.3.

We state a fact we need.

Lemma 9.2. *Let $\{(x_i, y_i) \in \Lambda^*\}$ be a sequence. Suppose that $\{x_i\} \rightarrow x$, $\{y_i\} \rightarrow y$. Then*

$$\{L(x_i, y_i)\} \rightarrow L(x, y), \{\tau(x_i, y_i)\} \rightarrow \tau(x, y) \text{ as } i \rightarrow \infty.$$

Proof. This follows by the fact that \mathcal{N} is continuous by Proposition 6.2. \square

Lemma 9.3. *Let $(x, y) \in \Lambda^*$. Then $\tau(x, y)$ is independent of $y \in \Lambda - \{x\}$, and $h(x, y) \not\subset \mathbb{S}$ for every $(x, y) \in \Lambda^*$.*

Proof. Suppose that x is an endpoint of two nonwandering geodesics l with the starting point y and another one l' with the starting point y' .

If $(x, y) \in \Lambda^*$, then $L(x, y)$ is a line in E by Proposition 6.2; $h(x, y)$ is not in \mathbb{S} .

Recall that $\Sigma''_+ = \Omega_+/\Gamma$ is a compact hyperbolic surface with geodesic boundary and the nonwandering geodesics are always in it. Choose a sequence $\{p_i\}$ of points \overline{xy}^o so that $p_i \rightarrow x$. There exists a sequence $\{\gamma_i^{-1}\}$ of elements of Γ so that $\gamma_i^{-1}(p_i) \in F$ for a fixed compact fundamental domain F of Ω_+ under Γ .

By choosing a subsequence, we may assume that $\{p_i\} \rightarrow p_\infty \in F$ and $\{\gamma_i^{-1}(\text{Cl}(l))\}$ converges to a segment s_∞ passing p_∞ and ending at two points of Λ . Let l_∞ denote the interior of s_∞ in Ω_+ . Let x_∞ and y_∞ denote the endpoints so that we may assume without loss of generality that $\{\gamma_i^{-1}(x)\} \rightarrow x_\infty$ and $\{\gamma_i^{-1}(y)\} \rightarrow y_\infty$.

Since Γ is not elementary, we may assume that $\{x, y\} \cap \{x_\infty, y_\infty\} = \emptyset$ by multiplying by $\gamma_i^{-1} \in \Gamma$ if necessary.

Since $\{\gamma_i^{-1}(x)\} \rightarrow x_\infty$ and $\{\gamma_i^{-1}(y)\} \rightarrow y_\infty$,

$$\{\text{Cl}(L(\gamma_i^{-1}(x), \gamma_i^{-1}(y))) = \gamma_i^{-1}(\text{Cl}(L(x, y)))\} \rightarrow \text{Cl}(L(x_\infty, y_\infty))$$

holds by continuity of \mathcal{N} by Proposition 6.2. Hence,

$$(33) \quad \{\gamma_i^{-1}(h(x, y)) = h(\gamma_i^{-1}(x), \gamma_i^{-1}(y))\} \rightarrow h(x_\infty, y_\infty)$$

holds.

By using perpendicular geodesics in the hyperbolic space \mathbb{S}_+ to the geodesic l , we can show that y_∞ is the attractor point of the sequence $\{\gamma_i^{-1}\}$ since γ_i^{-1} sends any precompact open set in $\text{bd}_{\mathbb{S}}\mathbb{S}_+ - \{x\}$ to an arbitrarily small neighborhood of y_∞ for sufficiently large i . Also, x is the repellor point of the sequence $\{\gamma_i^{-1}\}$ since γ_i sends any precompact open set in $\text{bd}_{\mathbb{S}}\mathbb{S}_+ - \{y_\infty\}$ to an arbitrarily small neighborhood of x for sufficiently large i using geodesics perpendicular to $\gamma_i^{-1}(l)$. Therefore, for an attracting fixed point a_i and repelling fixed point r_i of γ_i^{-1} for each i , it follows that $\{a_i\} \rightarrow y_\infty$ and $\{r_i\} \rightarrow x$ by Lemma 4.2 since $x \neq y_\infty$. $\overline{r_i a_i} \rightarrow \overline{x y_\infty}$ implies

$$(34) \quad \{\text{Cl}(\text{Axis}(\gamma_i^{-1})) = \text{Cl}(L(r_i, a_i))\} \rightarrow \text{Cl}(L(x, y_\infty)),$$

$$(35) \quad \{h(r_i, a_i)\} \rightarrow h(x, y_\infty).$$

From this, $y' \neq x$ and $y' \in \text{bd}_{\mathbb{S}}\mathbb{S}_+$ show

$$(36) \quad \gamma_i^{-1}(y') \rightarrow y_\infty,$$

$$(37) \quad \{\gamma_i^{-1}(h(x, y')) = h(\gamma_i^{-1}(x), \gamma_i^{-1}(y'))\} \rightarrow h(x_\infty, y_\infty).$$

Since $\partial h(x, y)$ is a great circle tangent to $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ at x and so is $\partial h(x, y')$, we obtain $\partial h(x, y) = \partial h(x, y')$. And $\gamma_i^{-1}(\partial h(x, y'))$ is tangent to $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ at $\gamma_i^{-1}(x)$. Thus, $\{\gamma_i^{-1}(\partial h(x, y'))\}$ converges to a great circle $\mathbf{S}_{x_\infty}^1$ tangent to $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ at x_∞ in \mathbb{S} .

Choose a point q in the interior of the 2-hemisphere corresponding to $h(x, y')$ bounded away from $\partial h(x, y) = \partial h(x, y')$:

$$\mathbf{d}_{\mathbf{S}^3}(q, \partial h(x, y)) \geq C_0, C_0 > 0.$$

Suppose that $q \in \mathcal{H} - h(x, y_\infty)$ for contradiction. Then q is uniformly bounded away from $h(r_i, a_i)$ for $i > I_0$ for a fixed integer I_0 by (34). That is, there exists a constant $C_1, C_1 > 0$, such that

$$\mathbf{d}_{\mathbf{S}^3}(q, h(r_i, a_i)) > C_1.$$

We claim that $\{\gamma_i^{-1}(q)\} \rightarrow y_\infty$ or $\{\gamma_i^{-1}(q)\} \rightarrow y_{\infty,-}$:

We write γ_i^{-1} in the matrix of the form of (13) with respect to coordinate change automorphisms $h_i \in \text{Aut}(\mathbf{S}^3)$ so that $h_i \gamma_i^{-1} h_i^{-1}$ is of form

$$(38) \quad \hat{\gamma}_i^{-1} := \begin{bmatrix} \lambda(\gamma_i^{-1}) & 0 & 0 & 0 \\ 0 & 1 & 0 & k(\gamma_i^{-1}) \\ 0 & 0 & \frac{1}{\lambda(\gamma_i^{-1})} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

where $\lambda(\gamma_i^{-1}) > 1, k(\gamma_i^{-1}) > 0$ and

$$h_i(a_i) = \mathbf{e}_1, h_i(\hat{s}_i) = \mathbf{e}_2, h_i(r_i) = \mathbf{e}_3, h_i(\hat{p}_i) = \mathbf{e}_4,$$

where we choose $\hat{p}_i \in \text{Axis}(\gamma_i^{-1})$ so that \hat{p}_i is uniformly bounded away from \mathbb{S} by (34), and \hat{s}_i is the saddle-type fixed point or the point corresponding to an end point of $\text{Axis}(\gamma_i^{-1})$ in \mathbb{S} . As above, h_i is uniformly bi-Lipschitz with respect to $\mathbf{d}_{\mathbb{S}^3}$ and can be assumed to be convergent. Also, $h(r_i, a_i)$ is the image of the 2-hemisphere in \mathcal{H} given by \mathbf{S}_0^2 of Lemma 7.1 under h_i^{-1} . (We are exactly in the situation of the part (I) of the proof of Proposition 7.2 in Section 7.) As before, $k(\gamma_i^{-1}) \leq C \log \lambda(\gamma_i^{-1})$ holds for the terms in the matrices of γ_i^{-1} by Lemma 7.3. By the first item of Lemma 7.1, we obtain

$$\{\gamma_i^{-1}(q)\} \rightarrow y_\infty \text{ or } \{\gamma_i^{-1}(q)\} \rightarrow y_{\infty-}$$

since h_i are uniformly bi-Lipschitz and

$$\mathbf{d}_{\mathbb{S}^3}(q, h(r_i, a_i)) > C_1 \text{ for } i > I_0 \text{ and a constant } C_1 > 0.$$

The proof is entirely same as in the part (I)(iv-b) of the proof of Proposition 7.2.

Since $\{\gamma_i^{-1}(q)\} \rightarrow y_\infty$ or $y_{\infty-} \in \mathbb{S}$ and $\{\gamma_i^{-1}(\partial h(x, y'))\}$ converges to a great circle $\mathbf{S}_{x_\infty}^1$ tangent to $\text{bd}_{\mathbb{S}}\mathbb{S}_+$ at x_∞ in \mathbb{S} , it follows that $\{\gamma_i^{-1}(h(x, y'))\}$ geometrically converges to a 2-hemisphere in \mathbb{S} by Lemma 2.3 up to a choice of subsequence. We obtained a contradiction to (36).

Therefore, $q \in h(x, y_\infty)$ and $h(x, y') = h(x, y_\infty)$ by geometry.

Similarly, we have $h(x, y) = h(x, y_\infty)$ by (33), and we conclude that $h(x, y) = h(x, y')$.

□

Now we complete the proof of Proposition 9.1: We have defined the function $\tau : \Lambda \rightarrow \mathcal{H}$ and the 2-hemisphere $h(x)$ for any $x \in \Lambda$ by

$$\tau(x) := \tau(x, y), h(x) := h(x, y) \text{ for any } y \in \Lambda - \{x\}.$$

Now, we show that τ is continuous.

Now suppose that $\{x_i\}$ is a sequence of arbitrary points in Λ so that $\{x_i\} \rightarrow x$. We choose $y_i \in \Lambda$ for each i so that $\{y_i\} \rightarrow y \in \Lambda$ for $y \neq x$. Then $\{h(x_i, y_i)\} \rightarrow h(x, y)$ holds by Lemma 9.2 and therefore, $\{\tau(x_i)\} \rightarrow \tau(x)$ holds by the independence of Lemma 9.3. Therefore, τ is a continuous function. □

9.2. Domains. For each $i \in \mathcal{J}$, let Δ_i be an open \mathbf{g}_i -invariant triangle disjoint from Ω_+ with an edge \mathbf{l}_i where \mathbf{g}_i is a corresponding element of Γ acting on \mathbf{l}_i . Hence its attracting and repelling fixed points a_i, r_i , at the ends and a saddle-type fixed point s_i are vertices of Δ_i . We chose

the triangle $\Delta_i \subset \mathcal{R}_i$ for the rectangular domain \mathcal{R}_i defined in Section 5.

In \mathbb{S} , at each point x of Λ , there is a closed 2-hemisphere $H_x \subset \mathbb{S}$ supporting the convex hull of Λ . We form an intersection $D(\Lambda) = \bigcap_{x \in \Lambda} H_x^o \subset \mathbb{S}$. Then $D(\Lambda)$ is an open properly convex domain in \mathbb{S} :

$$(39) \quad D(\Lambda) = \bigcup_{i \in \mathcal{J}} \Delta_i \cup \mathbb{S}_+ \subset \tilde{\Sigma}.$$

Hence, $D(\Lambda) \subset \tilde{\Sigma}$ holds. Moreover, $D(\Lambda)/\Gamma$ is an open surface obtained from Σ_+ by adding open elementary annuli $\Delta_i/\langle \mathbf{g}_i \rangle$ in corresponding annuli A_j . Thus, the closure of $D(\Lambda)/\Gamma$ in Σ is a compact surface in Σ with boundary components $\mathbf{c}_j \subset A_j$ for each $j = 1, \dots, b$.

Defining $D(\mathcal{A}(\Lambda)) = \mathcal{A}(D(\Lambda))$, we obtain closed curves $\mathbf{c}'_j \subset A_j$ for each $j = 1, \dots, b$ that are boundary components of $D(\mathcal{A}(\Lambda))/\Gamma$ in Σ . Note that $(\tilde{\Sigma} - D(\Lambda) - D(\mathcal{A}(\Lambda)))/\Gamma$ is a union of compact annuli in each of A_j , $j = 1, \dots, b$. We denote each component annulus by B_j so that $B_j \subset A_j$ where $\partial B_j = \mathbf{c}_j \cup \mathbf{c}'_j$. (Each B_j is a sum of two elementary annuli and is bounded by nonprincipal boundary components.)

Lemma 9.4. *Let N_Γ be the set of points where the nonwandering spacelike geodesics passes in E/Γ . Then for any finite index subgroup Γ'' of Γ , the natural map*

$$N_{\Gamma''} \subset E/\Gamma'' \rightarrow N_\Gamma \subset E/\Gamma$$

is a finite covering map.

Proof. If the ω -limits of a spacelike geodesic l in E/Γ'' is compact, then so is the one for its image geodesic l in E/Γ . Conversely, the inverse image of the compact ω -limit of a spacelike geodesic l in E/Γ has a compact inverse image and the lifting spacelike geodesic l must have the ω -limit in it. The same argument works for α -limits. Thus, for a covering $p_1 : E/\Gamma'' \rightarrow E/\Gamma$, we obtain $p_1^{-1}(N_\Gamma) = N_{\Gamma''}$, which implies the conclusion. \square

The boundary of a convex open domain in E is *asymptotic* to a set K in \mathbb{S}_+ if every unbounded sequence of points in it has all limit points in K .

Corollary 9.5. *Let Γ be a Margulis spacetime without parabolics. In E , there exists a Γ -invariant nonempty convex open domain \mathcal{D} whose boundary in E is asymptotic to $\text{bd}_{\mathbb{S}} D(\Lambda)$, homeomorphic to a circle. There exists another Γ -invariant convex open domain \mathcal{D}' whose boundary in E is asymptotic to $\mathcal{A}(\text{bd}_{\mathbb{S}} D(\Lambda))$ so that the closures of \mathcal{D} and*

\mathcal{D}' are disjoint. Moreover, every nonwandering spacelike geodesic is contained in a manifold

$$(\mathsf{E} - \mathcal{D} - \mathcal{D}')/\Gamma$$

with concave boundary.

Proof. First, we work with the index ≤ 2 subgroup Γ' of Γ with $\mathcal{L}(\Gamma') \subset \mathrm{SO}(2, 1)^\circ$. Let Λ denote the limit set of Γ' . For each $x \in \Lambda$, take the open half-space $\mathcal{H}_x \subset \mathsf{E}$ bounded by $h(x, y)$ for some $y \in \Lambda - \{x\}$ and containing Λ in its closure in the 3-hemisphere \mathcal{H} . Then we take $\mathcal{D} := \bigcap_{x \in \Lambda} \mathcal{H}_x$. The domain is a properly convex open domain since it has at least three supporting hyperplanes and \mathbb{S} in a general position.

We show that \mathcal{D} is not empty: That is, we claim that there exists a compact convex set $K \subset \mathsf{E}$ so that $\mathcal{H}_x \cap K \neq \emptyset$ for all $x \in \Lambda$ and hence there exists

$$z \in K \cap \bigcap_{x \in \Lambda} \mathrm{Cl}(\mathcal{H}_x)$$

by compactness. Then the join $J(z, \mathrm{bd}_{\mathbb{S}}\mathbb{S}_+)$ is a subset of $\mathrm{Cl}(\mathcal{H}_x)$ for every $x \in \Lambda$. Hence the topological interior of the join $J(z, \mathrm{bd}_{\mathbb{S}}\mathbb{S}_+)$ is a nonempty subset of \mathcal{D} .

Suppose that such a compact set K does not exist. Then there exists a sequence of 2-hemispheres $h(x_i, y_i), x_i, y_i \in \Lambda$ so that it bounds an open half-space \mathcal{H}_{x_i} in E and $\{\mathcal{H}_{x_i} \cap K = \emptyset\}$ for each compact set $K \subset \mathsf{E}$ if $i > I_K$ for an integer I_K depending on K . Since Λ is compact, this is equivalent to the statement that $\{h(x_i, y_i)\}$ converges to a 2-hemisphere in \mathbb{S} up to a choice of a subsequence. Let x_∞ be a limit point of the sequence $\{x_i\}$. Then $\tau(x_\infty)$ is a pair (x_∞, h_∞) for a 2-hemisphere h_∞ in \mathbb{S} by the continuity of τ contradicting Lemma 9.3.

Let $\mathrm{bd}_{\mathsf{E}}(\bigcap_{x \in \Lambda} \mathcal{H}_x)$ be the topological boundary in E . The closure of $\mathcal{D} = \bigcap_{x \in \Lambda} \mathcal{H}_x$ is a convex compact subset of \mathbf{S}^3 , and

$$\mathrm{bdCl}(\mathcal{D}) = \mathrm{Cl}(D(\Lambda)) \cup \mathrm{bd}_{\mathsf{E}}\left(\bigcap_{x \in \Lambda} \mathcal{H}_x\right).$$

Thus, every unbounded sequence of points in $\mathrm{bd}_{\mathsf{E}}(\bigcap_{x \in \Lambda} \mathcal{H}_x)$ will have limit points in $D(\Lambda)$. Since the limit points cannot be in $D(\Lambda)^\circ$, they are in $\mathrm{bd}_{\mathbb{S}}D(\Lambda)$.

For each \mathcal{H}_x , we can take $(\mathsf{E} - \mathcal{H}_x)^\circ$ and take the intersection over all $x \in \Lambda$. This gives us

$$\mathcal{D}' := \bigcap_{x \in \Lambda} (\mathsf{E} - \mathcal{H}_x)^\circ,$$

which is again a properly convex open domain and is nonempty. Also,

$$\text{bd}\mathcal{D}' = \text{Cl}(\mathcal{A}(\mathcal{D}(\Lambda))) \cup \text{bd}_{\mathbb{E}} \bigcap_{x \in \Lambda} (\mathbb{E} - \mathcal{H}_x)^o.$$

These follow as before for \mathcal{D} .

First, notice $\mathcal{D} \cap \mathcal{D}' = \emptyset$. If $\text{Cl}(\mathcal{D}) \cap \text{Cl}(\mathcal{D}') \neq \emptyset$ holds, then it is a compact convex subset of \mathbb{E} as the boundary lies in \mathbb{S} and hypersurfaces in \mathbb{E} and the fact that

$$\text{Cl}(D(\Lambda)) \cap \text{Cl}(\mathcal{A}(D(\Lambda))) = \emptyset.$$

Since $\text{Cl}(\mathcal{D}) \cap \text{Cl}(\mathcal{D}')$ is also Γ' -invariant, the Brouwer fixed-point theorem implies a contradiction that the intersection contains a fixed point for each element $g \in \Gamma'$. Hence, $\text{Cl}(\mathcal{D}) \cap \text{Cl}(\mathcal{D}') = \emptyset$.

Every nonwandering spacelike geodesic equals $L(x, y)$ for $(x, y) \in \Lambda^*$ by Theorem 1 of [34]. Since $L(x, y)$ is in $h(x, y)$ for some $(x, y) \in \Lambda^*$, and $h(x) \cap (\mathcal{D} \cup \mathcal{D}') = \emptyset$ holds, the result follows provided $\Gamma = \Gamma'$.

Suppose now that $\Gamma - \Gamma' \neq \emptyset$. Let $\phi \in \Gamma - \Gamma'$. Then ϕ induces an inner automorphism $g \mapsto g^\phi = \phi g \phi^{-1} : \Gamma' \rightarrow \Gamma'$. Thus, if $(x, h(x))$ is fixed by g , then $(\phi(x), \phi h(x))$ is fixed by g^ϕ with $\phi(x)$ in $\text{bd}_{\mathbb{S}} \mathbb{S}_-$ and is uniquely determined by g^ϕ . We obtain $\phi(h(x)) = h(\mathcal{A}(\phi(x)))$ since $\phi(h(x))$ contains $\text{Axis}(g^\phi) = \phi(\text{Axis}(g))$. Thus,

$$(\mathcal{A}(\phi(x)), \phi(h(x))) = \tau(\mathcal{A}(\phi(x))).$$

Since ϕ sends $h(x)$ to $\phi(h(x))$ but sends $\text{bd}_{\mathbb{S}} \mathbb{S}_+$ to $\text{bd}_{\mathbb{S}} \mathbb{S}_-$, it follows that ϕ sends the component \mathcal{H}_x^o of $\mathbb{E} - h(x)$ to the component $(\mathbb{E} - \mathcal{H}_{\mathcal{A}(\phi(x))})^o$ of $\mathbb{E} - \phi(h(x))$ for each fixed point x of an element of Γ . We obtain

$$\phi(\mathcal{H}_x^o) = (\mathbb{E} - \mathcal{H}_{\mathcal{A}(\phi(x))})^o \text{ for } x \in \Lambda.$$

Since the domain \mathcal{D} is determined by $h(x)$ for $x \in \Lambda$, we obtain $\phi(\mathcal{D}) = \mathcal{D}'$ and $\phi(\mathcal{D}') = \mathcal{D}$.

The last statement follows by Lemma 9.4. □

Remark 9.6. These domains are the invariant domains found by Mess [43] for the closed surface groups. See Barbot [3] and Bonsante [8] for details of Mess's work around this. Also, Barbot had shown the fact that the supporting hyperplanes contain all nonwandering geodesics.

10. GEOMETRIC FINITENESS

We define geometric finiteness and show that the Margulis spacetimes are geometrically finite. The geometrical finiteness here of course is already discovered by Goldman, Labourie, and Margulis [35], [34], and we only restate it in this form.

We start with

Theorem 10.1 (Scott [46]). *If M is an orientable 3-manifold and $\pi_1(M)$ is finitely generated, then there exists a compact submanifold N of M such that the inclusion homomorphism induces an isomorphism $\pi_1(N) \rightarrow \pi_1(M)$.*

If M is aspherical, as in Margulis spacetimes, then we can construct a compact submanifold so that the inclusion map $N \rightarrow M$ is a homotopy equivalence. We say that a compact submanifold with properties of N is a *core* of M . For E/Γ with a free group Γ of a finite rank, there exists a core.

Definition 10.2 (Geometric finiteness). Let Γ be the orientation-preserving Lorentzian finitely generated isometry group acting on the Lorenzian space E freely and properly discontinuously. Let E/Γ be given an arbitrary complete path-metric δ and a core M_0 . Let \mathcal{R} denote the set of all nonwandering geodesics in E/Γ . E/Γ is *geometrically finite* if

$$\bigcup_{c \in \mathcal{R}} c \subset N_C^\delta(M_0)$$

for a C - δ -neighborhood of M_0 for a constant $C > 0$.

In other words, for each pair of a nonwandering geodesic c and a point x on c in E/Γ , x is of a uniformly bounded δ -distance from M_0 ; that is, $\delta(x, M_0) \leq C$ for a constant C independent of x and c .

Actually, since M_0 is compact, we can replace M_0 by a base point x_0 in the above definition.

Lemma 10.3. *Fix an arbitrary point x_0 and an arbitrary Euclidean metric d_E on E considered as an affine space. A Margulis spacetime E/Γ is geometrically finite if and only if there exists a constant $C' > 0$ such that for every pair of a geodesic l mapping to a nonwandering geodesic in E/Γ and a point y on l in E , there exists $\gamma \in \Gamma$ so that $d_E(x_0, \gamma(y)) < C'$.*

Proof. Let $C > 0$ be the uniform bound for the path-metric δ of E/Γ used in Definition 10.2. A δ -ball of radius C in E maps to a bounded ball in E in the Euclidean metric d_E since it is compact. The Euclidean diameters of the δ -balls of radii equal to C with centers in a compact set K are bounded above by a constant $C' > 0$.

Let l be a geodesic with $y \in l$ in the premise. Suppose that l maps to a nonwandering geodesic c and y to a point $x \in c$. Then x is of the δ -distance bounded above by C from a point $x' \in M_0$ by definition of the geometric finiteness. Let F be the compact fundamental domain of M_0 in E . Assume without loss of generality that $x_0 \in F$. Choose

$\gamma \in \Gamma$ so that $\gamma(y)$ is of the δ -distance $\leq C$ from $x' \in F$. Thus, $\gamma(y)$ is of the Euclidean distance $\leq C'$ from x_0 .

Conversely, a Euclidean ball with center x_0 of radius C' has a δ -diameter uniformly bounded above. The proof of the converse is very similar to the above. \square

Theorem 10.4. *A Margulis spacetime E/Γ without parabolics is geometrically finite.*

Proof. Assume first $\mathcal{L}(\Gamma) \subset \mathrm{SO}(2, 1)^o$. Let x_0 be a base point in E . Consider the compact fundamental domain \hat{F} of $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ under the deck transformation group to $\mathbb{U}_{\mathrm{rec}}\Sigma_+$ which covers a closed subset of the compact set $\mathbb{U}\Sigma''_+$. Then $\pi_E \circ \tilde{N}(\hat{F})$ gives us a set A of geodesics of bounded distances from x_0 since it is an image of a compact set: that is, $\{\mathbf{d}_{\mathbb{S}^3}(x_0, l) | l \in A\} < C$ for a constant C .

Let \mathbf{u} denote the unit vector at t in E tangent to a geodesic l mapping to a nonwandering one in E/Γ . Then (t, \mathbf{u}) corresponds to (t', \mathbf{u}') for a point $t' \in \mathbb{S}_+$ and \mathbf{u}' on $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ via $\pi_E \circ \tilde{N}$ by Proposition 6.2. Since \hat{F} is a fundamental domain, $(\gamma(t'), D\gamma(\mathbf{u}'))$ is in the bounded set \hat{F} for a deck transformation γ and the differential $D\gamma$. Thus, $(\gamma(g), D\gamma(\mathbf{u}))$ is in the image of $\pi_E \circ \tilde{N}(\hat{F})$, a compact subset. By Lemma 10.3, it follows that E/Γ is geometrically finite.

Assume now that $\mathcal{L}(\Gamma)$ is not a subset of $\mathrm{SO}(2, 1)^o$. The full general case follows by Lemma 9.4. \square

Theorem 10.5. *There exists a core in a Margulis spacetime containing all nonwandering spacelike geodesics.*

Proof. Let Γ be the finitely generated free group of rank ≥ 2 acting freely and properly discontinuously on E . We construct $\tilde{\Sigma}$ as above.

First, assume $\mathcal{L}(\Gamma)$ is a subset of $\mathrm{SO}(2, 1)^o$.

Let M denote the compact handlebody $(E \cup \tilde{\Sigma})/\Gamma$ so that M^o is the Margulis spacetime E/Γ . We denote by \tilde{M} the space $E \cup \tilde{\Sigma}$. Recall the domains \mathcal{D} and \mathcal{D}' from the proof of Corollary 9.5. Then considering the closure $\hat{\mathcal{D}} = \mathrm{Cl}_{\tilde{M}}(\mathcal{D})$ of \mathcal{D} in \tilde{M} , it follows that Γ acts properly discontinuously on $\hat{\mathcal{D}}$ as well.

The topological boundary $\mathrm{bd}_{\tilde{M}}\hat{\mathcal{D}}$ in \tilde{M} is a closed invariant subset of E and is a connected properly imbedded 2-dimensional submanifold invariant under Γ . Since $\mathrm{bd}_{\tilde{M}}\hat{\mathcal{D}}$ equals $\mathrm{bd}\mathcal{D} - D(\Lambda)$, $\mathrm{bd}_{\tilde{M}}\hat{\mathcal{D}}$ is homeomorphic to a 2-cell. However, under the projection $p : \tilde{M} \rightarrow M$, it goes to a compact closed connected properly imbedded surface Σ_1 bounding the image of $\hat{\mathcal{D}}$. To explain more, $M_1 := \hat{\mathcal{D}}/\Gamma$ is imbedded in M and

is homotopy equivalent to M and the topological boundary $\text{bd}_M M_1$ in M is the compact surface Σ_1 .

Since Σ_1 is homotopic to the closure S_1 of $D(\Lambda)/\Gamma$ in Σ sharing boundary components in M_1 and $\partial M_1 = \Sigma_1 \cup S_1$, it follows that M_1 is homeomorphic to $\Sigma_1 \times I$ for an interval I by Theorem 10.2 of [38].

Notice also

$$\partial\Sigma_1 = \partial S_1 = \bigcup_{i=1}^{\mathbf{b}} \mathbf{c}_i$$

for the number of boundary components \mathbf{b} of S_1 and the boundary components \mathbf{c}_i of Σ_1 .

We can do similarly for the closure $\hat{\mathcal{D}}_- := \text{Cl}_{\tilde{M}}(\mathcal{A}(\mathcal{D}))$ and we obtain $M_2 := \hat{\mathcal{D}}_-/\Gamma$ imbedded into M disjoint from M_1 . Similarly M_2 is homeomorphic to $\Sigma_2 \times I$ for a surface $\Sigma_2 = \text{bd}_M M_2$ which is also homotopy equivalent to M .

Notice also

$$\partial\Sigma_2 = \bigcup_{i=1}^{\mathbf{b}} \mathbf{c}'_i$$

for the boundary components \mathbf{c}'_i of Σ_2 .

Let $\text{int}M_1$ and $\text{int}M_2$ denote the topological interiors. Since we are removing the product of I with surfaces, it follows that $M - \text{int}M_1 - \text{int}M_2$ is a compact 3-manifold homeomorphic to M bounded by $\Sigma_1 \cup \Sigma_2 \cup B_1 \cup \dots \cup B_g$. (Recall B_i from Section 9.2.)

By geometric finiteness, every nonwandering geodesic is of a uniformly bounded δ -distance from a base point x_0 in M . Therefore, we can isotopy all B_i to imbedded annuli in M° so that the resulting imbedded annuli B'_i are all disjoint from closed spacelike geodesics in M° and $\partial B'_i \subset \Sigma_1 \cup \Sigma_2$. Therefore, Σ_1 and Σ_2 and B'_1, \dots, B'_g bound a compact submanifold \hat{M} containing all closed spacelike geodesics.

Assume now that $\mathcal{L}(\Gamma)$ is not a subset of $\text{SO}(2, 1)^\circ$. The full general case follows since a coset representative $\phi \in \Gamma - \Gamma'$ induces a diffeomorphism of M sending M_1 to M_2 and vice versa and by Lemma 9.4. We can also choose B_i equivariantly. \square

REFERENCES

- [1] J. Anderson, P. Bonfert-Taylor, and E. Taylor, Convergence groups, Hausdorff dimension, and a theorem of Sullivan and Tukia, *Geom. Dedicata* **103** (2004), 51–67.
- [2] H. Abels, G.A. Margulis, and G.A. Soifer, Properly discontinuous groups of affine transformations with orthogonal linear part, *C. R. Acad. Sci. Paris Sér. I Math.* **324**(3) (1997), 253–258.

- [3] T. Barbot, Globally hyperbolic flat space-times, *J. Geom. Phys.* **53**(2) (2005), 123–165.
- [4] A. Beardon, *Geometry of discrete groups*, Graduate Text in Mathematics Vol. **91** Springer New York, 1995.
- [5] Y. Benoist, Convexes divisibles I, in: Dani S.G., Prasad G. (eds.) *Algebraic groups and arithmetic*, Tata Inst. Fund. Res. Stud. Math. 17, Tata Institute of Fund. Research, 2004, pp. 339–374.
- [6] J.-P. Benzécri, Sur les variétés localement affines et localement projectives, *Bull Soc. Math. France* **88** (1960), 229–332.
- [7] F. Bonahon, Bouts des variétés hyperboliques de dimension 3, *Ann. of Math.* (2) **124** (1986), no. 1, 71–158.
- [8] F. Bonsante, Flat spacetimes with compact hyperbolic Cauchy surfaces, *J. Differential Geom.* **69**(3) (2005), 441–521.
- [9] B. Bowditch, Notes on tameness, *Enseign. Math.* (2) **56**(3-4) (2010), 229–285.
- [10] W. Browder, J. Levine, and G.R. Livesay, Finding a boundary for an open manifold, *Amer. J. Math.* **87** (1965), 1017–1028.
- [11] H. Busemann and P. Kelly, *Projective geometry and projective metrics*, Academic Press Inc., New York, N. Y., 1953. viii+332 pp.
- [12] V. Charette, T. Drumm, and W. Goldman, Affine deformations of the three-holed sphere, *Geom. Topol.* **14** (2010), 1355–1382.
- [13] V. Charette, T. Drumm, and W. Goldman, *Finite-sided deformation spaces of complete affine 3-manifolds*, arXiv:1107.2862v1.
- [14] V. Charette, W. Goldman, and C. Jones, Recurrent geodesics in flat Lorentzian manifolds, *Canad. Math. Bull.* **47**(3) (2004), 332–342.
- [15] S. Choi, The Margulis Lemma and the thick and thin decomposition for convex real projective surfaces, *Adv. Math.* **122** (1996), 150–191.
- [16] S. Choi, Convex decompositions of real projective surfaces. I: π -annuli and convexity, *J. Differential Geom.* **40**(1) (1994), 165–208.
- [17] S. Choi, Convex decompositions of real projective surfaces. II: Admissible decompositions, *J. Differential Geom.* **40**(2) (1994), 239–283.
- [18] S. Choi and W. Goldman, The classification of real projective structures on compact surfaces, *Bull. Amer. Math. Soc. (N.S.)* **34**(2) (1997), 161–171.
- [19] D. Cooper, D. Long, and M. Thistlethwaite, Computing varieties of representations of hyperbolic 3-manifolds into $SL(4, \mathbb{R})$, *Exp. Math.* **15** (2006), 291–305.
- [20] D. Cooper, D. Long, and M. Thistlethwaite, Flexing closed hyperbolic manifolds, *Geom. Topol.* **11** (2007), 2413–2440.
- [21] T. Drumm, Fundamental polyhedra for Margulis space-times, Doctoral dissertation, University of Maryland (1990); *Topology* **31**(4) (1992), 677–683.
- [22] T. Drumm, Linear holonomy of Margulis space-times, *J. Differential Geom.* **38** (1993), 679–691.
- [23] T. Drumm and W. Goldman, The Geometry of Crooked Planes, *Topology* **38**(2) (1999), 323–351.
- [24] T. Drumm and W. Goldman, The isospectrality of flat Lorentz 3-manifolds, *J. Differential Geom.* **58** (2001), 457–465.
- [25] P. Eberlein, Geodesic flows on negatively curved manifolds, I, *Ann. of Math.*, 2nd Ser. **95** (1972), 492–510.
- [26] D.B.A. Epstein and A. Marden, Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces in *Analytical and geometric aspects*

- of hyperbolic space* (Coventry/Durham, 1984), pp. 113–253, *London Math. Soc. Lecture Note Ser.*, 111, Cambridge Univ. Press, Cambridge, 1987.
- [27] C. Frances, The conformal boundary of Margulis space-times, *C.R. Acad. Sci. Paris, Ser. I* **336** (2003), 751–756.
 - [28] D. Fried, W. Goldman, and M. Hirsch, Affine manifolds with nilpotent holonomy, *Comment. Math. Helv.* **56**(4) (1981), 487–523.
 - [29] D. Fried and W. Goldman, Three-dimensional affine crystallographic groups, *Adv. Math.* **47**(1) (1983), 1–49.
 - [30] W. Goldman, Affine manifolds and projective geometry on surfaces, Senior Thesis, Princeton University, 1977.
 - [31] W. Goldman, Projective structures with Fuchsian holonomy, *J. Differential Geom.* **25**(3) (1987), 297–326.
 - [32] W. Goldman, Convex real projective structures on compact surfaces, *J. Differential Geom.* **31** (1990), 791–845.
 - [33] W. Goldman, Projective geometry of manifolds, Lecture notes from a 1988 course, <http://www2.math.umd.edu/~wmg/pgom.pdf> 1988.
 - [34] W. Goldman and F. Labourie, Geodesics in Margulis space times, *Ergod. Theory Dynam. Systems* **32** (2012), 643–651.
 - [35] W. Goldman, F. Labourie, and G. Margulis, Proper affine actions and geodesic flows of hyperbolic surfaces, *Ann. of Math.* (2) **170** (2009), 1051–1083.
 - [36] W. Goldman, G. Margulis, and Y. Minsky, Complete flat Lorentz 3-manifolds and laminations on hyperbolic surfaces, Preprint, May 29, 2012
 - [37] C. McA. Gordon and R.A. Litherland, Incompressible surfaces in branched coverings, In *Smith Conjecture* (eds. Morgan and Bass). Academic Press 1984, pp. 139–151.
 - [38] J. Hempel, *3-manifolds*, Princeton University Press, Princeton, 1976.
 - [39] S. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Encyclopedia of mathematics and its applications vol. 54, Cambridge University Press, Cambridge 1995.
 - [40] N. Kuiper, On convex locally-projective spaces, in *Convegno Internazionale di Geometria Differenziale (Italia, 1953)*, pp. 200–213. Edizioni Cremonese, Roma, 1954.
 - [41] G.A. Margulis, Complete affine locally flat manifolds with a free fundamental group, In: *Automorphic functions and number theory*, II, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 134, 1984, pp. 190–205.
 - [42] A. Marden, The geometry of finitely generated kleinian groups, *Ann. of Math.* (2) **99** (1974), 383–462.
 - [43] G. Mess, Lorentz space-times of constant curvature, *Geom. Dedicata* **126** (2007), 3–45.
 - [44] R. Meyers, End reductions, fundamental groups, and covering spaces of irreducible open 3-manifolds, *Geom. Topol.* **9** (2005), 971–990.
 - [45] T. Nagano and K. Yagi, The affine structures on the real two-torus: I, *Osaka J. of Math.* **11** (1974), 181–210.
 - [46] P. Scott, Compact submanifolds of 3-manifolds, *J. Lond. Math. Soc.* (2) **7** (1973), 246–250.
 - [47] L. Siebenmann, The obstruction to finding a boundary for an open manifold of dimension greater than five, Ph.D. thesis, Princeton University, 1965.

- [48] W. Thurston, Geometry and topology of 3-manifolds, Lecture notes. Princeton University, 1979 <http://www.msri.org/publications/books/gt3m/>.
- [49] P. Scott and T. Tucker, Some examples of exotic noncompact 3-manifolds, *Q. J. Math. Ser. (2)* **40**(160) (1989), 481–499.
- [50] T. Tucker, Non-compact 3-manifolds and the missing-boundary problem, *Topology* **13** (1974), 267–273.

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON 305-701, SOUTH KOREA,

E-mail address: schoi@math.kaist.ac.kr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK,
MD 20742-4015, USA

E-mail address: wmg@math.umd.edu